




ACCEPTED  
29 FEB 1980  
Insecticide,  
Fungicide, and  
Herbicide Act  
No. of 226-243

# Copper-Sevin



Combination Dust  
For Control of Certain Insects and Diseases On  
Tomatoes, Beans  
and other Vegetables

ACTIVE INGREDIENTS  
COPPER...  
SEVIN...  
INERT INGREDIENTS

KEEP OUT OF REACH OF CHILDREN  
**CAUTION**  
STATEMENT OF PRACTICAL TREATMENT  
IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention immediately.  
IF INHALED: Remove victim to fresh air and apply respiration if indicated.  
IF ON SKIN: Wash promptly with soap and water. Rinse thoroughly.  
IF IN EYES: Rinse eyes for at least 15 minutes with water and call a physician immediately.  
NOTE TO PHYSICIAN: CARBARYL is a moderate reversible cholinesterase inhibitor. ATROPINE is antidotal.  
SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 226-243 EPA Est. 226-KY-1  
SEVIN is the Trade Mark of U.C.C. for CARBARYL

**TOBACCO STATES CHEMICAL CO., INC.**  
LEXINGTON, KENTUCKY 40580

**PRECAUTIONARY STATEMENTS**

**HAZARD TO HUMAN AND DOMESTIC ANIMALS:** Caution Harmful if swallowed, inhaled, or absorbed through the skin. May cause irritation of eyes, nose, and throat. Avoid breathing of dust. Do not take internally. Avoid contact with skin, eyes, and clothing. Do not feed treated plants to dairy or meat animals.

**ENVIRONMENTAL HAZARD:** This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treated area. This pesticide is toxic to aquatic organisms. Keep out of lakes, streams, or ponds. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use Product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

Avoid contamination of feed and food stuffs. Do not reuse empty container. Wrap container and put in trash container.

226-243

**HOW AND WHEN TO APPLY:** TASCOCOPPER-SEVIN COMBINATION DUST should be applied as it comes from the package with a hand duster. For best results, a thorough, even application should be made to upper and lower leaf surfaces and stems. It is not necessary to discolor the foliage with dust; the proper rate, evenly applied is adequate for control.

**INSECT Control** is best obtained by application before or at first signs of insect infestation, but preferably before damage from feeding occurs. Repeat as needed at 7 to 10 day intervals.

For **DISEASE Control** it is necessary to have the dust on the plants before the diseases occur.

Follow the application rates listed for the crops below. Observe the time interval indicated between last application and harvest of crop.

**NOTE:** To avoid POSSIBLE INJURY on TENDER FOLIAGE, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days.

Crop	Rate Per Acre	Insects Controlled	Diseases Controlled	Harvest Interval
Beans	10 lbs.	Mexican bean beetles	Downey mildew.	Do not apply within One day of Harvest
	20 lbs.	Leafhoppers, Bean leaf beetles, Cucumber beetles, Western bean cutworms, Flea beetles, and Japanese beetles.	Angular leaf spot, and Anthracnose	
	20 to 25 lbs.	Corn earworms and Velvet bean caterpillars		
	40 lbs.	Lygus and Stink bugs		
Broccoli, Cabbage, & Cauliflower	20 lbs.	Flea beetles and Harlequin bugs.	Downey mildew, Leaf spot.	Do not apply within Three days of Harvest.
	40 lbs.	Armyworms, Corn earworms & Imported cabbage worms.		
Cucumbers and Squash	10 lbs.	Pickleworms, Melonworms, Cucumber beetles, Squash bug, & Flea beetles.	Downey mildew, Angular leaf spot, and Anthracnose.	Do not apply within One day of Harvest
Potatoes	10 lbs.	Colorado potato beetles.	Early and Late blights	Do not apply within One day of Harvest.
	20 lbs.	Leafhoppers and Flea beetles.		
	40 lbs.	Lygus bugs.		
Tomatoes, and Peppers.	20 lbs.	Tomato fruit worms, Fall army worms, Flea beetles and Leafhoppers.	Early and Late blights, Septoria leaf spot, Anthracnose, and Leaf mold	Do not apply within One day of Harvest
	40 lbs.	Tomato hornworms, European corn borers, and Stink bugs.		

Tobacco States Chemical Co., Inc. makes no warranty of merchantability, fitness for any purpose, or otherwise, express or implied, concerning this product or its use, which extend beyond the statements on this label.

BEST DOCUMENT AVAILABLE

24