

TOBACCO STATES CHEMICAL CO., INC.
LEXINGTON, KENTUCKY 40580

DURSBAN®

2 lb. INSECTICIDE

CONTROLS PESTS OF HOUSEHOLDS (COCKROACHES, ANTS, BROWN DOG TICKS, SILVERFISH, SPIDERS). **TURF** (CHINCH BUGS, SOD WEBWORMS, ANTS, BROWN DOG TICKS, CUTWORMS, EARWIGS, CRICKETS, GRASSHOPPERS, CERTAIN WHITE GRUBS, HYPERODES WEEVIL and **ORNAMENTAL PLANTS:** APHIDS, BAGWORMS, GRASSHOPPERS, LEAFHOPPERS, SPITTLE BUGS, MEALYBUGS, WHITE FLIES, EXPOSED THRIPS, MITES, CUTWORMS.

Also Recommended for Mosquito Control and Area Control of Brown Dog Ticks.

ONLY FOR SALE TO: Use and Storage by Serviceperson

To Be Applied Only By or Under the Supervision of Pest Control Operators, Public Health Organizations, Mosquito Abatement Districts and Other Trained Personnel Responsible for Insect Control Programs.

(USE DIRECTIONS Continued from Back Panel)

Pests of Lawns and Other Ornamental and Recreational Turf Grass Areas. Use DURSBAN 2 Lb. to control the pests listed in the following table by application at the recommended dosage, and in accordance with the directions given below. Dilute DURSBAN 2 Lb. in enough water to obtain complete and uniform coverage of pest-infested areas. Apply as a coarse spray using suitable application equipment. Thoroughly water immediately after treatment to wash the insecticide into the turf and plants as noted. For best results, the lawn should be moist at time of treatment.

NOTE: To help prevent infestation of buildings by ants, crickets, earwigs, millipedes, and silverfish (pillbugs), treat a band 6 to 10 feet wide around a building to the turf and also the building foundation to a height of 2 to 3 feet where pests are active and may find entrance. Use 3 fluid ounces of DURSBAN 2 Lb. per 100 gals. of water and apply as a coarse spray to thoroughly and uniformly wet the band area.

Pests of Ornamentals. Use DURSBAN 2 Lb. to treat flowers, shrubs, vines, shade and flowering trees and evergreens found to be infested with the pests listed in the following table. Dilute DURSBAN 2 Lb. with water according to directions given in the table and apply using suitable hand or power spray equipment in a manner to provide complete and uniform coverage. For best results apply a wetting spray to both upper and lower leaf surfaces and infested limb and trunk areas. Attempt to penetrate dense foliage but avoid overspraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 day intervals if needed.

NOTE: Do not use on azaleas, camelias, poinsettias, rose bushes, or variegated ivy because of possible injury to these plants.

DOSAGE and DILUTIONS for USE on ORNAMENTAL PLANTS

Pests Controlled	Amount of DURSBAN 2 Lb. in Water to Make		
	3 gallons	11 gallons	100 gallons
Aphids, Bagworms, Eastern caterpillar, Fall webworm, Grasshopper, Mites, Orange-striped oakworm, Spittlebugs, Whiteflies, Yellow-necked caterpillar	6.5 fl. oz.	1.6 fl. oz.	1 pt. 1
Cutworms, Leafhoppers, Mealybugs, Mimosa webworm, Redhumped caterpillar, Thrips (exposed)	10 fl. oz.	3.2 fl. oz.	1 qt. 11
Ash borer, Lilac borer, Scale crawlers, Oak leafminer, Pine needle, Tea, White peach	20 fl. oz.	6.4 fl. oz.	2 qts. 11


i. Equivalent to 4 ounces of chlorpyrifos per 100 gal.
ii. Equivalent to 8 ounces of chlorpyrifos per 100 gal.
iii. Equivalent to 1 pound of chlorpyrifos per 100 gal.

NOTE: Mosquitoes coming to rest on areas treated to control of turf and ornamental plants will be killed for varying periods of time after treatment depending on exposure of treated areas to weathering conditions.

Attention (Lawn and Ornamental Plant Pests): Keep children and pets off treated areas until spray has dried. Keep out of lawn and garden fishponds and other bodies of water.

Area Control of Brown Dog Ticks and Chiggers: Use DURSBAN 2 Lb. to control Brown Dog ticks and chiggers infesting non-cropland areas such as roadsides, footpaths and trails, picnic and camping sites, parks and other recreational areas where these pests are present and create a nuisance or public health problem. Apply DURSBAN 2 Lb. in water at the rate of 1 pint per acre (equivalent to 1 fl. oz. per 270 sq. ft.) using a hydraulic sprayer, mist applicator, backpack sprayer or other suitable hand or power sprayer equipment. Treat low-lying brushy areas, weeds, and ground surface with pests during the early season to obtain thorough coverage. Use 4 to 10 gal. of water per acre.

(USE DIRECTIONS Continued on Side Panel)



ACTIVE INGREDIENTS

Chlorpyrifos (O,O-dichloro-2,2,6,6-tetrachloro-2-pyridyl)phosphorothioate) 26.6%
Aromatic petroleum derivative solvent 14.9%

INERT INGREDIENTS

TOTAL 58.5%

Contains 1 pound of Chlorpyrifos per gallon
Trademark of The DOW CHEMICAL CO

AGRICULTURAL CHEMICAL

Do Not Ship with Foods, Feeds, or Clothing

KEEP OUT OF REACH OF CHILDREN

WARNING

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Do not induce vomiting. Call a physician and or transport to the hospital.
IF INHALED: Remove to fresh air. If poisoning symptoms appear, transport to hospital.
IF ON SKIN: Immediately flush skin with plenty of water while removing and contaminated clothing. Call physician. Wash clothing before reuse.
IF IN EYES: Irrigate with water for 15 minutes. Transport to hospital.

Read Complete Precautions on Side Panel

EPA Registration No. 226-247 EPA Establishment 226-AY 1

**Net Contents:
One Gallon (3.79 L)**



NOTICE: See label areas for the product conforms to its chemical description and is ready for use for the purposes stated on the label. The label also contains directions for use, and other important conditions of use. But neither this warranty nor any other warranty of MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, expressed or implied, extends to the use of this product contrary to the instructions or under abnormal conditions, or under conditions not reasonably foreseeable to retailer and buyer assumes the risk for such use.