

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL (10 Points)

Do not contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Pesticide Disposal: Pesticide, spray mixture or rinse water that cannot be used according to label directions must be disposed of according to applicable Federal, state or local procedures.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or dispose of in a sanitary landfill or by other approved state and local procedures.


MIXING DIRECTIONS

1. Add sufficient water to the mixing tank to allow proper agitation by pump or paddles.
2. Add other desired pesticides as listed mixing thoroughly prior to the addition of oil.
3. Add oil under agitation when 3/4 full, topping off with water to form a milky solution.
4. Maintain agitation until solution is used.
5. In small equipment lacking agitators. Use the following table to convert to small spray volumes.

STIR OR SHAKE DILUTED SPRAY FREQUENTLY DURING APPLICATION

Spray Dilution Conversion

Gallons Scale-X per 100 Gallon of Spray	0.5	0.7	1.0	1.3	1.6	2.0	3.0
Tablespoons Scale-X per Gallon of Spray	1 1/4	1 8/10	2 1/2	3 1/3	4 1/10	5 1/10	7 7/10



Timing the Treatment: See tables below for general use. You must determine final timing to fit local growth and climatic conditions.

DO NOT EXCEED MAXIMUM RATES OR APPLY LATER THAN RECOMMENDED

CROP	PEST	GALLONS SCALE-X PER 100 GALLONS SPRAY	TIME OF APPLICATION (STAGE OF DEVELOPMENT)
APPLES	Apple Red Bug, European	2	Dormant
	Fruit Lecanium Scale		
	Fruit Tree Leafroller	3	Dormant
	European Red Mite	2	Green Tip to Delayed Dormant 1/2" Green.
	San Jose Scale	2	Delayed Dormant, 1/2" Green.
	Scurfy Scale	3	Delayed Dormant, 1/2" Green.
PEARS	Fruit Tree Leafroller	3	Dormant
	Pear Blister Mite		
	Pear Psylla	2	Dormant
PECANS	Obscure Scale	3	Dormant
PEACHES	Colony Peach Scale	3	Dormant
PLUMS & PEARNS	European Red Mite	2	Dormant to Delayed Dormant
	European Fruit Lecanium Scale		1/2" Green

CITRUS

ON CITRUS: Do not apply when trees are in or near wilting. Oil applied during August and September may affect solids content and retard color. Oil applied after October 1 may increase cold damage to trees. Observe all cautions and limitations on labels of all products used in mixtures. Parathion and Chlorobenzilate are "Restricted Use Pesticide" products. Do not use Scale-X in combination with or immediately preceding or following applications of Dinitro compounds, Sulfurs, Captan, Folpet, Dyrene, Karathane, Morestan, or Sevin. Check tank mixtures for chemical and physical compatibility.

CITRUS PEST	GALLONS OF SCALE-X PER 100 GALLONS OF DILUTE SPRAY	PESTICIDE	POST BLOOM
Scales: Glover (long), Chaff Purple, Red, Yellow, Snow, Brown Soft Scale, California Red Scale	1.0-1.3 1.3-1.6		March & April
Scales listed above & Rust Mite Greasy Spot & Sooty Mold (Fla. Only) California Red Scale	1.0-1.3 1.0-1.3 1.3-1.6	1 Lb Zineb 75% WP or 1/2 pt Chlorobenzilate 4	March & April
Scales listed above & Spider Mites, Black Scale, White Fly, California Red Scale (Do not use in Calif.)	0.5-0.7	1/2 Pt Ethion 4% 1/2 Pt Parathion 4% 1 Pt Malathion 5%	Summer (June- July 15)
Scale Insects in post bloom for oil, Greasy Spot & Sooty Mold (Fla. Only)	1.0-1.3		Summer (June- July 15)

* Add Zineb or Chlorobenzilate as listed for Rust Mite. Ethion - Do not repeat applications within 30 days. Parathion - Do not apply within 30 days of Harvest. Malathion - Do not apply within 7 days of Harvest.

Tobacco States Chemical Co., Inc. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PURPOSE, OR OTHERWISE, EXPRESS OR IMPLIED, concerning this product or its use, which extend beyond the statement on this label.