## WARNING

May cause skin irritation. Harmful if swallowed. Do not take internally. Avoid inhaling spray mist. Avoid contact with skin, eyes or clothing.

This product will kill or seriously injure many desirable forms of vegetation. Do not apply directly to flowers, fruits, vegetables, grapes, ornamentals, cotton or other desirable plants. Do not use when there is hazard from drifting mists. (Coarse sprays are less likely to drift.)

Vapors from this product may injure susceptible plants in the immediate vicinity.

Avoid contamination of water used for domestic purposes and irrigation ditches. Excessive amounts of this product in the soil may temporarily inhibit seed germination and plant growth.

This product not for use on lawns.

This product is toxic to fish. Keep out of any body of water. Apply this product only as specified on this label.

Do not graze dairy animals on treated areas for 7 days after treatment.

Do not store near fertilizers, seeds, insecticides or fungicides.

Spray equipment used in applying this product should be thoroughly cleaned before using for any other purpose. Use repeated flushing with soap and warm water or suitable chemical cleaner. Rinse equipment and containers and dispose of wastes or spillage by burying in noncrop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes. It is best to use a separate sprayer for application of insecticides and fungicides.



Contains 4 Lbs. 2,4,5-T Esters

ACTIVE INGREDIENTS: \*Butyl Ester 2,4,5-Trichlorophenoxyacetic Acid \*\*Isooctyl Ester 2,4,5-Trichlorophonoxyacetic Acid INERT INGREDIENTS

TOTAL \*Equivalent to 25.4% 2,4,5-Trichlorophenoxyacetic Acid

\*\*Equivalent to 9.6% Trichlorophenoxyacetic Acid > Contains 3.14 lbs. 2,4,5-T Acid Per Gallon

## CAUTION: **KEEP OUT OF REACH OF CHILDREN**

## DIRECTIONS FOR USE

Use in non-crop areas as rangeland, pasture, fence rows, drainage ditch banks, around buildings, along roadways, power lines, right of ways, and other similar waste places.


Use 2-3 guarts WEEDESTROY 2,4,5-T in 100 gallons of water. Apply spray when plants are in full leaf and actively growing. Thoroughly wet all foliage and bark. The actual volume needed will depend on the density and size of the brush. Best results are obtained by using high pressures and applying a minimum of 100 gallons of spray per acre. Repeat applications may be necessary to control hard to kill species.

For knapsack sprayers dilute 1/2 to 1 pint WEEDESTROY 2,4,5-T in 5 gallons of water and apply as a drenching spray. These lower concentrations have been used successfully on the most susceptible woody plants like sassafras, cherry, and willow. Brush is best controlled when less than six feet high. Brush tailer than six feet is better controlled if cut and the stubble sprayed thoroughly with a solution of 2-3 quarts of WEEDESTROY 2,4,5-T in 10 gallons of kerosene or fuel oil.

STUMP TREATMENT - Stump treatment at any time of the year is now possiblbe with WEEDESTROY 2,4,5-T. Use 2-3 pints in 10 gallons Diesel oil or kerosene and apply with knapsack sprayer equipped with a fan type nozzle. Spray tops and sides of stumps thoroughly. This treatment has reduced resprouting as much as 75% using a single application. Retreat the few resprouting stumps the following spring when sprouts are in full leaf.

DORMANT SPRAY - Use 2 quarts in 10 gallons fuel, diesel, kerosene or transformer oil. Spray trunks and brush from ground line to 3 ft. Apply spray to run-off point completely circling trunk. In large trees axe incisions are helpful. CHICCEDTIDI E DI ANITE

| SUSCEPTIBLE P<br>Alder<br>Aspen<br>American<br>Crab Apple<br>Barberry<br>Birch<br>Blackberry<br>Buckbrush<br>Catalpa<br>RESISTANT PLA | Cedar, salt<br>Cherry, choke<br>Cherry, wild<br>Chinaberry<br>Cottonwood<br>Currant<br>Dogwood<br>Elderberry<br>Elder, box | Elm<br>Green briar<br>Gum, black<br>Gum. Sweet<br>Hackberry<br>Hawthorne<br>Hazel<br>Hickory<br>Honeysuckle | Herculesclub<br>Hemlock<br>Hornbeam<br>Horsechestnut<br>Juneberry<br>Juniper<br>Locust, black<br>Locust, honey<br>Mulberry | Oak<br>Osage, Orange<br>Pine<br>Poison ivy<br>Poison, oak<br>Poplar<br>Prickly Ash<br>Raspberry<br>Rose Cherokee | Sagebrush<br>Sassafras<br>Silverberry<br>Skunkbrush<br>Shadbush<br>Snowberry<br>Spicebush<br>Sumac<br>Sweetfern | Sycamore<br>Tree of Heaven<br>Trumpet vine<br>Tuliptree<br>Virginia creeper<br>Walnut<br>Willow<br>Winterberry<br>Witchhazel |
|---|--|---|--|--|---|--|
|   | sswood Beech   | Maple, F  | Red Mesquite   | Persimmon  | Rhododendron  | Spruce   |



iodenaron Spruce

## NOTICE

The use of this material being subject to conditions beyond their control, neither Riverdale Semical Company or Seller liable for the resepresentation or warranty, express or implied, witherespect to results from such use, whether print used in accordance with directions. The We accepts and uses this material subject to his terms and shall not hold either Riverdale Elemical Company or seller liable for the refulls of any use of this material. Any buyer not accepting the foregoing terms is requested to return to seller this container unopened and in good condition within five (5) days from the date of purchase. Seller is hereby authorized by Riverdale Chemical Company to refund the purchase price.

Any buyer who shall not have returned this container in such time and manner shall be conclusively presumed to have accepted all terms of this notice.

NET CONTENTS

GAL.

U.S.D.A. Reg. No. 228-16

MANNE FAC THIRE C. S. RIVERDALE CHEMICAL COMPANY CHICAGO HEIGHTS ILLINOIS