

228-126

04/29/2004

1/14

RIVERDALE®

SOLUTION™ EMULSIBLE

A SELECTIVE WEED KILLER

FOR CONTROL OF MANY BROADLEAF WEEDS AND BRUSH CONTROL IN NON-CROP AREAS SUCH AS LAWNS, PASTURES, RANGELANDS, FENCE ROWS, RIGHTS-OF-WAY, ALSO FOR USE IN CORN AND SMALL GRAINS.

ACTIVE INGREDIENT:

Isooctyl (2-ethylhexyl) Ester of 2,4-Dichlorophenoxyacetic Acid\* . . . . . 94.0%

INERT INGREDIENTS:\*\* . . . . . 6.0%

TOTAL . . . . . 100.0%

Isomer Specific AOAC Method, Equivalent to:

\*2,4-Dichlorophenoxyacetic Acid . . . . . 62.4%, 5.9 lbs./gal.

\*\*This product contains Petroleum Distillates

Riverdale is a Registered Trademark of Riverdale - A Nufarm Company

Solution is a Trademark of Riverdale - A Nufarm Company

KEEP OUT OF REACH OF CHILDREN

CAUTION - CAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

(If you do not understand the label, find someone to explain it to you in detail.)

SEE SIDE PANELS FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

NET CONTENTS

GALS.

EPA REG. NO. 228-126

EPA EST. NO. 228-IL-1

MANUFACTURED BY NUFARM AMERICAS, INC., BARR RIDGE, ILLINOIS 60527-0866

ACCEPTED

APR 29 2004

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

228-126

NOTE: Spanish Language is optional.

Revised A/O 8/8/02 More 2-EHE/GDE Revisions.

A/O 5/15/03 Changed First Aid per PR 2001-1 and switched competitor product names for five of our new product names via Amendment. Updated 9/19/03

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION - CAUCION**

Harmful if swallowed, inhaled or absorbed through skin. Avoid inhalation of vapors or spray mist, and contact with skin, eyes and clothing. Remove saturated clothing as soon as possible and shower. If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron. If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

**NON-WPS TURF USES:** Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) -- in general, only agricultural plant uses are covered by the WPS -- must wear: long pants, long-sleeved shirt, socks, shoes and chemical resistant gloves. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications to turf per treatment site is 2 per year.

**NON-WPS INDUSTRIAL USES:** When mixing, loading or applying this product or repairing or cleaning equipment used with this product, wear face shield, goggles or safety glasses and chemical resistant gloves, long-sleeved shirt, long pants, socks and shoes. It is recommended that safety glasses include front, brow and temple protection. For aerial applicators in an enclosed cockpit and applicators applying this product from a tractor that has a completely enclosed cab, eye protection is not required. Wash hands, face and arms with soap and water as soon as possible after mixing, loading or applying this product. After work, remove all clothing and shower using soap and water. Do not reuse clothing worn during the previous day's mixing and loading or application of this product without cleaning first. Clothing must be kept and washed separately from other household laundry.

**WPS USES:** Personal Protective Equipment - Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) -- in general, agricultural plant uses are covered -- must wear: long-sleeved shirt and long pants; chemical-resistant gloves such as Barrier Laminate, Nitrile Rubber ≥ 14 mils, Neoprene Rubber ≥ 14 mils, and Viton ≥ 14 mils; shoes plus socks; and protective eyewear. If this container is over one gallon and less than five gallons, mixers and loaders who do not use a mechanical system (such as a probe and pump or spigot) to transfer contents of this container must wear coveralls or a chemical resistant apron in addition to the other required PPE. Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry. After each day of use, clothing or PPE must not be reused until it has been cleaned.

**Engineering Controls Statements:** If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal. If the mechanical system is used in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

**USERS SHOULD:** Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**FIRST AID STATEMENT**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

**ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Vapors from this product may injure susceptible plants in the immediate vicinity. Use care to avoid spray contact or drift to 2,4-D susceptible plants such as cotton, tomatoes, flowers, okra, grapes, fruit trees and ornamentals. Do not permit spray mist containing this product to drift onto them. Do not spray when the wind is blowing towards susceptible crops or ornamental plants. Use coarse sprays and/or low spray pressure to minimize drift. Do not apply with hollow cone type insecticide or other nozzles that produce fine spray droplets. Spray drift can be lessened by keeping the spray boom as low as possible by spraying when wind velocity is low, by decreasing the pounds of pressure of the nozzle tips, and by stopping all spraying when wind exceeds 6 to 7 miles per hour. On cropland and along roadsides, do not exceed 20 psi pressure. Do not apply when a temperature inversion exists. If questions exist pertaining to the existence of an inversion, consult with local weather services before making an application. Do not use the same spray equipment for applying other materials to 2,4-D susceptible crops as injury may result. It is best to use a separate sprayer for application of insecticides and fungicides. Clean and rinse spray equipment using soap or detergent and water or suitable chemical cleaner, and rinse thoroughly before reuse for other spraying. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate domestic or irrigation waters. However, treated water may be used for watering turf grasses immediately after application. Do not use in or near a greenhouse. Excessive amounts of this product in the soil may temporarily inhibit seed germination and plant growth.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

When using on Pastures and Rangeland Grasses there is a (1) 7 day pre-grazing interval for dairy cattle; (2) 30 day preharvest interval for grass cut for hay; and (3) 3 day pre-slaughter interval for meat animals.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls, chemical-resistant gloves such as Barrier Laminate, Nitrile Rubber  $\geq$  14 mils, Neoprene Rubber  $\geq$  14 mils, and Viton  $\geq$  14 mils; shoes plus socks, and protective eyewear. No Worker Protection Standard worker entry restrictions or worker notification requirements apply when this product is directly injected into agricultural plants.

#### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

For Turf use do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

#### GENERAL INFORMATION

This product is a low volatile ester especially prepared for use on crops and weeds where a susceptible crop in the near vicinity may be injured by a more volatile product. It is recommended for control of numerous broadleaf weeds and certain 2,4-D susceptible woody plants without injury to most established grasses. In cropland, 2,4-D Ester is more effective than amines for controlling hard-to-kill weeds such as bindweed, thistle, smartweeds, wild garlic, curly dock, tansy ragwort, and wild onions. For best results apply this product as a water or oil spray during warm weather when young succulent weeds or brush are actively growing. Application under drought conditions often will give poor results. The lower recommended rates will be satisfactory on susceptible annual weeds. For perennial weeds and conditions such as the very dry areas of the western states, where control is difficult, the higher recommended rates should be used. Deep-rooted perennial

weeds such as Canada thistle and field bindweed and many woody plants usually require repeated applications for maximum control.

Unless otherwise recommended, suggested application rates may be up to 10 gallons of total spray by air or 5 to 25 gallons by ground application equipment. If band treatment is used, base the dosage rate on the actual area to be sprayed. Although water quantities may vary due to different types of application equipment, sufficient water must be used to provide for complete and uniform coverage. High water gallonage may be used if desired to improve spray coverage. In all cases, use the same recommended amount of 2,4-D per acre. When product is used for weed control in crops, the growth stage of the crop must be considered. For crop uses, do not mix with oil, surfactants, or other adjuvants unless specifically recommended on label. To do so may reduce herbicide's selectivity and could result in crop damage. If you are not prepared to accept some degree of crop injury, do not use this product. Crop varieties vary in response to 2,4-D and some are easily injured. Apply this product to varieties known to be tolerant to 2,4-D. If you are uncertain concerning tolerant varieties of local use situations that may affect crop tolerance to 2,4-D, consult your seed company, State Agricultural Extension Service or qualified crop consultant for advise.

Aerial applications should be used only when there is no danger of drift to susceptible crops. Many states have regulations concerning aerial application of 2,4-D formulations. Consult local regulatory authorities before making applications. Although this product is a low volatile formulation, at temperatures above 90°F vapors may damage susceptible crops growing nearby.

**TO PREPARE THE SPRAY:** (1) Fill the spray tank about half full with water, then add the required amount of this product with agitation, and finally the rest of the water. NOTE: This product in water forms an emulsion which tends to separate unless the mixture is kept agitated. Continue agitation during application until spray tank is empty. (2) If oil is added, first mix this product and the oil and then add this mixture to the water. However, with adequate agitation, the oil can be added after the product is mixed in water. (3) If straight oil is used, a solution is formed and separation does not occur. Do not allow any water to get into the oil-herbicide mixture to avoid formation of an invert emulsion.

Riverdale Solution™ Emulsible will kill or control the following weeds in addition to many other noxious plants susceptible to 2,4-D.

Alder, Alfalfa, Artichoke, Aster, Austrian fieldcress, Beggartick, Bident, Bindweed, Bitterweed, Bitter wintercress, Blackeyed Susan, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckbrush, Buckhorn, Bull thistle, Bur ragweed, Burdock, Burhead, Buttercup, Canada thistle, Carpetweed, Catnip, Chamise, Cherokee rose, Chickweed, Chicory, Cinquefoil, Coastal redstem sage, Cockle, Cocklebur, Coffee bean, Coffeeweed, Common sowthistle, Cornflower, Coyotebrush, Creeping jenny, Croton, Curly Indigo, Dandelion, Devil's claw, Dock, Dogbane, Dogfennel, Elderberry, Fanweed, Fiddle neck, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Goosefoot, Ground ivy, Gumweed, Halogeton, Hawkweed, Healall, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Indiana mallow, Indigo, Ironweed, Jerusalem artichoke, Jewelweed, Jimsonweed, Klamathweed, Knotweed, Kochia, Lambsquarter, Locoweed, Lupine, Mallow, Manzanita, Marijuana, Many flowered aster, Marshelder, Mexican weed, Milkvetch, Morningglory, Musk thistle, Mustard, Nettle, Nutgrass, Orange hawkweed, Parsnip, Pennycress, Penneywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Rabbitbrush, Ragweed, Redstem, Russian thistle, Sagebrush, Salsify, Sand shinnery oak, Shepherdspurse, Sicklepod, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, St. Johnswort, Starthistle, Stinging nettle, Stinkweed, Sumac, Sunflower, Sweet clover, Tansymustard, Tansyragwort, Tanweed, Tarweed, Texas blueweed, Thistles, Toadflax, Tumbleweed, Velvetleaf, Vervain, Vetch, Virginia creeper, Wild buckwheat, Wild carrot, Wild garlic, Wild lettuce, Wild onion, Wild parsnip, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Wormseed, Wormwood, Yellow rocket, Yellow starthistle and other broadleaf weeds which may be listed elsewhere on this label.

Some of these species may require repeat applications and/or use of higher rate recommended on this product label even under ideal conditions for applications. Control of pigweeds in the High Plains area of Texas and Oklahoma may not be satisfactory with this product.

**SELECTIVE WEEDING IN CROPS**

**USE IN LIQUID NITROGEN FERTILIZER:** This product may be combined with liquid nitrogen fertilizer suitable for foliage application on corn, grass, pastures, or small grains in one operation. Use product according to directions on this label for those crops. Use liquid nitrogen fertilizer at rates recommended by supplier or Extension Service Specialist. Mix the product and fertilizer according to the following instructions: Fill the spray tank approximately half full with the liquid nitrogen fertilizer. Add the product while agitating the tank. Add the remainder of the fertilizer while continuing to agitate. Apply immediately maintaining agitation during application until tank is empty. **Do not apply during cold (near freezing) weather.** Spray mixture must be used immediately and may not be stored. Do not allow mixture to stand overnight.

**NOTE:** If good, continuous agitation is not maintained, separation of the spray mixture and/or clogging of the nozzles is likely to occur. Fertilizers can increase foliage contact burn of herbicides. Reducing the fertilizer rate and concentrate will reduce the hazard of leaf burn.

**CORN (Field, Sweet and Popcorn):**

Pre-plant -	2/3 to 1-1/3 pints
Pre-emergent -	Average Conditions - 1-1/4 to 2-1/4 pints
Emergent -	2/3 pint
Post-emergent -	Average Conditions - 1/4 pint Dry Conditions* - 1/4 to 1/2 pint
Pre-harvest -	1/2 to 1-1/3 pints

\*For western states - Arizona, Idaho, Montana, Nevada, Oregon, Utah, Washington and Wyoming

All with recommended amounts of water to make per acre applications. Use lower rates of product for easily-killed weeds, on inbreds, and when corn is growing rapidly. Do not cultivate for about 2 weeks after treatment while corn is brittle.

**Pre-Plant:** To control emerged broadleaf weed seedlings or existing cover crops prior to planting corn. Apply 7 to 14 days before planting. Do not use on light, sandy soil, or where soil moisture is inadequate for normal weed growth. Use high rate for control of less susceptible weeds or cover crops such as alfalfa.

**Pre-emergent:** Apply product to emerged weeds from 3 to 5 days after planting but before corn emerges. Do not use on very light, sandy soils. Use the higher rates on heavy soils. Plant corn as deep as practical. Product will not control weeds which have not emerged.

**Emergent:** Apply in 10 to 30 gallons of water per acre just as corn plants are breaking ground.

**Post-emergent:** As soon as corn is over 8 inches tall, use drop nozzles to keep spray off corn foliage as much as possible; direct spray over tops of weeds but not over the corn. Do not apply from tasseling to dough stage. If corn is growing rapidly and temperature and soil moisture is high, use 1/4 pint per acre to reduce possibility of crop damage. Delay cultivation for 8 to 10 days to prevent stalk breakage due to temporary brittleness caused by 2,4-D. Application rates of up to 1/2 pint per acre may be used to control some hard-to-control weeds. However, the possibility of injury to the corn is increased. Do not use with atrazine, oil or other adjuvants. Since the tolerance to 2,4-D of individual hybrids varies, consult your seed supplier, local Extension Service, Agricultural Experiment Station, or University Weed Specialist for information.

**Pre-harvest:** After the hard dough or denting stage, apply 1/2 to 1-1/3 pints in 20 to 50 gallons of water per acre by air or ground equipment to suppress perennial weeds, decrease weed seed production, and control tall weeds such as bindweed, cocklebur, dogbane, jimsonweed, ragweed, sunflower, velvetleaf and vines that interfere with harvesting. The high rate will be needed for tough weeds under stress.

**SMALL GRAINS (barley, oats, wheat, rye), not underseeded with a legumes:**

Wheat, Barley, Rye - Annual weeds -	Average Conditions - 1/4 to 1/2 pint;
	Dry Conditions (Western States) - 1/2 to 1/4 pints
Perennial weeds -	Average Conditions - 1/2 pint;
	Dry Conditions (Western States) - 3/4 - 1-1/4 pints
Pre-harvest -	Average Conditions - 1/2 to 1-1/4 pints

Oats --- Spring - 1/4 pint and Fall - 1/4 to 1/3 pint

For aerial application on grain, it is suggested to use this product in 2 or more gallons of water per acre and for ground application use a minimum of 10 gallons of water per

acre.

Make application in the Spring when the grain is fully tillered or stooled (usually about 4 to 8 inches high), but before jointing. Do not spray before the tiller stage nor from early boot to dough stage.

Use lower rate of product for easily-killed seedling weeds, and higher rate for older and more tolerant weeds. Do not treat grains under-seeded with legumes, and do not spray winter grains in the Fall. To control large weeds that will interfere with harvest or to suppress perennial weeds, pre-harvest treatment can be applied when grain is in the dough stage. Higher rates may be needed to handle difficult weed problems in certain areas such as under dry conditions especially in western areas. However, do not use unless possible crop injury will be acceptable. For the high rates on Spring wheat and barley as well as winter wheat and rye, consult State Agricultural Experiment Station or Extension Service weed specialists for recommendations or suggestions to fit local conditions.

**For emergency weed control in wheat:** Perennial broadleaf weeds - apply 1-3/4 pints per acre when weeds are approaching bud stage. Do not spray grain in the boot to dough stage. The 1-3/4 pint per acre application can produce injury to wheat. Balance the severity of your weed problem against the possibility of crop damage. Where perennial weeds are scattered, spot treatment is suggested to minimize the extent of crop injury. Use lower rate if small annual and biennial weeds are the major problem. Use the higher rate if perennial weeds or annual and biennial weeds are present which are in the hard-to-kill categories as determined by local experience. The higher rates increase the risk of grain injury and should be used only where the weed control problem justifies the grain damage risk. Do not apply Solution Emulsible to grain in the seedling stage. For aerial application on grain, apply this product in 2 to 10 gallons of water per acre. For ground application use a minimum of 10 gallons of water per acre.

**Spring Seeded Oats:** Use 1/4 pint per acre with recommended amount of water to give good coverage. Apply after the fully tillered stage, except during the boot to dough stage.

**Fall Seeded Oats (Southern):** Apply 1/6 to 3/4 pint per acre with recommended amount of water after full tillering but before early boot stage. Some difficult weeds may require the higher rates of 1/3 to 3/4 pint per acre for maximum control but injury may result. Do not spray during or immediately following cold weather.

**Pre-harvest Treatment:** Apply 2/3 to 1-1/3 pints with recommended amount of water per acre when grains are in the hard dough stage to control large weeds that may interfere with harvest. Best results will be obtained when soil moisture is sufficient to cause succulent weed growth.

NOTE: Oats are less tolerant to 2,4-D than wheat or barley and more likely to be injured.

**Wheat and Barley:** Control of Wild Garlic and Wild Onion.

For improved control of difficult weeds including Wild Garlic and Wild Onion, apply 1/2 to 1-1/4 pints of product per acre. Since these rates may injure the crop do not use unless possible crop damage is acceptable. For the higher rates on Spring wheat and barley, consult your local State Agricultural Experiment Station or Extension Service Weed Specialist for recommendations or suggestions to fit local conditions.

**Control of Wild Garlic in Stubble Grain and Corn Fields:** Following the harvest of small grains, and corn, Wild Garlic often produces new Fall growth. This should be sprayed with 2-1/2 to 3-3/4 pints of product in 20 to 40 gallons of water per acre. This is a useful practice as one part of Wild Garlic control program. Do not plant any crop for three months after treatment.

**SORGHUM (Milo):** For Post-emergent control in average conditions, use 1/4 pint; dry conditions (Western States) use 1/4 to 1/3 pint with suggested volume of 5 gallons of water by air or 6 to 20 gallons with ground equipment to make per acre applications. Apply to sorghum when crop is 5 to 15 inches high to top of canopy with secondary roots well established. If sorghum is taller than 8 inches, use drop nozzles to keep the spray off the foliage as much as possible. Do not apply boot, flowering or early dough stage. Rates of up to 1/2 pint per acre may be used to control some hard-to-control weeds. However, the chance of crop injury is increased with the higher rates. Do not use with oil. Because temporary injury may occur if conditions of high temperature and high soil moisture exist, use lower rate. Varieties vary in tolerance to 2,4-D and some hybrids are quite sensitive. Spray only varieties known to be tolerant to 2,4-D. Contact seed company or your Agricultural Experiment Station or Extension Service weed specialists for this information.

**GRASS SEED CROPS:** Apply 1/2 to 2-1/2 pints of product in up to 30 gallons of water per acre by air or ground equipment in the Spring or Fall to control broadleaf weeds in grass being grown for seed. Do not apply from early boot to milk stage. Spray seedling grass only after the five leaf stage, using 1/3 to 1/2 pint per acre to control small seedling weeds.

After the grass is well established, higher rates of up to 2-1/2 pints per acre can be used to control hard-to-control annual or perennial weeds. For best results, apply when soil moisture is adequate for good growth. Do not use on Bent unless injury can be tolerated.

**NO-TILL APPLICATION:** Solution™ Emulsible may be used in the broadcast method with a normal boom or with direct pipes set 12" apart in 36" rows. When using Solution Emulsible, apply at a rate of 9 oz. in 10 gallons of water per acre. Maintain uniform pressure and speed when applying.

**FOR USE IN CROP RESIDUE MANAGEMENT SYSTEMS  
IN SOYBEANS (Preplant only)**

**GENERAL INFORMATION:** Riverdale SOLUTION EMULSIBLE is a herbicide that provides control of many emerged susceptible annual and perennial broadleaf weeds. River SOLUTION EMULSIBLE may be applied prior to planting soybeans to provide foliar burndown control of susceptible annual and perennial broadleaf weeds and certain broadleaf cover crops such as those listed on this label. Riverdale SOLUTION EMULSIBLE should only be applied preplant to soybeans in situations such as reduced tillage production systems, where emerged weeds are present. Apply only according to the application instructions given below. Do not use any tillage operations between application of Riverdale SOLUTION EMULSIBLE and planting soybeans.

**MIXING INSTRUCTIONS:** Compatible crop oil concentrates, agricultural surfactants and fluid fertilizers approved for use on growing crops may increase the herbicidal effectiveness of 2,4-D on certain weeds and may be added to the spray tank. Read and follow all directions and precautions on this label and on all labels of adjuvants or fertilizers mixed with this product.

**APPLICATION PROCEDURES:** Apply using air or ground equipment in sufficient gallonage to obtain adequate coverage of weeds. Use 2 or more gallons of water per acre in aerial equipment and 10 or more gallons of water per acre in ground equipment.

**APPLICATION TIMING AND USE RATES**

2,4-D Formulation	Maximum Rate	When to Apply (Days prior to planting soybeans)
SOLUTION™ EMULSIBLE	5/8 pint (10.0 fl. oz.) (0.5 lb. a.e./acre) 1-1/4 pint (20.0 fl. oz.) (1.0 lb. a.e./acre)	NOT LESS THAN 7 DAYS  NOT LESS THAN 30 DAYS

**WEEDS CONTROLLED:** Alfalfa\*, Bindweed\*, Bullnettle, Bittercress-smallflowered, Buttercup-smallflowered, Carolina geranium, Cinquefoil-common and rough, Clover-red\*, Cocklebur-common, Dandelion\*, Eveningprimrose-cutleaf, Garlic-wild\*, Horseweed or marestail, Ironweed, Lambsquarters-common, Lettuce-prickly, morningglory-annual, Mousetail, Mustard-wild, Onion-wild\*, Pennycress-field, Peppergrass\*, Purslane-common, Ragweed-common, Ragweed-giant, Shepherdspurse, Smartweed-Pennsylvania\*, Sowthistle-annual, Speedwell, Thistle-Canada\*, Thistle-bull, Velvetleaf, Vetch-hairy\*, Virginia copperleaf.

\*These species are only partially controlled.

For best weed control at time of treatment, weeds should be small, actively growing and free of stress caused by extremes in climatic conditions, diseases, or insect damage. The response of individual weed species to Riverdale SOLUTION EMULSIBLE is variable. Consult your local county or state AGRICULTURAL Extension Service or crop consultant for advice.

**APPLICATION RESTRICTIONS AND PRECAUTIONS:** Important Notice - Unacceptable injury to soybeans planted in fields treated with Riverdale SOLUTION EMULSIBLE may occur. Whether or not soybean injury occurs and the extent of the injury will depend on weather (temperature and rainfall) from herbicide application until soybean emergence and agronomic factors such as the amount of weed vegetation and previous crop residue present. Injury is more likely under cool rainy conditions and where there is less weed vegetation and crop residue present.

Do not use on low organic sandy soils ( $\leq 1.0\%$ ).

Apply maximum of one application per growing season regardless of the treatment rate.

Do not apply Riverdale SOLUTION EMULSIBLE when weather conditions such as temperature air inversions or wind favor drift from treated areas to susceptible plants.

**Livestock Grazing Restriction:** Do not feed hay, forage or fodder. Restrict livestock from grazing treated fields. Livestock should be restricted from feed/grazing of treated cover crops.

In fields treated with Riverdale SOLUTION EMULSIBLE, plant soybean seed as deep as practical or at least 1.0 inch deep. Adjust the planter, if necessary, to ensure that planted seed is completely covered.

Do not apply Riverdale SOLUTION EMULSIBLE prior to planting soybeans, if you are not prepared to accept the results of soybean injury, including possible loss of stand and yield.

Do not replant fields treated with Riverdale SOLUTION EMULSIBLE in the same growing season with crops other than those labeled for 2,4-D use.

#### **SELECTIVE WEEDING IN NON-CROP AREAS**

**ORNAMENTAL TURF** such as Lawns, Golf Courses (Fairways, Aprons, Tees and Roughs), Sod Farms, Cemeteries, and Parks:

Use 1.25 to 2.7 pints of product in 40 to 180 gallons of water to give good coverage to one acre on established stands of perennial grasses. Usually 2.5 pints per acre provides good weed control under average conditions. On turf, apply a maximum of 2.7 pints of this product per acre per application per site. Treat when weeds are young and actively growing. Do not apply to newly seeded grasses until well established. Use higher rate for hard-to-kill weeds. Use higher rate when using higher volume of water per acre. Do not exceed specified application dosages for any area. Deep-rooted perennial weeds may require repeated treatments in the same season or in subsequent years. Spray when air temperature is between 50° and 85°F. Avoid applying during excessively dry or hot periods unless irrigation (watering) is used before treatment. Do not apply if rainfall is expected within 48 hours, nor should lawns be irrigated for 48 hours following application. For optimum results, turf should not be mowed for 1 to 2 days before and after application. Reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Maximum kill of weeds will be obtained by applying in Spring and early Fall when weeds are actively growing. Do not use on golf greens nor on dichondra or other broadleaf herbaceous ground covers. Do not use on creeping grasses such as bent and St. Augustine except for spot treating, nor on newly seeded turf until grass is well established.

**FALLOW LAND:** Use 1-1/4 to 3-3/4 pints of this product in a recommended minimum of 10 gallons of water per acre for ground application and recommended minimum of 2 gallons for aerial application of water per acre on annual broadleaf weeds and up to 4 pints per acre on established perennial species such as Canada thistle and Field bindweed. Use lower rate when annual weeds are small (2 to 3" tall) and growing actively. Use the higher rate on older and drought-stressed plants. Spray musk thistles and other biennial species while in seedling to rosette stage, and before flower stalks are initiated. The lower rate can be used in Spring during rosette stage. In Fall or after flower stalks have developed, use highest rate. Spray perennial weed in bud to bloom stage, or in good vegetative growth. Do not disturb treated area for at least 2 weeks after treatment, or until weed tops are dead. Do not plant any crop for 3 months after treatment or until chemical has disappeared from soil.

**ESTABLISHED PASTURES AND RANGELANDS:** Use 1/3 to 2-1/2 pints of product in sufficient water to give good coverage to one acre depending on type of weeds and stage of growth. Use only on established stands of perennial grasses. Do not use on bentgrass, alfalfa, clover, or other legumes. Do not use on newly seeded areas until grass is well established. Do not use from early boot to milk stage when grass seed productions is desired.

**Bitterweed, Broomweed, Croton, Docks, Kochia, Marshelder, Musk thistle and Other Broadleaf Weeds:** Use 2.5 to 2.7 pints of this product in 10 to 30 gallons of water per acre. If

weeds are young and growing actively, 1-1/4 pints per acre will provide control of some species. Deep-rooted perennial weeds may require repeated treatments in the same year or in subsequent years.

**Weed control in Newly Sprigged Coastal Bermudagrass:** Apply 1.25 to 2.7 pints of this product in 20 to 100 gallons of water per acre pre-emergence and/or post-emergence.

**WILD GARLIC AND WILD ONION CONTROL:** Apply up to 2.7 pints of product per acre making two applications, Fall-Spring or Spring-Fall, starting in the late Fall or early Spring.

**GENERAL WEED CONTROL (Airfields, Roadsides, Vacant Lots, Fencerows, Industrial Sites, Rights-of-Way, and similar areas):**

Use 1-1/4 to 3-3/4 pints of product per acre. Apply when most annual broadleaf weeds are still young and growing vigorously. Apply when perennial and biennial weeds are actively growing and near the bud stage, but before flowering. For best results on tansy ragwort and musk thistle, treat in rosette stage, before bolting. A second application is usually needed for best results on thistle, nettle, and bindweed. Treat wild onion or garlic in early Spring and in Fall when they are young and growing actively. Mix 2-1/2 pints of this product in 2 quarts kerosene or diesel oil then add this mixture to 100 gallons of water. Apply 300 to 500 gallons of spray per acre, depending on the stand. The addition of a wetting agent (spray adjuvant) is suggested. Usually 2-1/2 pints per acre will give adequate control. Do not use on herbaceous ground covers or creeping grass such as Bent. Legumes will usually be damaged or killed. Deep-rooted perennials may require repeat applications. Do not use on freshly seeded turf until grass is well established. Delay reseeding for 30 days.

**CONTROL OF SOUTHERN WILD ROSE:** On roadsides and fencerows, use 2-3/4 quarts of this product plus 4 to 8 ounces of an agricultural surfactant per 100 gallons of water and spray thoroughly as soon as foliage is well developed. Two or more treatments may be required. On rangeland, apply a maximum of 2.7 pints of this product per acre per application.

**SPOT TREATMENT IN NON-CROP AREAS:** To control broadleaf weeds in small areas with a hand or back pack sprayer, use 2-1/2 fluid ounces of this product per gallon of water and spray to thoroughly wet all foliage.

**GRASSES IN CONSERVATION RESERVE PROGRAM AREAS:** To control annual broadleaf weeds apply when weeds are actively growing. Use 1/4 to 1/2 pint per acre when weeds are small; use higher rates on older weeds. Excessive injury may result if applied to your grasses with fewer than 6 leaves to prior to grasses being well established. To control biennial and perennial broadleaf weeds in established grasses apply at a rate of 1-1/4 to 2-1/2 pints per acre. Apply to actively growing weeds. Treat when biennial weeds are in the seedling to rosette stage and before flower stalks become apparent. Treat perennial weeds in the bud to bloom stage.

**NOTE:** Suggest at least 2 gallons of water per acre by air and 5 gallons of water per acre by ground. Do not harvest or graze treated Conservation Reserve Program areas. Do not apply to grasses in the boot to dough stage if grass seed production is desired.

**BRUSH CONTROL**

The maximum application rate for forestry site preparation is 2.7 quarts per acre per application per site.

**WOODY PLANT CONTROL:** To control woody plants susceptible to 2,4-D such as Alder, Buckbrush, Elderberry, Sumac, Cherokee rose, Japanese honeysuckle, Virginia creeper, Wild grape and Willow on non-crop areas such as rights-of-way, fence rows and roadsides, use 1-1/4 to 1-3/4 quarts of product per acre in 30 to 100 gallons of water. Lower volume of water can be used unless applying through such equipment as Directa-Spra, Wobbler, Mini Wobbler, Spirometer. Spray brush 5 to 8 feet tall after Spring foliage is well developed. Wet all parts of the plants thoroughly, including stem and foliage, to the point of runoff. Higher volumes of up to 300 to 500 gallons of spray per acre may be necessary where the brush is very dense and over 6 to 8 feet high. Spraying can be effective at anytime up to 3 weeks before frost as long as soil moisture is sufficient for active growth of the brush. Control will be less effective in mid-summer during hot dry weather when soil moisture is deficient and plants are not actively growing. Oil or wetting agent may be added to the spray, if needed for increased effectiveness.

Hard-to-control species may require re-treatment next season. In general, it is better to cut tall woody plants and spray sucker growth when 2 to 4 feet tall.

**SAND SHINERY OAK AND SAND SAGEBRUSH:** On the oak, use 1-1/3 pints of this product in 5 gallons of oil or in 4 gallons of water plus 1 gallon of oil per acre. Apply by aircraft between May 15 and June 15. On the sagebrush, use 1-1/3 pints in 3 gallons of oil per acre and apply by aircraft when foliage is fully expanded and the brush is actively growing.

**BIG SAGEBRUSH AND RABBITBRUSH:** Use 1-1/3 to 4-1/3 pints in 2 to 3 gallons of oil or in 3 to 5 gallons of oil-water emulsion spray. For rabbitbrush, the 4-1/3 pints rate is usually required. Brush should be leafed out and growing actively when treated. Retreatment may be needed.

Chamise, Manzanita, Buckbrush, Coastal Sage, Coyotebrush and certain other Chaparral Species: Use 1-1/3 to 4-1/3 pints per acre in 5 to 10 gallons of water. One gallon of fuel oil may be included in the spray mixture for added effectiveness. Make applications by aircraft or ground equipment to obtain uniform spray coverage. For effective control, the brush must be fully leafed out and growing actively when sprayed. Retreatment may be needed. Consult state or local brush control specialists for most effective rate, volume and timing of spray application.

#### USES IN FOREST MANAGEMENT

**Conifer Release:** For control of Alder, apply 1 to 1-1/4 quarts of product per acre in 8 to 25 gallons of water, and apply as a foliage spray. Treat when 3/4 of the brush foliage has attained full size leaves and before new conifer growth reaches 2" in length. This is usually between early May and mid-June. Adjust treatment date depending on stage of growth and brush species. This may cause leader deformation on ex-posed firs, but they should overcome this during the second year after spraying. To control susceptible brush species such as ceanothus spp., chinquapin, madrone, manzanita, oil and tanoak and to release Douglas fir, hemlock, Sitka spruce or grand fir, apply 2 quarts of product per acre before new growth on Douglas fir is 2" long. To control manzanita and ceanothus in ponderosa pine, apply 1-3/4 quarts of Solution™ Emulsible before pine growth begins in Spring.

To increase performance, add 2 to 4 quarts of diesel, fuel oil, kerosene, or a suitable approved agricultural surfactant at recommended label rate.

After northern conifers, jack pine, red pine, black spruce, and white spruce cease growth and "harden off" (usually in Mid-July), a spray of 3/4 to 1-3/4 quarts of product in 8 to 25 gallons of water per acre may be applied by air to control certain competing hardwood species such as Alder, Aspen, Birch, and Willow. Since this treatment may cause occasional conifer injury, do not use if such injury cannot be tolerated. Consult your regional or extension forester or state herbicide specialist for recommendations to fit local conditions.

**Tree Injections (Pine Release):** To control hardwoods, such as Oaks, Hickory, Maple, Pecan, Elm, Sumac, Sweetgum and Hawthorn in forest and other non-crop areas, apply undiluted Solution™ Emulsible in a concentrate tree injector calibrated to apply .6 ml. per injection. Space injections 2" apart, edge to edge, completely around the tree and close to the base. The injector bit must penetrate the inner bark. On hard-to-kill species such as Hickory, Dogwood, Red maple, Blue beech and Ash make injections 1-1/2" apart, edge to edge. Treatment may be made at any time of the year. For best results, injections should be made during growing season, May 15 to October 15. For dilute injections, mix 2/3 gallon of Solution™ Emulsible in 19 gallons of water.

**Dormant Application (other than pine):** For the control of susceptible deciduous brush species such as Alder, cascara, cherry poplar and serviceberry apply up to 2 quarts of product per acre in sufficient diesel, fuel oil or kerosene for good coverage.

**Pine Only:** Make application while pine buds are still dormant. Apply 1-1/4 quarts of product per acre in sufficient water for good coverage by air or ground equipment. Do not use this application unless some pine injury is acceptable. Use of diesel, kerosene, or other oil, or addition of surfactants to spray mix may cause unacceptable pine injury.

**Herbaceous Weed Control:** To control over-wintering susceptible weeds such as false dandelion, klamath weeds, plantain, tansy ragwort apply 2/3 to 2 quarts of product in sufficient water for good coverage. Make application at rates and timing indicated above if pines are present. For control of hazel brush and similar species in the Lake States area, apply 1-1/4 quarts of product per acre in 8 to 25 gallons of water, when new shoot growth of Hazel is complete (usually mid-July).

**Site Preparation:** (As Budbreak Spray) - For control of Alder prior to planting seedlings,

apply 1-1/4 to 2-1/2 quarts of product per acre in 8 to 25 gallons of water, after Alder budbreak but before foliage is 1/4 full size. Application may be made by air or ground. If desired, diesel, fuel oil or kerosene may be substituted for water, air or ground. (As Foliage Spray) - For control of Alder prior to planting seedlings, apply 1-1/4 quarts of product per acre in 8 to 25 gallons of water, after most Alder leaves are full size. To increase penetration, 2 to 4 quarts per acre of diesel, fuel oil, kerosene, or a suitable approved agricultural surfactant at recommended label rates, may be added to the spray mixture.

#### TANK MIXES

Read and follow the label of each tank mix product used for precautionary statements, directions for use, geographic and other restrictions.

Using Riverdale Solution Emulsible and Buctril® ME4 for weed control on cereal grains (wheat, barley and rye): Buctril ME4 Broadleaf Herbicide will control some annual weeds that are resistant to Solution Emulsible and may be tank mixed with Solution Emulsible for broader spectrum weed control on small grains. In cereal areas except Washington, Oregon and Idaho, use 1/4 to 1/2 pint of Solution Emulsible plus 1/2 to 3/4 pint of Buctril ME4 per acre. In Washington, Oregon and Idaho: use 1/4 to 1/2 pint of Solution Emulsible plus 3/4 to 1 pint Buctril ME4 per acre. First mix the Solution Emulsible in water then add the Buctril ME4. Use the higher rates for larger weeds or where weed growth is slow due to dry or cold weather. Apply before weeds are 6 inches high. Use 10 to 20 gallons total spray volume per acre with ground equipment or 5 to 10 gallons total spray volume with air application. Use higher volume on larger weeds.

Using Riverdale Solution Emulsible and Sencor as knockdown herbicides for no-till: Solution Emulsible with Sencor DF alone or in combination with Dual, Lasso, Surflan or Prowl may be applied as an early preplant surface application for the control of certain broadleaf weeds and grasses in soybeans in minimum or no-till products. Application is recommended 30 days prior to planting. Apply at rate of 1-1/3 pints Solution Emulsible (1# A.I.) per acre with labeled rates of Sencor. Where grass herbicide is used in tank mix, apply at the rates specified on that product's label.

Using Riverdale Solution Emulsible and Turbo 8EC in reduced-tillage or no-till systems: Solution Emulsible may be applied in combination with Turbo 8EC for the control of annual grasses and broadleaf weeds and the suppression of emerged perennial weeds when soybeans are directly seeded into a stale seedbed, cover crop or in previous crop residues. Special precautions: poor weed control and/or crop injury may result if directions are not followed. Do not use a rib-type press wheel on your no-till planter or crop injury may result. Apply at a rate of 1-1/3 pints Solution Emulsible (1# A.I.) per acre with labeled rates of Turbo 8EC. Application is recommended 30 days prior to planting.

Using Riverdale Solution Emulsible and Poast as a burndown prior to planting soybeans: for broad spectrum post-emergence weed control, a tank mix application of Solution Emulsible with Poast may be made for control of emerged broadleaf and grass weeds before planting soybeans. Apply at a rate of 2/3 pints Solution Emulsible (1/2# A.I.) per acre with labeled rates of Poast.

Using Riverdale Solution Emulsible and Tahoe™ 3A tank Mixtures for Non-Crop Areas:  
Broadleaf Weed Control: Use 1-1/4 to 2-1/2 pints Solution Emulsible plus 2 to 6 pints Tahoe™ 4E (or 3 to 8 pints Tahoe 3A) per acre. For wider spectrum control of broadleaf weeds and woody plants. Apply as a broadcast spray in enough water to deliver 20 to 100 gallons total spray per acre. Apply when broadleaf weeds are actively growing. Woody Plant Control Broadcast Foliar Spray: Use 1/2 to 1-1/4 gallons Solution Emulsible plus 1-1/2 to 3 quarts Tahoe 4E (or 2 to 4 quarts Tahoe 3A) per acre. Apply as a broadcast spray in enough water to deliver 20 to 100 gallons total spray per acre. Apply when woody plants are actively growing. Woody Plant Control High Volume Leaf-Stem Treatment with Ground Equipment: Use 1/2 to 5 quarts Solution Emulsible plus 1-1/2 to 12 pints Tahoe 4E (or 2 to 16 pints Tahoe 3A) per acre. Mix 1/2 to 2 qts. Solution Emulsible plus 1-1/2 to 3 pts. Tahoe 4E (or 2 to 4 pints Tahoe 3A) in enough water to make 100 gallons of spray. Apply at a volume of 100 to 400 gallons of total spray per acre depending on size and density of woody plants. Thoroughly wet all leaves, stems, and roots collars of plants to be controlled. Woody Plant Control Aerial Application (Helicopter only): Use 1/2 to 1-1/4 gallons Solution Emulsible plus 3 to 4 quarts Tahoe 4E (or 4 to 6 quarts Tahoe 3A) per acre. Apply in a total spray volume of 210 to 30 gallons per acre using drift control equipment such as Microfoil boom or an effective drift control agent such as Lo-Drift

Spray Additive. Use the higher rates and volumes when plants are dense or under drought conditions.

Using Riverdale Solution Emulsible and Diablo™ Herbicide tank mixtures for Non-Crop Areas: Annual broadleaf weeds: Use 1-1/4 to 2-1/2 pints Solution Emulsible plus 1/2 to 1-1/2 pints Diablo. For wider spectrum control of broadleaf weeds and woody plants. Apply as a broadcast spray in enough water to deliver 20 to 100 gallons total spray per acre. Apply when broadleaf weeds are actively growing. Use the higher rates when treating dense or tall vegetative growth. Perennial and Biennial Broadleaf Weeds: Use 2 to 4 pints Solution Emulsible plus 1/2 to 6 pints Diablo. Apply as a broadcast spray in enough water to deliver 20 to 100 gallons total spray per acre. Apply when broadleaf weeds are actively growing but prior to flowering. Use the lower rates for biennials less than 3 inches rosette diameter. Use the higher rates for perennial weeds or for biennial weeds past the 3 inch rosette stage. Woody Plant Control Broadcast, High Volume, Stem Foliage or Aerial Application: Use 1/2 to 1-1/4 gallons Solution Emulsible plus 2 to 8 quarts Diablo. Apply as a broadcast spray in enough water to deliver 20 to 100 gallons total spray per acre or apply as a high volume stem foliage spray in enough volume to thoroughly wet leaves, stems and root collars (100 to 400 gallons per acre) or apply aerially in enough water to deliver total spray volume of 10 to 30 gallons per acre using drift control equipment such as the Microfoil Boom or an effective drift control agent such as Lo-Drift Spray Additive. Use the higher rates and volumes when plants are dense or under drought conditions.

Using Riverdale Solution Emulsible and Patriot, Spyder and Telar: To improve control of some target species, this product may also be tank mixed with Escort, Oust, and Telar herbicides for post-emergent weed control. Tank mixes have shown improved control where resistant bio-types are present.

NOTE: All intended tank mix combinations should be used only in recommended areas on the same broadleaf weed species found on both labels. For application methods and other use specifications, use the most restricted limitations from labeling of both products.

Local conditions may affect the use of this chemical. Consult State Agricultural Extension or Experiment Station weed specialist for specific recommendations for local weed problems and for information on possible lower dosages.

**STORAGE AND DISPOSAL**

**STORAGE:** Always use original container to store pesticides in a secured warehouse or storage building. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stock cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides. Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If container is damaged or if pesticide has leaked contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Direct-Spray, Wobbler, Mini Wobbler, Spirometer, Lo-Drift, and Bucril are registered trademarks of Rhone-Poulenc Ag Company  
 Surflan is a trademark of the Dow AgroSciences LLC  
 Escort, Oust and Telar are trademarks of E.I. DuPont de Nemours & Co. (Inc.)  
 Poast is a trademark of BASF Corp.  
 Dual is a trademark of Ciba-Geigy Corp.  
 Lasso is a trademark of Monsanto Agri Co.  
 Sencor is a trademark of Bayer AG  
 Turbo is a trademark of Mobay Corp.  
 Prowl is a trademark of American Cyanamid Co.  
 Patriot, Spyder, Diablo and Tahoe are trademarks of Riverdale - A Nufarm Company

**WARRANTY**

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risk of use, storage or handling of this material not in strict accordance with directions given herewith. (RV 051503A)