TELEPHONE (800) 345-3330 (708) 754-3330 FAX (708) 754-0314



CHEMICAL COMPANY

OFFICE: 425 WEST 194th STREET, GLENWOOD, IL 60425-1584 PLANT: 220 EAST 17th STREET, CHICAGO HEIGHTS, IL 60411-3699

June 4, 1993

Document Processing Desk (Notification) Registration Division - H7505C U. S. Environmental Protection Agency 401 M Street S.W. Washington, DC 20460

Dear Sir/Madam:

Subject: PR Notice 93-5 dated March 9, 1993

Riverdale Butox / 2,4-D Low Volatile Ester

EPA Reg. No. 223-141 OPP Identifier No. 196223

Please find enclosed the necessary paperwork to notify the Agency that we are revising our label in accordance with the above subject PR Notice. Be advised that this product is part of our "John Doe" private label program. As such, we do not market this product under the Riverdale name. Thus, there will not be any final printed label. If there is anything else we should do at this time, please contact us immediately.

Sincerely yours,

RIVERDALE CHEMICAL COMPANY

By Russell F. Sawyer Regulatory Affairs Manager

RFS:1s

Enc.





| | | | | | | | ζ |
|---------------------------------------|--|--|---------------------|--|---------------------|---|---------|
| Please read Instructions on rev | United States Environmenta | | m Approved | | _ | opproval expires to OPP Identifier Nur | |
| $^{(A)}$ Ω \square \square | rams (H7505C) | | Registrat Amendm | ion | | | |
| | 20460 r Pesticide: | | | 196223 | | | |
| | Sec | tion I | بها | | . . | | |
| . Company/Product Number | | 2. EPA Produ | ct Manager | | 3. Prop | osed Classificati | ion |
| 228-141 | . <u>_</u> | | I. Mill | er | ↓ — . | . — | |
| I. Company/Product (Name) | PM# | | | | | | |
| Riverdale Butoxy 2 | ,4-D Low Volatile Es | ster | 23 | | <u> </u> | | |
| . Name and Address of Applicar | | 6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling | | | | | |
| RIVERDALE CHEMICAL | COMPANY | to: | DOUCT IS SIT | Har of Igentic | ai in con | ірозіноп апо іа | Deinig |
| 425 West 194th Stre | | | | | | | |
| Glenwood, IL 6042 | EPA Reg. N | EPA Reg. No | | | | | |
| Check if this is a n | ew address | Product Nar | me | | | | |
| | Section | on II | | | | | |
| Amendment - Explain below | | Final printed labels in response to Agency letter dated | | | | | |
| Resubmission in response to | | "Me Too" Application. | | | | | |
| Notification - Explain below. | Other - | Other - explain below. | | | | | |
| Explanation: Use additional pa | ige(s) if necessary. (For section | and Section II.) | * | | | | |
| To comply with PR I | | | | DATE | 1111 | | |
| | Section | n lil | | <u>. </u> | | | |
| 1. Material This Product WIN B | | Maria - Calabia Deale | | 2. Type of Co | antainar | | |
| | Packaging | Water Soluble Packa | ging | 1 — | | | |
| Yes* | Yes | Yes | | · | vletal Plastic | | |
| L N₀ L | No | ∐ No | | 1 | Glass | | |
| lar s | es," No. per | if "Yes," | No. per | | Paper Other (Spe | ecity) | |
| * Certification must be submitted. | Package wgt. container | Package wgt. | container | 1 | | | • |
| 3. Location of Net Contents Infor | mation 4. Size(s) of R | letail Container | 5. l | ocation of Lab | el Directio | ens . | |
| Label Contai | ner | | IE | On Label On Labeling | a accomp | anying groduct | |
| 6. Manner In Which Label Is Affin | ced To Product Lithogra | | Other (| | <u>g 0000p</u> | , | • • • • |
| | Paper of Stendik | | | | | | • |
| | | lon IV | | | _ | 4 | |
| 1. Contact Point (Complete item | | | ntacted, if ne | | | | |
| Name | Ti | tle | | T | elephone | No. (Include Are | ea Gode |
| Russell F. Sawyer | , | Regulatory Af | faire Ma | nager | 708/ | 754-3330 | • • • |
| | Certification | | | | | 5. Date Applicati | en • |
| | ave made on this form and all at ngly false or misleading stateme | tachments thereto are | | | e | Received • (Stamped) | ••• |
| 2. Signature | | Title | | | | | |
| en 1 | _ | | | | | | |
| 7 '+1 Jan | }\ | Regulatory Af | fairs Ma | anager | | | |
| Typed Name | () 5 | Date | | | - 1 | | |

Russell F. Sawyer EPA Form 8570-1 (Rev. 12-90)

Previous editions are obsolete.

June 4, 1993

White -EPA File Copy (original) Yellow - Applicant copy

RIVERDALE

BUTOXY 2.4-D LOW VOLATILE ESTER

Isomer Specific AOAC Method, Equivalent to: *2,4-Dichlorophenoxyacetic Acid 42.8%, 3.84 lbs./gal.

KERP OUT OF REACH OF CHILDREN

CAUTION - CAUCION

Si Usted no entiende la etiqueta, busque a alguien para que se la explique a Usted en detalle.

SEE SIDE AND BACK PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS
AND STATEMENT OF PRACTICAL TREATMENT

NET CONTENTS

GAL.

EPA REG. NO. 228-141

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid inhalation of vapors or spray mist and contact with skin, eyes and clothing. When mixing, loading or applying this product or repairing or cleaning equipment used with this product, wear eye protection (face shield or safety glasses), chemical resistant gloves, long-sleeved shirt, long pants, socks and shoes. It is recommended that safety glasses include front, brow and temple protection. For aerial applications in an enclosed cockpit and applicators applying this product from a tractor

NOTE: Spanish Language is optional.

Revised A/O 3/9/93

HOTIFICATION
LABEL NOT REVEYUD
PER PRI NOTICE 64-6
DATE. 7/1/93

MIVELIGATE DICONA TO A DI DOM TOTALITE PROCES

that has been a completely enclosed cab, eye protection is not required. Wash hands, face and arms with soap and water as soon as possible after mixing, loading or applying this product. Wash hands, face and arms with soap and water before eating, smoking or drinking. Wash hands and arms before using toilet. After work, remove all clothing and shower using soap and water. Do not reuse clothing worn during the previous day's mixing and loading or application of this product without cleaning first. Clothing must be kept and washed separately from other household laundry. Remove saturated clothing as soon as possible and shower. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Get medical attention immediately. This product

contains petroleum distillates. DO NOT induce vomiting

HABE -

or give anything by mouth to an unconscious person.

IF ON SKIN: Wash thoroughly with plenty of soap and water. Ge

medical attention if irritation persists.

IF IN EYES: Flush eyes with water for 15 minutes and get medical

attention.

If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron. If this container is five gallons or more in capacity, then a mechanical system (probe and pump) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

Please see additional Precautionary Statements in Directions For Use instructions.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Drift or runoff may adversely affect fish and non-target plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Vapors from this product may injure susceptible plants in the immediate vicinity. Use care to avoid spray contact or drift to 2,4-D susceptible plants such as cotton, tomatoes, flowers, grapes, fruit trees and ornamentals. Do not permit spray mist containing this product to drift onto them. Do not spray when the wind is blowing towards susceptible crops or ornamental plants. Use coarse sprays and/or low spray pressure to minimize drift. On cropland and along roadsides, do not exceed 20 psi pressure. Spray drift can be lessened by keeping the spray boom as low as possible, by spraying when wind velocity is low, and by stopping all spraying when wind exceeds 6 to 7 miles per hour. Do not use the same spray equipment for applying other materials to 2,4-D susceptible crops as injury may result. It is best to see a separate sprayer for application of insecticides and fungicides. Clean and rinse spray equipment using soap or detergent and water or suitable chemical cleaner, and rinse thoroughly before reuse for other spraying. Do not contaminate water when disposing of equipment

> Killedator Grayes in 1985 Markeyes (1985)

> > Service 12 (12 Aft)

washwaters. Do not apply this product through any type of irrigation system. Do not contaminate domestic or irrigation waters. Excessive amounts of this product in the soil may temporarily inhibit seed germination and plant growth.

Most groundwater contamination involving phenoxy herbicides such as 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

When using on Pastures and Rangeland Grasses there is a (1) 7 day pregrazing interval for dairy cattle: (2) 30 day preharvest interval for grass cut for hay; and (3) 3 day pre-slaughter interval for meat animals.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

RE-ENTRY STATEMENT

Do not enter treated areas without protective clothing until sprays have dried. Because certain states may require more restrictive re-entry intervals for various corps treated with this product, consult your State Department of Agriculture for further information.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Warnings should state "Do not enter treated areas unless wearing chemical resistant, full body clothing, including NIOSH approved respirator, goggles, and gloves until sprays have dried". When oral warnings are given, warnings shall be given in language customarily understood by workers. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. Written warnings must include the following information: "CAUTION: Area treated with 2,4-D Butoxy Low Volatile Ester on date of application. Do not enter without appropriate protective clothing until sprays have dried. In case of accidental exposure follow precautionary statements on label."

ADDITIONAL PRECAUTIONARY STATEMENTS FOR AGRICULTURAL CROP USES AND INDUSTRIAL APPLICATIONS

The rates of applications for pastures and rangelands are per acre per application per site. Do not enter or allow worker entry into treated areas during restricted-entry interval of 12 hours. There isn't any reentry restrictions for Industrial applications nor non-crop uses of pastures and rangelands.

GENERAL INFORMATION

This product is a low volatile ester especially prepared for use on crops and weeds where a susceptible crop in the near vicinity may be injured by a more volatile product. It is recommended for control of numerous broadleaf weeds and certain 2,4-D susceptible woody plants without injury to most established grasses. For best results apply this product as a water or oil spray during warm weather when young, succulent weeds or brush are actively growing. Application under drought conditions often will give poor results. The lower recommended rates will be satisfactory on susceptible annual weeds. For perennial weeds and conditions such as the very dry areas of the western states, where control is difficult, the higher recommended rates should be used. Deep-rooted perennial weeds such as Canada thistle and field bindweed and many woody plants usually require repeated applications for maximum control.

Unless otherwise directed, application rates should be 1 to 10 gallons of total spray by air or 5 to 25 gallons by ground application equipment. Higher gallonage may be used if desired to improve spray coverage. If band treatment is used, base the dosage rate on actual area to be sprayed. In either case, use the same amount of 2,4-D recommended per acre. Although water quantities may vary due to different type of application equipment, sufficient water must be used to provide for complete and uniform coverage. When product is used for weed control in crops, the growth stage of the crop must be considered. For crop uses, do not mix with oil, surfactants, or other adjuvants unless specifically recommended on label. To do so may reduce herbicide's selectivity and could result in crop damage.

Aerial applications should be used only when there is no danger of drift to susceptible crops. Many states have regulations concerning aerial application of 2,4-D formulations. Consult local regulatory authorities before making applications. Although this product is a low volatile formulation, at temperatures above 90°F vapors may damage susceptible crops growing nearby.

Riverdale Butoxy 2,4-D Low Volatile Ester is especially prepared for use on crops and weeds where a susceptible crop in the near vicinity may be injured by a more volatile product.

SOME WEEDS READILY KILLED DURING EARLY STAGES OF GROWTH

ANNUALS

PERENNIALS

| Beggarsticks | Kochia | Hedge bindweed | Healall |
|------------------|--------------------|----------------|-----------------------|
| Black madic | Marsh elder | Buckhorn . | Nettle, stinging |
| butter print | Mustards | Bull thistle | Lawn pennywort |
| or Velvetleaf | Pigweeds | Burdock | Plantains |
| Cocklebur | Puncture vine | Chickweed | Sunflower (perennial) |
| Fanweed | Ragweed | mouseear | Vervains |
| Hemp (mcrijuana) | Sunflower (annual) | Dandelion | Vetch |

TREATMENT IN CORN AND SORGHUMS - Use from 1/2 to 1 pint Riverdale Butoxy 2,4-D Low Volatile Fster with recommended amounts of water to make per acre applications. Time of application should be determined by weed growth with the ideal time after emergence of majority of weeds but before they have damaged the crop. The crop is most sensitive when growing fast and temperatures and moisture favorable for plant growth. As a general rule, the crop is less susceptible from 3 inches up to 10 inches. Use

drop nozzle when corn or sorghum is over 10 inches tall. Do not apply from tasseling to dough stage.

TREATMENT IN SMALL GRAINS - WHEAT, OATS, BARLEY AND RYE - Use from 1/2 to 1 pint Riverdale Butoxy 2,4-D Low Volatile Ester with recommended amounts of water to make per acre applications for ground equipment and airplane application. Good results have also been obtained with airplane application using diesel oil as the diluent instead of water. Make application after the crop is fully tillered until jointing is under way. Stop treatment before boot stage is reached. Spraying in small grain is frequently desirable after the crop has reached the milk stage. At such time annual and perennial weeds often interfere with harvesting and produce seeds which lower quality of threshed grain. Under such conditions use from 1 to 1-1/2 pints Riverdale Butoxy 2,4-D Low Volatile Ester with recommended amounts of water to make per acre applications. Ground equipment will cause some mechanical damage to the crop. When applying with plane use dosage in 2 to 3 gallons diesel oil.

GENERAL BROADLEAF CONTROL ON PASTURES AND ON NON-CROPPED LAND - Do not apply to clover or newly seeded pastures.

ANNUAL WEEDS - Most annual broadleaf weeds can be controlled with 1 to 1-1/2 pints in 50 to 100 gallons of water per acre. Make application when weeds are young and actively growing.

New seedings of grass should not be treated until they are well established. For controlling annual weeds in farm yards, fence rows along highways and similar locations use 1 quart in 5 to 10 gallons water per acre. Possible injury may occur to bentgrass, St. Augustine, grass and clover.

PERENNIAL WEEDS - Many of the perennial weeds in pastures, including ironweed, vervain and asters are moderately tolerant to 2,4-D. For control of these type weeds use I quart Riverdale Butoxy 2,4-D Low Volatile Ester in 5 to 10 gallons water per acre. Make application not later than the bud stage and repeat treatment the following season.

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near seeds, fertilizers, insecticides or fungicides. Containers should be opened in well ventilated areas. Do not contaminate water, food or feed by storage or disposal. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If container is damaged or if pesticide has leaked contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for re-

cycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic containers also disposable by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Local conditions may affect the use of this chemical. Consult State Agricultural Extension or Experiment Station weed specialists for specific recommendations for local weed problems and for information on possible lower dosages.

WARRANTY: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risk of use, storage or handling of this material not in strict accordance with directions given herewith. (PR933 042194)