

228-145

1/18/2006

Page 183

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

JAN 18 2006

Theodore D. Head
Nufarm Americas Inc.
1333 Burr Ridge Parkway, Suite 125A
Burr Ridge, IL 60527-0866

Dear Mr. Head:

SUBJECT: Label Amendment
Weedestroy AM 40
EPA Registration No. 228-145

The label amendment referred to above, submitted in accordance with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records. Please submit one (1) copy of your final printed labeling before you release the product for shipment. This amended labeling supersedes all previously accepted ones.

Sincerely yours,

/s/

Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

7505C								
Miller								
1/17/06								

273

SUPPLEMENTAL LABELING

READ AND FOLLOW THE ENTIRE LABEL BOOKLET FOR RIVERDALE WEEDDESTROY AM 40 BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for this product and this supplement.

ACCEPTED

JAN 18 2006
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

228-145



Riverdale
WEEDestroy AM
40

EPA Reg. No. 228-149

Maximum Use Rates For Forestry and Non-Crop Sites

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

All applicable directions and precautions in the label booklet for this product must be followed.

Also refer to additional "GENERAL INFORMATION" and "WEEDS CONTROLLED" in the label booklet. INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY, DO NOT USE THIS PRODUCT. Product should not be used in or near greenhouses.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR, Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. For Turf use, do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

young and growing vigorously. Apply when perennial and biennial weeds are actively growing and near the bud stage, but before flowering. Thoroughly wet weeds when applying this mixture. For best results on Tansy ragwort and Musk thistle, treat in rosette stage, before bolting. Treat Wild onion or garlic in early Spring and in Fall when they are young and growing actively. The addition of a wetting agent (spray adjuvant) is suggested. Usually 4 pints per acre will give adequate control. Do not use on herbaceous ground covers or creeping grass such as Bent. Legumes will usually be damaged or killed. Deep-rooted perennials such as Bindweed, Whitetop, Perennial sowthistle, Blue lettuce, Nettle, Bur ragweed, Canada thistle and other noxious perennials somewhat resistant to 2,4-D may require repeat applications to kill. Do not use on freshly seeded turf until grass is well established. Delay reseeding for 30 days. For chemical mowing applications on roadside and utility rights-of-way, using low volume spray equipment such as the "Lucas" 64 system use 1/2 gallon of this product in 1 to 5 gallons of water per acre.

Seasonal: The maximum seasonal application rate to non-crop sites is 4 pounds 2,4-D Acid equivalent per acre per application site.

CONTROL OF SOUTHERN WILD ROSE: On roadsides and fencerows, use 1/2 gallon of this product plus the recommended rate of a high quality surfactant per 100 gallons of water and spray thoroughly as soon as foliage is well developed. Two or more treatments may be required.

TANK MIXES

Using Weedestroy AM-40 and Tahoe™ 4E or Tahoe™ 3A tank mixtures for Non-Crop Areas: Broadleaf Weed Control - Use 2 to 4 pints of this product plus 2 to 6 pints of Tahoe 4E (or 3 to 8 pints of Tahoe 3A) per acre. For wider spectrum control of broadleaf weeds and woody plants, apply as a broadcast spray in enough water to deliver 20 to 100 gallons total spray per acre. Apply when broadleaf

GENERAL WEED CONTROL (Airfields, Roadsides, Vacant Lots, Drainage Ditchbanks, Fencerows, Industrial Sites, Rights-of-Way, Utility Power Lines, Railroads, and similar areas): Use 1/2 gallon of product per acre. Use sufficient gallonage for thorough and uniform coverage. Apply when most annual broadleaf weeds are still

weeds are actively growing. **Woody Plant total Broadcast Foliar Spray** - Use 1 gallon of this product plus 1-1/2 to 3 quarts of Tahoe 4E (or 2 to 4 quarts of Tahoe 3A) per acre. Apply as a broadcast spray in enough water to adequately wet foliage. This may require 20 to 100 gallons of water per acre. Apply when woody plants are actively growing. **Woody Plant Control High Volume Leaf-Stem Treatment with Ground Equipment** - Use 1 gallon of this product per acre plus 1-1/2 to 12 pints of Tahoe 4E (or 2 to 16 pints of Tahoe 3A). Mix 1 to 3 quarts of this product per acre plus 1-1/2 to 3 pints of Tahoe 4E (or 2 to 4 pints of Tahoe 3A) in enough water to wet all parts of the brush foliage, stem and bark. This may require 100 to 400 gallons of water per acre depending on size and density of woody plants. Thoroughly wet all leaves, stems, and root collars of plants to be controlled. **Woody Plant Control Aerial Application (Helicopter only)** - Use 1 gallon of this product per acre plus 3 to 4 quarts of Tahoe 4E (or 4 to 6 quarts of Tahoe 3A). Thoroughly wet all leaves, stems, and root collars of plants to be controlled. This may require 10 to 30 gallons of water per acre using drift control equipment such as Microfoil boom or an effective drift control agent such as Lo-Drift Spray Additive. Use the higher rates and volumes when plants are dense or under drought conditions.

NOTICE

Before buying or using this product, read "WARRANTY LIMITATIONS AND DISCLAIMER" in the label booklet. If terms are not acceptable, return unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by the Buyer or any other User constitutes acceptance of the terms under the "WARRANTY LIMITATIONS AND DISCLAIMER".

**Manufactured By
Nufarm Americas Inc.
Burr Ridge, IL 60521**

09/05

Using Weedestroy AM-40 and **Diablo® Herbicide** tank mixtures for Non-Crop Areas: **Annual broadleaf weeds** - Use 2 to 4 pints per acre of this product plus 1/2 to 1-1/2 pints of Diablo. For wider spectrum control of broadleaf weeds and woody plants - Apply as a broadcast spray in enough water to wet all parts of the brush foliage, stem and bark. This may require 20 to 100 gallons of water per acre. Apply when broadleaf weeds are actively growing. Use the higher rates when treating dense or tall vegetative growth. **Perennial and Biennial Broadleaf Weeds** - Use 3 to 6 pints of this product per acre plus 1/2 to 6 pints Diablo. Apply as a broadcast spray in enough water to wet all parts of the brush foliage, stem and bark. This may require 20 to 100 gallons water per acre. Apply when broadleaf weeds are actively growing but prior to flowering. Use the lower rates for biennials less than 3 inches rosette diameter. Use the higher rates for perennial weeds or for biennial weeds past the 3 inch rosette stage. **Woody Plant Control Broadcast, High Volume, Stem Foliage or Aerial Application** - Use 1 gallon of this product per acre plus 2 to 8 quarts of Diablo. Apply as a broadcast spray in enough water to wet all parts of the brush foliage, stem and bark. This may require 20 to 100 gallons water per acre or apply as a high volume stem foliage spray in enough volume to thoroughly wet leaves, stems and root collars (100 to 400 gallons water per acre) or apply aerially in enough water to wet all parts of the brush foliage, stem and bark. This may require 10 to 30 gallons of water per acre using drift control equipment such as the Microfoil Boom or an effective drift control agent such as Lo-Drift Spray Additive. Use the higher rates and volumes when plants are dense or under drought conditions. Tank mixing 2,4-D with Diablo Herbicide will improve control of late, post-emergent applications in corn, Fall and Spring seeded wheat, Fall seeded barley, pastures, rangelands, non-crop use, turfs and lawns.

This product will either kill, control or suppress the weeds listed in the label booklet for this product. Some of these species may require repeat spot applications even under ideal conditions.