



Riverdale

TURF WEEDSTROY-D

For control of broadleaf weeds in turf, golf courses, park grounds and similar non-crop turf areas

ACTIVE INGREDIENT

*Dimethylamine salt of 2,4-Dichlorophenoxyacetic Acid

25.5%

INERT INGREDIENTS

74.5%

TOTAL

100.0%

*2,4-Dichlorophenoxyacetic Acid Equivalent: 21.2% - 1.92 Lbs./Gal. Isomer Specific by AOAC Method No. 6 D015

CAUTION: KEEP OUT OF REACH OF CHILDREN See Additional Precautions on Side Panels

CONTROLS THESE TURF WEEDS:

Buck horn	Red Sorrel
Plantains	Wild Garlic
Curled Dock	Wild Onion
Dandelion	Many Others

DIRECTIONS FOR USE

Use 2 to 3 quarts per acre (1½ to 2½ ounces in 1-2 gallons waters, per 1,000 sq. ft.) in sufficient water for uniform spray coverage. Repeat applications may be necessary. Use lower rate for easy-to-kill species, i.e. dandelion and plantains. Use higher rate for more effective general control of weeds.

Use preferably in the spring or fall when weeds are actively growing. The killing action is more rapid when applied on warm clear sunny days than when weather is cool, cloudy or wet.

NOTE: Do not use on newly seeded lawns. Do not mow turf for several days before or after treatment. In some areas Dichondra, clover, bent grasses, carpet buffalo, and St. Augustine grasses may be injured.

NET CONTENTS

GALLONS

CAUTIONS

May cause skin irritation.
Harmful if swallowed.
Avoid contact with eyes, skin or clothing.
Avoid inhaling spray mist.
Do not use on light sandy soils.
Do NOT store with foodstuffs.

This product will kill or seriously injure many desirable forms of vegetation. Do not apply directly to flowers, fruits, vegetables, grapes, ornamentals, cotton or other desirable plants. Do not use when there is hazard from drifting mists. (Coarse sprays are less likely to drift.)

Vapors from this product may injure susceptible plants in the immediate vicinity. Avoid contamination of water used for domestic purposes and irrigation purposes. Excessive amounts of this product in the soil may temporarily inhibit seed germination and plant growth.

Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not store near fertilizers, seeds, insecticides or fungicides.

Avoid contamination of feed and foodstuffs.

Do not reuse empty drums. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies. Do not burn.

Spray equipment used in applying this product should be thoroughly cleaned before using for any other purpose. Use repeated flushing with soap and warm water or suitable chemical cleaner. It is best to use a separate sprayer for application of insecticides and fungicides.

Follow metho
condit
not sp
influen
duct
Buyer
handli
corda

ACC
22
Under the
Fungicide
as an
registered
EPA Reg

RI



Riverdale

TURF WEEDESTROY-D

For control of broadleaf weeds in turf, golf courses, park grounds, and similar non-crop turf areas.

ACTIVE INGREDIENT

Dimethylamine salt of 2,4-Dichlorophenoxyacetic Acid 25.5%

INERT INGREDIENTS

25.5%

74.5%

TOTAL

100.0%

2,4-Dichlorophenoxyacetic Acid Equivalent 21.2% 192 Lbs. Gal. Form. Specific by AOAC Method No. 6-D015

CAUTION: KEEP OUT OF REACH OF CHILDREN See Additional Precautions on Side Panels

CONTROLS THESE TURF WEEDS:

- | | |
|-------------|-------------|
| Buck horn | Red Sorrel |
| Plantains | Wild Garlic |
| Curled Dock | Wild Onion |
| Dandelion | Many Others |

DIRECTIONS FOR USE

Use 2 to 3 quarts per acre (1 1/2 to 2 1/2 ounces in 1-2 gallons waters, per 1,000 sq. ft.) in sufficient water for uniform spray coverage. Repeat applications may be necessary. Use lower rate for easy-to-kill species, i.e. dandelion and plantains. Use higher rate for more effective general control of weeds.

Use preferably in the spring or fall when weeds are actively growing. The killing action is more rapid when applied on warm clear sunny days than when weather is cool, cloudy or wet.


NOTE: Do not use on newly seeded lawns. Do not mow turf for several days before or after treatment. In some areas Dichondra, clover, bent grasses, carpet buffalo, and St. Augustine grasses may be injured.

NET CONTENTS GALLONS

NOTICE

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

DO NOT Store with Foodstuffs.



EPA Reg. No. 228-148
EPA Est. No. 228-IL-1



59