

RIVERDALE

25-3-3

WEED AND FEED

FERTILIZER PLUS 2,4-D & MCPP
FOR ORNAMENTAL TURF USE

Treats (or covers) 5000 Sq. Feet (50' x 100')

This product is a remarkably effective lawn fertilizer which combines both fast and long-lasting feed action, plus effective weed control. This product produces a thick, green turf while killing weeds to make way for new turf growth.

Riverdale 25-3-3 Weed and Feed is designed for the control of dandelion, clover, and many other broadleaf weeds. Weeds contacted by this product gradually die and disappear from the lawn. See back panel for a list of weeds controlled by Riverdale 25-3-3 Weed and Feed.

ACTIVE INGREDIENTS:

Isooctyl Ester of 2,4-Dichlorophenoxyacetic Acid* 0.87%
Isooctyl Ester of 2-(2-Methyl-4-chlorophenoxy)propionic Acid** 0.90%

INERT INGREDIENTS: 98.23%
TOTAL 100.00%

ISOMER SPECIFIC AOAC METHOD, EQUIVALENT TO:

*2,4-Dichlorophenoxyacetic Acid 0.57%
**2-(2-Methyl-4-chlorophenoxy)propionic Acid 0.59%


GUARANTEED ANALYSIS:°

Total Nitrogen (N) 25%
 1.0% Ammoniacal Nitrogen
 24.0% Urea Nitrogen
Available Phosphoric Acid (P₂O₅) 3%
Soluble Potash (K₂O) 3%

Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated Urea, Urea, Triple Super Phosphate, and Muriate of Potash.

Chlorine (Cl) Maximum 2.5%

The urea nitrogen in this product has been coated to provide 6.8% coated slow release urea nitrogen.



NOTIFICATION
LABEL NOT REVIEWED
FOR FIF NOTICE 88-6
DATE JUL 6 1990

KEEP OUT OF REACH OF CHILDREN

CAUTION

**SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
AND STATEMENT OF PRACTICAL TREATMENT**

NET WEIGHT LBS.

EPA REG. NO. 228-152

EPA EST. NO. 228-II-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584

(Back Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Avoid inhalation of dust and contact with eyes, skin, or clothing. Wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT

- IF SWALLOWED: Get Medical Attention.
- IF IN EYES: Flush Eyes with Water for 15 Minutes. Get medical attention.
- IF ON SKIN: Wash skin with Soap and Water.


ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

This product may be used on Bluegrass, Fescue, Rye, Common Bermuda, Bahia, and Zoysia Lawns. DO NOT USE on Dichondra, Carpet and St. Augustine grasses. DO NOT USE this



product where desirable clovers are present or allow the product to "wind drift" on flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth in lawns and control many broadleaf weeds such as: Blue lettuce, Buckhorn, Burdock, Chickweed, Chicory, Cinquefoil, Clover, Cocklebur, Croton, Dandelion, Dock, Fanweed, Galinsoga, Goldenrod, Ground ivy, Hawkweed, Healall, Henbit, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarters, Mallow, Morningglory, Mustard, Nettles, Oxalis, Pennywort, Peppergrass, Pigweed, Plantain, Povertyweed, Purslane, Ragweed, Shepherdspurse, Smartweed, Speedwell, Spotted spurge, Thistle, Vervain, Vetch, Wild aster, Wild lettuce, Wild radish, and Yarrow.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Apply when grass is moist (after a rain, sprinkling or dew). For most effective broadleaf weed killing action, avoid unnecessary disturbances, including watering, of treated areas for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours following treatment. Avoid contact with desirable shrubs, plants, vegetables, or flowers. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. Do not cut grass 1 to 2 days before and after application.


HOW TO APPLY: This 20 (or 18) pound bag should be used on 5000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds (3.6 for 18# bag) per 1000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 sq. ft. and adjust to apply at the rate of 1 lb. (.9 lb. for 18# bag) per 250 sq. ft. Clean spreader thoroughly after use.

SPREADER SETTINGS
For One Pass Application°

<u>MAKE</u>	<u>SETTINGS</u>	<u>MAKE</u>	<u>SETTINGS</u>
Cyclone	5-1/4 - 5-3/4	Montgomery Wards	6-3/4
Scotts 35	5-1/2	1091,1093,1095	
Scotts C-1	J - L	Montgomery Wards	5
Central	7 - 8	1080,1081	
K-Mart	7 - 8	Precision	6

Sweep all fertilizer materials from concrete surfaces immediately after application. Do not wash material from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.



Consumer Statement (for bags less than 50#)

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near other feed or foodstuffs. To prevent cross-contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides or seeds.

PRODUCT DISPOSAL: To dispose of bag which contains unused product, securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discard in trash.

Commercial Statement (bags 50# and over)

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near seeds, fertilizers, insecticides or fungicides. Do not stack more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay away from smoke.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

The following note will not be part of the actual label copy.

°NOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers.

