

l

08/17/2005

(Commercial Label)

RIVERDALE

GRANULAR LAWN WEED KILLER

This product is designed for the control of dandelion, clover, and many other broadleaf weeds.

Treats (or covers) 10,000 Square Feet (100' x 100')

 ACTIVE INGREDIENTS:
 Isooctyl (2-ethylhexyl) Ester of 2,4 Dichlorophenoxyacetic Acid*
 1.57%

 2-(2-Methyl-4-Chlorophenoxy)propionic Acid
 0.58%

 INERT INGREDIENTS:
 97.85%

 TOTAL
 100.00%

ISOMER SPECIFIC AOAC METHOD, EQUIVALENT TO: *2,4-Dichlorophenoxyacetic Acid

Riverdale is a Registered Trademark of Riverdale - A Nufarm Company

KEEP OUT OF REACH OF CHILDREN

CAUTION

SHE BELOW FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Avoid inhalation of dust and contact with eyes, skin, or clothing. Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

- IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
- IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN Take off contaminated clothing. Rinse skin immediately with plenty of OR CLOTHING: water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

Revised 4/18/03 Changed First Aid Statement per PR 2001-1 and Disposal Statement per PR 2001-6 and added an Environmental Statement both via Amendment. 7/9/03 Corrected Ornamental Sites per EPA's Request 5/28/03 Meeting. 1/3

1.05%

AUG 1 7 2005

Under the Federal Insectioide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

228-154

f

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters, nor pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

It is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas.

This product may be used on Bluegrass, Fescue, Rye, Common Bermuda, Bahia, and Zoysia Lawns. DO NOT USE on Dichondra, Carpet and St. Augustine grasses. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" on flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed for control of broadleaf weeds in lawns such as: Bindweed, Black medic, Blue lettuce, Buckhorn, Burdock, Chickweed, Chicory, Cinquefoil, Clover, Cocklebur, Croton, Dandelion, Dock, Fanweed, Galinsoga, Goldenrod, Ground ivy, Hawkweed, Healall, Henbit, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarters, Mallow, Morningglory, Mustard, Nettles, Oxalis, Pennywort, Peppergrass, Pigweed, Plantain, Povertyweed, Purslane, Ragweed, Shepherdspurse, Smartweed, Speedwell, Spotted spurge, Thistle, Vervain, Vetch, Wild aster, Wild lettuce, Wild radish, and Yarrow.

WHEN TO APPLY: Best result are obtained when weeds are growing actively in Spring or early Fall. Apply when grass is moist (after a rain, sprinkling or dew). For most effective broadleaf weed killing action, avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours following treatment. Avoid contact with desirable shrubs, plants, vegetables, or flowers. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. Do not cut grass 1 to 2 days before and after application.

HOW TO APPLY: This 20 pound bag should be used on 10,000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is equal to one application.

The following settings are <u>approximate</u> for applying contents of this package at a suggested rate of 2 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 500 square feet and adjust to apply at the rate of 1 pound per 500 square feet. Clean spreader thoroughly after use.

SPREADER SETTINGS For One Pass Application

MAKE	<u>SETTINGS</u>	MAKE	SETTINGS
Cyclone Scotts 35	5-1/4 to 5-3/4 5-1/2	Montgomery Wards 1091,1093,1095	6-3/4
Scotts C-1	J - L	Jackson (Drop Type)	
Central	7 - 8	Modern Show 'N Tell	€ 3−1/2
K-Mart	78	Model 105	7

Environmental Benefits

ť

1

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increated mawing height and frequency will produce smaller grass clippings that will decompose more readily so nultients go back into your lawn.

✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day

✔ Reduces Glare & Reduces Noise

- ✓ Traps Dust, Smoke Particles & other Poliutants
- Putties Water & Modifies Temperature
- ✓ Improves Soll Condition & Controls Erosion

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not stack more than two pallets high. Do not store near open containers of fertilizer, seed, or other pesticide. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay away from smoke.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions. (RV 070903)

NET WEIGHT: 20 LBS

EPA REG. NO. 228-154

EPA EST. NO. 228-IL-1

MANUFACTURED BY NUFARM AMERICAS INC., BURR RIDGE, ILLINOIS 60527-0866

The following note will not be part of the actual label copy

NOTE Spreader Settings will vary among various commercial fertilizer manufacturers