

 <p align="center"><b>U.S. ENVIRONMENTAL PROTECTION AGENCY</b>  <b>Office of Pesticide Programs</b>  <b>Registration Division (7505P)</b>  <b>Ariel Rios Building</b>  <b>1200 Pennsylvania Ave., NW</b>  <b>Washington, D.C. 20460</b></p>	EPA Reg. Number: 228-156	Date of Issuance: SEP 17 2008
	Term of Issuance:	
	Name of Pesticide Product: MCPA L.V. 4 Ester	
<p align="center">NOTICE OF PESTICIDE:</p> <p align="center"> <input type="checkbox"/> Registration  <input checked="" type="checkbox"/> Reregistration          (under FIFRA, as amended)       </p>		
Name and Address of Registrant (include ZIP Code): Nufarm Americas, Inc. 150 Harvest Dr., Suite 200 Burr Ridge, Illinois 60521		
<p><small>Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.</small></p>		
<p>On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.</p> <p>This product is reregistered in accordance with FIFRA section 4(g)(2)(C) provided you:</p> <ol style="list-style-type: none"> <li>1. Submit and/or cite all data required for registration/reregistration review of your product when the Agency requires all registrants of similar products to submit data.</li> <li>2. Per the acute toxicity review, the "IF SWALLOWED" statements must be revised to read:             "IF SWALLOWED:            Immediately call a poison control center or doctor.            Do not induce vomiting unless told to by a poison control center or doctor.            Do not give any liquid to the person.            Do not give anything by mouth to an unconscious person."</li> </ol> <p>Based on toxicity ranking per the acute toxicity review, the First Aid statements should be placed on the label in the following order:             "IF ON SKIN OR CLOTHING:...            IF SWALLOWED:...            IF INHALED:...            IF IN EYES:..."</p>		
Signature of Approving Official:  Joanne I. Miller Product Manager 23 Herbicide Branch Registration Division (7505P)	Date:  SEP 17 2008	

2) Per the acute toxicity review, the Note to Physician must be revised to read:

“Note to Physician

May pose an aspiration pneumonia hazard. Contains petroleum distillate.”

3) Per the acute toxicity review, the Hazards to Humans and Domestic Animals must be revised to read:

“WARNING

May be fatal if absorbed through skin. Causes skin irritation. Harmful if swallowed. Do not get in eyes, on skin, or on clothing. Prolonged or frequently repeated skin contact may cause allergic reaction in some individuals.”

4) Separate PPE for WPS and Non-WPS uses is not acceptable. Per the acute toxicity review and the RED, the PPE section must be revised to read:

“Some materials that are chemical-resistant to this product are made of barrier laminate, nitrile rubber, neoprene rubber, or viton. If you want more options, follow the instructions for category E on an EPA chemical-resistance category selection chart.

Personal Protective Equipment (PPE):

Pilots must wear:

Long sleeved-shirt and long pants, and  
Shoes and Socks.

Mixers, loaders and all other handlers must wear:

Coveralls over short-sleeved and short pants,  
Socks and chemical-resistant footwear, and  
Chemical-resistant gloves.

In addition to the above PPE, mixers and loaders supporting aerial application to rangelands, pasture lands, or noncropland must also wear:

- a chemical-resistant apron, and
- a NIOSH-approved respirator with a dust/mist filter with MSHA/NIOSH approval number prefix TC-21C or any N, R, P or HE filter.

See engineering controls for additional requirements.”

5) Per the RED, the following text must be added to the engineering control text currently on the label:

“Pilots must use an enclosed cockpit that meets the requirements listed in the WPS for agricultural pesticides [40 CFR 170.240(d)(6)].”

6) Per the RED label table, the text in **bold type** below must be added to the following User Safety Requirements:

“...If no such instructions for washables **exist**, use detergent and hot water...”

7) "This pesticide may be toxic to fish, aquatic invertebrates and aquatic plants," as originally required in the RED is not needed for MCPA and may be deleted from the Environmental Hazard section.

8) Per the RED, the early entry PPE must be revised to read:

- Coveralls,
- Shoes plus socks, and
- Chemical-resistant gloves made of any waterproof material."

9) A Non-Agricultural Use Requirements box must be added to label with the following entry restriction text (that was erroneously omitted from the RED but must be added to the label):

**"NON-AGRICULTURAL USE REQUIREMENTS**

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to product agricultural plants on farms, forests, nurseries, or greenhouses. Do not enter or allow others to enter the treated area until sprays have dried."

10) Per the RED, the spray drift text must be revised to read:

**"SPRAY DRIFT MANAGEMENT**

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment and weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles. Apply only when the wind speed is 2-10 mph at the application site.

*For aerial application:*

The boom length must not exceed 75% of the wingspan or 90% of the rotor blade diameter. Release spray at the lowest height consistent with efficacy and flight safety. Do not release spray at a height greater than 10 feet above the crop canopy.

When applications are made with a crosswind, the swath will be displaced downwind. The applicator must compensate for this displacement at the downwind edge of the application area by adjusting the path of the aircraft upwind.

Do not make applications into temperature inversions.

*For ground boom application:*

"Do not apply with a nozzle height greater than 4 feet above the crop canopy."

11) The following revisions are needed to the Directions for use:

Pasture/Grassland:

Per the RED, the following rate restriction text must be added to the label "Do not apply more than 1.5 lb ae per acre per year. Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days."

Noncropland:

Per the RED, the following rate restriction text must be added to the label "Do not apply more than 3.0 lbs ae per acre per year."

Small grains:

The rate restriction for small grains (barley, rye, wheat and oats) "Do not apply more than 1.75 pints (.8 lb ae) per acre per year" exceeds the rate of 0.75 lb ae per acre per year as specified in the RED. The label must be revised.

12) The MCPA RED erroneously omitted the rate restriction for occupational use to lawns and turfgrass. The rate restrictions appearing on the label for golf courses has to be revised to also apply to lawn and turfgrass.

13) It is unclear if this product is intended for sod farm use. If this product is not intended for sod farm use, the following text should be added to the label:

"Not for use on turf grown for sale or other commercial use as sod, or for research purposes."

14) Assure that the required acid equivalents per acre (lbs ae/A) restrictions are expressed as product volume or product weight per unit area that are in the same units as the registered application rate already on the label. Assure that the maximum application restrictions are identical to the highest currently registered rate for each appropriate application site.

A stamped copy of your labeling is enclosed for your records. Submit one copy of the revised final printed label for the record before you release the product for shipment. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

# MCPA L.V. 4 Ester

FOR THE SELECTIVE CONTROL OF BROADLEAF WEEDS IN FLAX, GRASSES, PASTURES, RANGELANDS, SMALL GRAINS, AND NON-CROP AREAS

<b>ACTIVE INGREDIENT:</b>	
Isooctyl (2-ethylhexyl) Ester of 2-Methyl-4-Chlorophenoxyacetic Acid*	68.7%
<b>OTHER INGREDIENTS:</b>	31.3%
TOTAL:	
	100.0%

Contains petroleum distillates

Isomer Specific by AOAC Method, Equivalent to:  
\*2-Methyl-4-Chlorophenoxyacetic Acid ..... 44.1%, 3.7 lbs./gal.

**KEEP OUT OF REACH OF CHILDREN**  
**WARNING - AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(If you do not understand the label, find someone to explain it to you in detail.)

**SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS**

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300  
For Medical Emergencies Only, Call (877) 325-1840

**ACCEPTED**  
**with COMMENTS**  
**In EPA Letter Dated:**  
**SEP 17 2008**

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

MANUFACTURED BY  
NUFARM AMERICAS INC.  
BURR RIDGE, IL 60527



EPA REG. NO. 228-156  
EPA EST. NO. 228-IL-1

NET CONTENTS

228-156

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**WARNING - AVISO**

May be fatal if absorbed through skin. Harmful if swallowed. Do not get in eyes, on skin, or on clothing.

**PERSONAL PROTECTIVE EQUIPMENT (PPE):**

**NON-WPS USES:** Applicators and other handlers who handle this pesticide for any use NOT covered by the Worker Protection Standard (40 CFR Part 170) -- in general, only agricultural plant uses are covered by the WPS -- must wear: protective clothing and rubber gloves.

**WPS USES:** Personal Protective Equipment - Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category E on an EPA chemical resistance category selection chart.

Applicators and other handlers who handle this pesticide for any use covered by the Worker Protection Standard (40 CFR Part 170) -- in general, agricultural plant uses are covered -- must wear: Long-sleeved shirt and long pants; chemical-resistant gloves when mixing, loading, or using any hand-held equipment, shoes plus socks. Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment (PPE). If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Engineering Controls Statements:** When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

<b>USER SAFETY RECOMMENDATIONS</b>	
<b>Users Should:</b>	
<ul style="list-style-type: none"> <li>• Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.</li> <li>• Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.</li> <li>• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.</li> </ul>	

<b>FIRST AID</b>	
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical treatment information.	
<b>NOTE TO PHYSICIAN</b>	
Probable mucosal damage may contraindicate the use of gastric lavage.	

**ENVIRONMENTAL HAZARDS**

This pesticide may be toxic to fish, aquatic invertebrates and aquatic plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater or rinsate. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

Spray equipment used in applying this product should be thoroughly cleaned before using for any other purpose. Use repeated flushing with soap and warm water or suitable chemical cleaner. It is best to use a separate sprayer for application of insecticides and fungicides. This product will kill or seriously injure many desirable forms of vegetation. Do not apply directly to flowers, fruits, grapes, tomatoes, ornamentals, cotton or other desirable plants. Vapors from this product may injure susceptible plants in the immediate vicinity. Do not apply when weather conditions favor drift from target

area. Avoid use of small-diameter nozzles.(Coarse sprays are less likely to drift.) Excessive amounts of this product in the soil may temporarily inhibit seed germination and plant growth. Most cases of groundwater contamination involving phenoxy herbicides such as MCPA have been associated with mixing/loading and disposal sites. Caution should be exercised when handling MCPA pesticides at such sites to prevent contamination of groundwater.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval (REI). The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls over short-sleeved shirt and short pants, chemical-resistant gloves such as Barrier Laminate, Nitrile Rubber ≥ 14 mils, Neoprene Rubber ≥ 14 mils and Viton ≥ 14 mils; chemical-resistant footwear plus socks, and chemical-resistant headgear for overhead exposure.

**WEEDS CONTROLLED**

This product is effective on a large number of broadleaf weeds. On Mustard, Thistle, and White top or Hoary cress, it has superiority over 2,4-D. The following are some of the weeds killed:

- |                 |                           |                        |                         |
|-----------------|---------------------------|------------------------|-------------------------|
| Beggarticks     | Field bindweed            | Pepperweed             | Sunflower               |
| Burcucumber     | Galinsoga                 | Perennial morningglory | Thornapple              |
| Burdock         | Goatsbeard                | Perennial ragweed      | Tree-of-heaven          |
| Buttercup       | Goldenrod                 | Pigweed                | Vetch                   |
| Canada thistle  | Halbertleaved salt brush  | Poison hemlock         | Whitebrush              |
| Carpetweed      | Hempnettle                | Prickly lettuce        | Wild carrot             |
| Catsear         | Hoary cress               | Puncturevine           | Wild gooseberry         |
| Chervil         | Honeysuckle               | Purslane               | Wild jute               |
| Cocklebur       | Jimsonweed                | Red root               | Wild marigold           |
| Croton          | Knotweed                  | Redstem                | Wild petunia            |
| Daisy           | Kochia                    | Shepherdspurse         | Wild radish             |
| Dandelion       | Lambsquarters (Goosefoot) | Sicklepod              | Wild sage               |
| Dock            | Marshelder                | Small plantain         | Witchweed               |
| Dragonhead mint | Mexican poppy             | Sneezeweed             | Yellow charlock-mustard |
| Dwarf nettle    | Miner's lettuce           | Sowthistle             | Yellow daisy            |
| Fanweed         | Narrow leaf plantain      | Spanish needle         | Yellow rocket           |
| Fat hen         | Nutgrass                  | Stinging nettle        |                         |
| Fennel          | Pennycress                | Stinkweed              |                         |

**SELECTIVE SPRAYING**

**NOTE:** When using on grains and flax - do not forage or graze dairy and meat animals on treated areas within seven days of slaughter. Also, except for small underseeded grains, use at least 8 to 10 gallons of water per acre for ground application and at least 3 to 5 gallons of water per acre for aerial application.

**SMALL GRAINS:** Barley, Rye, Wheat (For small grains underseeded with legumes, see the small grains paragraph below.) Spring planted - not seeded to legumes. Apply as a water mix spray by ground sprayer or airplane. Use 0.5 to 1.5 pints of this product per acre for the more susceptible weeds and up to 1.75 pints per acre for less susceptible weeds. Treat when fully tillered grains have reached the 3 to 4-leaf stage (8 to 10 inches tall). Do not spray grain in the boot to dough stage.

Fall planted - Fall application - not seeded to legumes. The safest time to treat is when fall planted small grains are fully tillered and have reached the 3 to 4-leaf to the boot stage (8 to 10 inches tall). Infestations of broadleaf weeds germinating in fall and winter may, however, justify treatment at an earlier stage. Fall planted barley, rye or wheat can be sprayed with 1/2 to 1 pint of this product per acre from the 3 to 4-leaf stage to the boot stage.

Spring application - Apply in early spring when annual broadleaf weeds are small and grain is fully tillered but before grain is in the jointed stage

**SMALL GRAINS - Underseeded with Alfalfa, Birdsfoot Trefoil, Lespedeza, Red and White Clover:** For emergency control of serious infestations of Mustard, Yellow rocket, and other susceptible broadleaf weeds, apply 0.25 to 0.5 pint in not more than 5 to 10 gallons of water per acre. Make application after cereal is well tillered (4 to 8 inches tall) and before reaching boot stage. The nurse crop and weeds should provide a protecting canopy which, together with the use of low gallonage applied at low pressure, will reduce the risk of damage to the legumes. Do not apply to small grains underseeded with vetch or sweet clover, which are very susceptible. There is also some risk to other legumes if only thinly protected by a canopy.

**RESTRICTIONS AND LIMITATIONS FOR USE ON SMALL GRAINS**

Do not apply more than 1.75 pints per acre per year.

**OATS:** Spray after oats are fully tillered (4 to 6 inches tall) but before they are in the boot to dough stage. Apply the lower rate of 1/2 pint where annual weeds are the problem, the higher rate of 1 pint for perennial weeds.

**RESTRICTIONS AND LIMITATIONS FOR USE ON OATS**

Do not apply more than 1.75 pint per acre per year.

**FLAX:** Use 0.25 to 0.5 pint per acre. Use lower rate where susceptible weeds such as Mustard are the main problem. Use higher rate when weeds such as Cocklebur, Lambsquarters, Pigweed, Ragweed and Stinkweed are the problem. Apply only when weeds are up and when flax is 2 to 8 inches high and before it comes into bud. Treatment after early bud stage may result in severe damage. If Canada thistle is present, it may be necessary to go as high as 0.5 pint per acre to prevent seed head production. Some injury to flax may result.

**RESTRICTIONS AND LIMITATIONS FOR USE ON FLAX**

Do not apply more than 0.5 pint per acre per year.

**RANGE AND TIMBERLANDS:** For control of Whitebrush - Use 1-1/4 quarts in mixture of 1 gallon of diesel oil and sufficient water to make 8 gallons of solution per acre. Apply in spring or fall under good moisture conditions, full leaf, before blossoms begin to fall.

**GRASSES:**

**Established Golf Courses, Lawns and similar Turf Grasses:** Use 1 to 3 pints per acre in 10 to 100 gallons of water to give thorough coverage. Use higher rate for White top and Canada thistle. On smaller areas, use 1 fluid ounce (4 tablespoonfuls) mixed in 3 to 5 gallons of water and apply uniformly over 2,000 square feet. Do not mow within 2 days before or after application. Spring and fall are best times to treat. Do not use on lawns of creeping grasses, such as bent, except for spot spraying, nor on freshly-seeded turf until the grass has become well established.

**RESTRICTIONS AND LIMITATIONS FOR USE ON GOLF COURSES**

Do not apply more than 3.25 pints per acre per year. Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

**Established Grassland and Pastures:** Use 1 to 3 pints per acre in sufficient water (10 to 100 gallons) in airplane or ground spray application and give thorough coverage. Use higher rate for Canada thistle, White top and other less easy-to-kill weeds. Spray perennials in early bud to full bloom stage and regrowth in fall; other weeds in spring or fall, when actively growing.

**Grasses Grown for Seed:** Use 1 to 2 pints per acre in 5 to 50 gallons of water by air or ground sprayer application. Use higher rate where weed stands are heavy. In established grasses, apply in spring before head comes into boot stage and on seedling grass after grass has tillered.

**NOTE:** For weed control in grasses, repeat treatment may be needed for less susceptible weeds. White clover and other legumes may be temporarily injured or killed. In some areas, bent, buffalo, carpet, centipede, dichondra and St. Augustine may also be injured by the treatment.

**RESTRICTIONS AND LIMITATIONS FOR USE ON GRASS GROWN FOR SEED**

Do not apply more than 3.25 pints per acre per year. Do not apply more than 2 applications per year with a minimum retreatment interval of 21 days.

**NON-CROP SPRAYING:** Canada thistle, Meadow buttercup and White top. (In Non-Crop Areas such as Fence rows, Rights-of-way, Roadsides and similar places): For spot treatment use 1/4 pint to 3 to 4 gallons of water or 3/4 gallon per acre in 12 to 20 gallons of water to give coverage for most extensive areas. Spray to wet weeds thoroughly when in bud to early bloom and again on fall regrowth.

**SPRAY DRIFT**

Avoiding spray drift at the application site is the responsibility of the applicator. The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

Apply only as a medium or coarser spray (ASAE standard 572) or a volume mean diameter of 300 microns or greater for spinning atomizer nozzles.

Apply only when the wind speed is 2 to 10 mph at the application site.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Always store pesticides in a secured warehouse or storage building. Do not store near open containers of fertilizer, seed or other pesticides. Containers should be opened in well-ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities.

**WARRANTY DISCLAIMER**

The directions for use of this product must be followed carefully. THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER. TO THE EXTENT PERMITTED BY LAW, MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR

9/a.

**ANY PARTICULAR TRADE USAGE.** UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO CROP INJURY OR INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

**LIMITATION OF LIABILITY**

TO THE EXTENT PERMITTED BY LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL. MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BY WAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS.

**If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.**

(RV092906)