

Riverdale

Pre-Emergent Crabgrass Control Concentrate

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid breathing dust or spray mist. Avoid contact with skin, eyes, or clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Use with care when applying in areas adjacent to any body of water. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of waste.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Do not apply when weather conditions favor runoff from target area.

Read the following carefully before application. Use this product only for recommended purposes and at recommended dosages.

For use on ESTABLISHED turfs listed below which are tolerant to the herbicide in this product when applied as directed. See Precautions.

Bluegrass (Perennial)	Bermudagrass
Ryegrass (Perennial)	Centipedegrass
Fescue (Except Fine Leaved)	St. Augustinegrass
Bahiagrass	Zoysiagrass

WHEN TO APPLY:

Apply to established turf before germination of annual weeds listed below. The general times of application are covered below. The precise time of application for a specific area will vary significantly because of climate and individual cultural practices. Consult State Agricultural Experiment Station Weed Specialists or local Turf Authorities for any additional information regarding specific weed germination dates in your area.

APPLICATION DIRECTIONS:

Application Equipment

This product is to be mixed with water and applied as a spray. Apply in 50 to 250 gallons of water per acre (broadcast basis) using any properly calibrated spray unit.

Weeds Controlled

This product will control annual bluegrass (Poa Annual), smooth crabgrass, hairy crabgrass, goosegrass, (silver crabgrass or crowfoot), water grass (barnyard grass), yellow and green foxtail. It will not control established weeds.

ACTIVE INGREDIENT:

N-Butyl-N-ethyl-alpha, alpha, alpha, trifluoro-2,6, dintro-p-toluidine 10%

INERT INGREDIENTS 90%

TOTAL 100%

MINIMUM GUARANTEED ANALYSIS

Total Nitrogen (N) 34%

ACCEPTED

JUL 23 1982

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 228-158

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger or, if available, by administering syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air.

IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash with plenty of soap and water.

NET WEIGHT 20 POUNDS

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY
CHICAGO HEIGHTS, ILLINOIS

DIRECTIONS FOR USE — continued

APPLICATION RATES:

Southern (warm season) grasses

For pre-emergent control of annual bluegrass (Poa Annual) apply in late summer to early fall, prior to annual bluegrass seed germination. Additional applications at the same rate can be made 4 to 6 months later in early spring, prior to soil warming and subsequent germination, in areas heavily infested in thin turf areas.

For pre-emergent control of crabgrass, foxtail, barnyard grass and goosegrass apply in late winter or early spring prior to annual weed grass germination. A second application may be made 4 to 6 months after initial treatment for continued control of annual weeds which may germinate into the summer or early fall.

AMOUNT TO APPLY:

Use 0.69 lbs. per 1,000 sq. ft. or 30 pounds per acre (equivalent to 3 pounds of active ingredient per acre). Mix with sufficient water for adequate coverage.

APPLICATION RATES:

Northern (cool season) grasses.

For pre-emergent control of annual bluegrass (Poa Annual), apply in late summer to early fall (mid-July to September) or in early spring prior to annual bluegrass germination.

For pre-emergent control of crabgrass, foxtail, barnyard grass and goosegrass, apply in early spring prior to annual weed grass germination.

AMOUNT TO APPLY:

Use 0.46 pounds per 1,000 sq. ft. or 20 pounds per acre (equivalent to 2 lbs. active ingredient per acre). Mix with sufficient water for adequate coverage.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL

Pesticide, spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or Local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL

Dispose of bags according to approved Federal, State or Local procedures under the Resource Conservation and Recovery Act.

PM 25

228-158

7/23/82

PM 181