

228-160

02/25/2003

1/4

RIVERDALE®

3 PLUS 3 AMINE
TURF HERBICIDE

Extra Copy

A SELECTIVE HERBICIDE TO CONTROL BROADLEAF WEEDS

GET THE OPTICAL ADVANTAGE®

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	33.95%
Dimethylamine Salt of (+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**†	17.06%

INERT INGREDIENTS:.....	48.99%
TOTAL	100.00%

Isomer specific Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid	28.2%, 2.70 lbs./gal.
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid.....	14.1%, 1.35 lbs./gal.

†CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p.

Riverdale and The Optical Advantage are Registered Trademarks of Riverdale - A Nufarm Company

KEEP OUT OF REACH OF CHILDREN

CAUTION - CAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.

SEE SIDE PANELS FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS


EPA REG. NO. 228-160

NET CONTENTS

GALS.

EPA EST. NO. 228-IL-1

MANUFACTURED BY NUFARM AMERICAS INC., BURR RIDGE, ILLINOIS 60527-0866



NOTE: Spanish language is optional

(Side Panels)

STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION - CAUCION**

Do not get in eyes. Harmful if swallowed or absorbed through skin. Avoid inhalation of vapors and contact with skin and clothing. Wear face shield or goggles, long pants, long-sleeved shirt, socks, shoes and rubber gloves when mixing, loading and applying this product. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron. If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES: Hold eyelids open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates or nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. When cleaning equipment, do not pour washwaters on the ground; spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes.

Most cases of groundwater contamination involving phenoxy herbicides such as MCPP and 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

GENERAL INFORMATION

3 Plus 3 Amine is a post-emergence herbicide especially formulated for use on turfgrass for selective control of many broadleaf weeds commonly growing in lawns without injury to desirable grasses. This product may be used on such types of turf as Marion, Kentucky bluegrass, Bermudagrass, fescues and rye grasses. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

USE PRECAUTIONS

CARE MUST BE TAKEN TO AVOID DRIFT OF SPRAY MISTS TO NON-TARGET susceptible plants such as cotton, vegetables, flowers, ornamentals, shrubs, trees, and other desirable plants. Do not spray near these plants. Do not spray on buffalo, carpetgrass, dichondra, St. Augustine, nor on grasses where desirable clovers are present. This product has been used successfully on bentgrass; however, it should not be used unless prior tests have shown it to be non-damaging under the particular conditions of use. To eradicate broadleaf weeds, it is better to use another selective herbicide for fine lawn grasses. Apply in Spring or Fall when weeds are actively growing. Use a lawn type sprayer with coarse spray as wind drift is less likely. Avoid fine mists. Do not spray exposed roots of desirable plants. Use the higher rate for hard-to-kill weeds. Do not exceed specified application dosages for any area. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 90°F. Avoid applying during excessively dry or hot periods unless irrigation is used. For optimum results, (1) turf should not be mowed 1 to 2 days before and following application; (2) Do not apply if rainfall is expected within 48 hours; nor should lawns be irrigated (watered) for 48 hours following application. Reseed no sooner than 3 to 4 weeks after application of this product.

WEEDS CONTROLLED BY 3 PLUS 3 AMINE

The following is a partial list of weeds controlled:

Buckhorn	Lambsquarters	Red clover
Clover	Mallow	*Red sorrel
Common chickweed	Mouseear chickweed	Stitchwort
Curley dock	Morningglory	*Speedwell
*Dandelion	Pigweed	White clover
*English daisy	Plantain	*Wild garlic
Ground ivy	Purslane	*Wild onion
Knotweed	Ragweed	Yarrow

Use high rate on chart for weeds marked with an asterisk () in the above list. Knotweed is controlled more easily in Spring or early Summer when weeds are young.

NOTE: MCPP-p acts more slowly than 2,4-D so that some weeds will be controlled before others. Allow 3 to 4 weeks for complete weed control action to occur.

<u>Amount to Apply</u>	<u>Per 1,000 Sq. Ft.</u>	<u>Per Acre</u>
Low rate	1-1/4 fl. ozs.	1-2/3 qt.
High rate	1-1/2 fl. ozs.	2 qts.

PREPARATION OF THE SPRAY: This product must be mixed with water prior to use. For best results use 2 to 4 gallons of water per 1,000 sq. ft. for smaller areas; 20 to 40 gallons of water per acre for larger areas. Apply by means of a sprayer to give uniform coverage. Repeat applications may be necessary especially with heavy infestations of the less susceptible weeds indicated with an asterisk in above list. 2 days before or after application, it is wise to properly fertilize the lawn so as to encourage the lawn to heal as rapidly as possible, filling in the bare areas where the weeds have died away. Once the weeds have been removed, the bare areas should be reseeded.

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near open containers of fertilizer, seed, or other pesticide. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well-ventilated area. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate ground water. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and local authorities by burning. If burned, stay out of smoke.

WARRANTY

Nufarm warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixtures with other chemicals, and factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use. (RV 013103)