221-163

Please read instructions on reverse before completing form. Form Approved. OMB No. 2070-0060. Approval expires 11-30-93 OPP Identifier Number United States Environmental Protection Agency Registration Office of Pesticide Programs (H7505C) **Amendment** Washington, DC 20460 165885 · **Application for Pesticide:** Other Section I 3. Proposed Classification 2. EPA Product Manager 1. Company/Product Number n 228–163 Joanne I. Miller Restricted > 1 None 4. Company/Product (Name) Riverdale Premium Weed & Feed 6. Expedited Review. In accordance with FIFRA Section 3(c)(3) 5. Name and Address of Applicant (Include ZIP Code) (b)(i), my product is similar or identical in composition and labeling RIVERDALE CHEMICAL COMPANY 425 West 194th Street 60425-1584 Glenwood, IL EPA Reg. No. 172175 Check if this is a new address. Product Name Section I i Final printed labels in response to Amendment - Explain below Agency letter dated_ Resubmission in response to Agency letter dated_ "Me Too" Application. Notification - Explain below. Other - explain below. Explanation: Use additional page(s) if necessary. (For section I and Section II.) To comply with PR Notice 93-3 Section III -1. Material This Product Will Be Packaged in: Child-Resistant Packaging Unit Packaging 2. Type of Container Water Soluble Packaging Metal Yes* Yes . Yes **Plastic** Glass Paper lf "Yes," Unit Package wgt. , No. per. Other (Specify container Package wgt. , , , , container Certification must be submitted. 5. Location of Label Directions 3. Location of Net Contents Information 4. Size(s) of Retail Container __ Container On Labeling accompanying groducing pagitaming ingrésis que 6. Manner In Which Label Is Affixed To Product Lithograph (2017) Paper glued ... Stenciled as year Section IValence by reside 1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application of Telephone No. (Include Area Code) Russell F. Sawyer Regulatory Affairs Manage: 1 2 708/754-3330 Certification Carter to the Transfer of 6. Date Application Received I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or (Stamped) both under applicable law. 2. Signature Regulatory Affairs Manager 4. Typed Name 5. Date

May 24.

1993

Russell F. Sawyer

(Non-Commercial Label)

RIVERDALE

PREMIUM WEED & FEED X - X - X

FOR LAWNS AND ORNAMENTAL TURF TREATS (OR COVERS) 5,000 SQ. FT. (50' x 100')

This weed and feed product is a remarkably effective lawn fertilizer which combines fast and long-lasting feeding action, plus effective weed control. This product produces a thick, green turf while killing weeds to make way for new turf growth.

Riverdale Premium Weed & Feed is designed for the control of dandelion, clover, and many other broadleaf weeds. Weeds contacted by this product gradually die and disappear from the lawn. See back panel

for a list of weeds controlled by Riverdale Weed & Feed.
Formulated for your area: Nitrogen to help promote green thick lawns. Includes slow-release nitrogen for extended lawn feeding; Phosphoric Acid to help provide strong, vigorous roots; and, Potash to help increase resistance to disease and drought.

ACTIVE INGRELIENTS:

| Isooctyl (2-ethylhexyl | Ester of 2,4-Dichlorophenoxyacetic Acid* 1.35% | |
|------------------------|--|--|
| Isooctyl (2-ethylhexyl |) Ester of 2-(2-Methyl-4-Chlorophenoxy) | |
| propionic Acid** | 1.37% | |

TOTAL 100.00%

Isomer Specific AOAC, Equivalent to:

GUARANTEED ANALYSIS: °

XZ Urea Nitrogen

Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated Urea, Urea, and Muriate of Potash

Chlorine (C1) Maximum X%

The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

NET WEIGHT - 15 LBS.

EPA REG NO. 228-163

EPA EST. NO. 228-IL-1

MANUFACTURED BY
RIVERDALE CHEMICAL COMPANY

Revised A/O 3/9/93

LAPTE TO THE TYPE PER TON TO THE PER TON TO THE PER TON TO THE PER TO THE PER

ATÉ ____ --

(Back Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Avoid inhalation of dust and contact with eyes, skin, or clothing. Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Get medical attention.

IF IN EYES: Flush eyes with water for 15 minutes. Get medical attention.

IF ON SKIN: Wash skin with soap and water.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

This product may be used on Bluegrass, Fescue, Rye, common Bermuda, Bahia and Zoysia lawns. Do not use on dichondra, carpetgrass and St. Augustine grasses. Do not use this product where desirable clovers are present or allow the product to "wind drift" or flowers, vegetables or ornamental shrubbery. When using a spinner type (rotary) spreader, use extreme caution to avoid any contact with desirable plants.

This product may be used for control of broadleaf weeds in lawns such as: Bindweed, Black medic, Blue lettuce, Buckhorn, Burdock, Chickweed, Chicory, Cinquefoil, Clover, Cocklebur, Croton, Dandelion, Dock, Fanweed, Galinsoga, Goldenrod, Ground ivy, Hawkweed, Healall, Henbit, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarters, Mallow, Morningglory, Mustard, Nettles, Oxalis, Pennywort, Peppergrass, Pigweed, Plantain, Povertyweed, Purslane, Ragweed, Shepherdspurse, Smartweed, Speedwell, Spotted spurge, Thistle, Vervain, Vetch, Wild aster, Wild lettuce, Wild radish and Yarrow.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Apply when grass is moist (after a rain, sprinkling or dew). For most effective broadleaf weed killing action, avoid unnecessary disturbances, including

watering, of treated areas for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours following treatment. Avoid contact with desirable shrubs, plants, regetables, or flowers. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. Do not cut grass 1 to 2 days before and after application.

HOW TO APPLY: This 15 pound bag should be used on 5000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is equal to one application.

The following settings are approximate for applying contents of this package at a suggested rate of 3.0 pounds per 1000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 sq. ft. and adjust to apply at a rate of .75 lb. per 250 sq. ft. Clean spreader thoroughly after use.

SPREADER SETTINGS For One Pass Application°

| SETTING | MAKE | SETTING |
|---------|------------------|---|
| 3-1/2 | Central | 5 |
| 4 | K-Mart | 5 |
| E | Jackson Drop | 5 |
| 5 | Jet Spray | 3-1/2 |
| 5 | Sears Rotary | 3 |
| | SETTING 3-1/2 | 3-1/2 Central 4 K-Mart E Jackson Drop 5 Jet Spray |

Sweep all fertilizer materials from concrete surfaces immediately after application. Do not wash material from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.

STORAGE AND DISPOSAL

STORAGE: Store in ...ginal container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near other feed or foodstuffs. To prevent cross-contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides or seeds.

PRODUCT DISPOSAL: To dispose of bag which contains unused product, securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discardin trash.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(ERP 061594) (PRO31 042194)

The following note will not be part of the actual label copy.

*NOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers.

(Commercial Paper)

RIVERDALE

PREMIUM WEED & FEED X - X - X

FOR LAWNS AND ORNAMENTAL TURF TREATS (OR COVERS) 5,000 SQ. FT. (50' x 100')

This weed and feed product is a remarkably effective lawn fertilizer which combines fast and long-lasting feeding action, plus effective weed control. This product produces a thick, green turf while killing weeds to make way for new turf growth.

Riverdale Premium Weed & Feed is designed for the control of dandelion, clover, and many other broadleaf weeds. Weeds contacted by this product gradually die and disappear from the lawn. See back panel

for a list of weeds controlled by Riverdale Weed & reed.
Formulated for your area: Nitrogen to help promote green thick lawns. Includes slow-release nitrogen for extended lawn feeding; Phosphoric Acid to help provide strong, vigorous roots; and, Potash to help increase resistance to disease and drought.

ACTIVE INGREDIENTS:

| Isooctyl (2-ethylhexyl) Ester of 2,4-Dichlorophenoxyacetic Acid* 1.35% |
|--|
| Isooctyl (2-ethylhexyl) Ester of 2-(2-Methyl-4-Chlorophenoxy) |
| propionic Acid** 1.37% |
| INERT INGREDIENTS: |
| TOTAL 100.00% |

Isomer Specific AOAC, Equivalent to:

| *2,4-Dichlorophenoxyacetic Acid | 0.90% |
|--|-------|
| **2-(2-Methyl-4-Chlorophenoxy)propionic Acid | 0.90% |

GUARANTEED ANALYSIS: °

| Total Nitrogen (N) | XZ. |
|--|------------|
| XZ Ammoniacal Nitrogen | |
| XX Urea Nitrogen | |
| Available Phosphoric Acid (P ₂ O ₅) | XZ |
| Soluble Potash (K ₂ O) | X % |

Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated Urea, Urea, and Muriate of Potash

Chlorine (C1) Maximum XX

The urea nitrogen in this product has been coated to provide XX coated slow release urea nitrogen.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
AND STATEMENT OF PRACTICAL TREATMENT

NET WEIGHT - 15 LBS.

EPA REG NO. 228-163

EPA EST. NO. 228-IL-1

NOTIFICATION
LAPT MOTOR PER FH NO 1993
DATE JUN 10 1993

Revised A/O 3/9/93

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY GLENWOOD, ILLINOIS 60425-1584

(Back Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Avoid inhalation of dust and contact with eyes, skin, or clothing. Wear long pants, long-sleeved shirt, socks, shoes and gloves when loading and applying this product. After using this product remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast application per treatment is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Get medical attention.

IF IN EYES: Flush eyes with water for 15 minutes. Get medica! attention.

IF ON SKIN: Wash skin with soap and water.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters, nor pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and MCPP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

This product may be used on Bluegrass, Fescue, Rye, common Bermuda, Bahia and Zoysia lawns. Do not use on dichondra, carpetgrass and St. Augustine grasses. Do not use this product where desirable clovers are present or allow the product to "wind drift"

on flowers, vegetables or ornamental shrubbery. When using a spinner type (rotary) spreader, use extreme caution to avoid any contact with desirable plants.

This product may be used for control of broadleaf weeds in lawns such as: Bindweed, Black medic, Blue lettuce, Buckhorn, Burdock, Chickweed, Chicory, Cinquefoil, Clover, Cocklebur, Croton, Dandelion, Dock, Fanweed, Galinsoga, Goldenrod, Ground ivy, Hawkweed, Healall, Henbit, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarters, Mallow, Morningglory, Mustard, Nettles, Oxalis, Pennywort, Peppergrass, Pigweed, Plantain, Povertyweed, Purslane, Ragweed, Shepherdspurse, Smartweed, Speedwell, Spotted spurge, Thistle, Vervain, Vetch, Wild aster, Wild lettuce, Wild radish and Yarrow.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Apply when grass is moist (after a rain, sprinkling or dew). For most effective broadleaf weed killing action, avoid unnecessary disturbances, including watering, of treated areas for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours following treatment. Avoid contact with desirable shrubs, plants, vegetables, or flowers. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. Do not cut grass 1 to 2 days before and after application.

HOW TO APPLY: This 15 pound bag should be used on 5000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is equal to one application.

The following settings are approximate for applying contents of this package at a suggested rate of 3.0 pounds per 1000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 sq. ft. and adjust to apply at a rate of .75 lb. per 250 sq. ft. Clean spreader thoroughly after use.

SPREADER SETTINGS For One Pass Application°

| MAKE | SETTING | MAKE | SETTING | |
|---------------|---------|--------------|-----------|-------------|
| Cyclone | 3-1/2 | Central | 5 • • ; | • |
| Scotts Drop | 4 | K-Mart | 5. | * * * * * * |
| Scotts Rotary | E | Jackson Drop | 5 | *** |
| Show-N-Tell | 5 | Jet Spray | 3-1/2 *** | 42:11 |
| Sears Drop | 5 | Sears Rotary | 3 • • • • | • |

Sweep all fertilizer materials from concrete surfaces immediately after application. ... Do not wash material from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near seeds, fertilizers, insecticides or fungicides. Do not stack more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

PRODUCT DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay away from smoke.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(ERP 061594) (PR933 042194)

The following note will not be part of the actual label copy.

*NOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers.