

RIVERDALE

WEDESTROY

WEED AND FEED

This product is designed for the control of dandelion and many other broadleaf weeds. See back panel for a complete list of weeds controlled by this product.

ACTIVE INGREDIENT:	By Weight
*Isooctyl Ester of 2,4-Dichlorophenoxy-acetic Acid.....	1.50%
INERT INGREDIENTS.....	98.50%
<i>0.99493 OK</i>	Total.... 100.00%
*Equivalent to 1.00% 2,4-Dichlorophenoxyacetic Acid Isomer Specific by AOAC Method 6.D01-5	

1.50975 OK

ACCEPTED
 SEP 13 1982
 228-164

MINIMUM GUARANTEED ANALYSIS

Total Nitrogen	10.00%
Available Phosphoric Acid (P ₂ O ₅)	6.00%
Soluble Potash (K ₂ O)	4.00%

KEEP OUT OF REACH OF CHILDREN
 CAUTION
 SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

MANUFACTURED BY
 RIVERDALE CHEMICAL COMPANY
 Chicago Heights, Illinois

Net Weight - 15 lbs.
 Treats 5,000 Sq.Ft. (50x100)

EPA Reg. No. 228-

EPA EST No. 228-IL-1

BEST DOCUMENT AVAILABLE

E

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS (& DOMESTIC ANIMALS)

CAUTION

Avoid contact with skin, eyes, or clothing.
Avoid breathing dust.
Wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT
In case of contact wash skin with soap and water; for eyes
flush with water for 15 minutes and get medical attention.

ENVIRONMENTAL HAZARDS
Keep out of lakes, streams, or ponds. Do not apply when weather
conditions favor drift from target area. Do not contaminate
water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

READ THE FOLLOWING CAREFULLY BEFORE APPLICATION

Use this product only for recommended purposes and dosage.

GENERAL INFORMATION

NOTE: Do not apply on Lippia, St. Augustine, Bentgrass or Dichondra. When properly used this product will not burn turf.

Do not re-seed for at least two weeks after application. Do not apply to a newly seeded lawn until after 3 mowings.

This product is designed to promote lush green growth and control many broadleaf weeds in lawns, such as:

Bahia	Fanweed	Lambsquarters	Pigweed
Bindweed	Goldenrod	Mallow	Plantain
Burkhorn	Ground Ivy	Morning Glory	Purslane
Burdock	Heal All	Mustard	Ragweed
Chickweed	Henbit	Nettles	Smartweed
Chicory	Jewelweed	Oxalis	Speedwell
Cocklebur	Jimson Weed	Pennywort	Thistle
Dandelion	Kochia	Pepper Grass	Yarrow

And many other weeds susceptible to 2,4-D

WHEN TO APPLY: For best results apply when weeds are young and actively growing.

Applications made when the grass is wet from dew, rain, or sprinkling will be more effective.

Do not water for at least 24 hours after application. If rain occurs during 24 hours after application, a second application may be necessary.

HOW TO APPLY: Use a fertilizer spreader. The following settings are approximate for applying contents of this package at suggested rate of 4 lbs. per 1000 sq. ft. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed.

SPREADER SETTINGS:	MAKE	SETTING
	Cyclone	5 $\frac{1}{4}$ - 5 $\frac{3}{4}$
	Scott's 35	5 $\frac{1}{2}$
	Scott's	C-1 J-1
	Central	7-8
	K-Mart	7-8
	Montgomery Ward's	6 $\frac{1}{2}$

DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.