

RIVERDALE 2D + 2DP LOW VOL

with comments

ACTIVE INGREDIENTS:

*Isooctyl Ester of 2,4-dichlorophenoxyacetic Acid....	32.8%
**Isooctyl Ester of 2-(2,4-dichlorophenoxy) propionic Acid.....	2.8%

228-167

INERT INGREDIENTS:..... 25.7%

TOTAL..... 100.00%

*2,4-dichlorophenoxyacetic Acid Equivalent 21.8%, 1.8 lbs./Gallon

**2-(2,4-dichlorophenoxy) propionic Acid Equivalent 2.9%, 1.87 lbs./Gallon.

** Isomer Specific by AOAC Method.

CAUTION

KEEP OUT OF REACH OF CHILDREN.

See Side Panel for Additional
Precautionary statements.

ONLY FOR SALE TO, USE AND STORAGE BY LAWN CARE, LANDSCAPING PERSONNEL, AGRICULTURAL, PROFESSIONAL AND SERVICE PERSONS.

MANUFACTURED BY
RIVERDALE CHEMICAL COMPANY
Chicago, Ill., Illinois

E.P.A. Reg. No. 225-1

E.P.A. Est. No. 228-IL-1

Net Contents _____ Gal.

(2)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION:

Harmful if swallowed or absorbed through the skin. Avoid contact with eyes, skin or clothing.

Avoid breathing spray mist.

In case of contact, immediately flush eyes with plenty of water for at least 15 minutes. Seek medical attention. Wash exposed skin gently with plenty of soap and water.

If swallowed, DO NOT induce vomiting. If vomiting occurs spontaneously, keep away clean. Never give anything by mouth to an unconscious person. Seek medical attention.

ENVIRONMENTAL HAZARDS:

DO NOT apply directly to water. DO NOT contaminate water by cleaning of equipment or disposal of wastes. DO NOT contaminate irrigation ditches or water used for domestic purposes. This product is toxic to fish. Use care to avoid spray contact or drift to 2,4-D susceptible plants such as cotton, tomatoes, flowers, grapes, fruit trees and ornamentals. DO NOT permit spray mist containing this product to drift onto them, since even very small quantities of the spray, which may not be visible, can cause severe injury during both growing and dormant periods. DO NOT spray when the wind is blowing towards susceptible crops or ornamental plants. Use coarse sprays to minimize drift. Spray drift can be lessened by keeping the spray boom as low as possible by applying 20 gallons or more spray per acre: by using no more than 20 pounds spraying pressure with flat fan or flooding flat fan nozzle tips; by spraying when wind velocity is low; and by stopping all spraying when wind exceeds 6 to 7 miles per hour. DO NOT apply with hollow cone-type insecticide or other nozzles that produce a fine droplet spray. Although this product is much less volatile than butyl or isopropyl esters, at high temperatures (above 95°F) vapors from this product may injure susceptible plants growing nearby. DO NOT use in a greenhouse. Flush sprayer out on suitable non-crop area after use. DO NOT use the same spray equipment for applying other materials to 2,4-D susceptible crops as injury may result.

NOTE: DO NOT graze dairy animals on treated areas within 6 weeks after application.

DO NOT graze meat animals on treated areas with 2 weeks of slaughter.

Read entire label before using this product.

GENERAL INFORMATION:

Lawns and other ornamental turf grass areas: Riverdale 20 + 2DP
Low Vol is recommended for control of broadleaf weeds in lawns and

similar turf areas. This treatment may injure bentgrass., St. Augustinegrass, centipede grass, carpetgrass and newly seeded lawns. If necessary to control weeds in such turf, use half the recommended rate in chart and repeat application in 2 to 3 weeks. Do not use on bentgrass greens or tees.

The following is a partial list of weeds controlled in turf by RIVERDALE 20 + 20P LOW VOL.

MORE SUSCEPTIBLE WEEDS:

- | | | |
|---------------------------------|----------------|---|
| Black medic | Dandelion | Little starwort |
| Buttercup | *English daisy | Oxalis |
| Chickweeds
(common mouseear) | Ground ivy | (wood sorrel) |
| ✓ Clovers | Healall | Plantains (narrow or buckhorn: broadleaf) |
| Cudweed | Henbit | *Spurge |
| | Knotweed | |
- *Treat in spring and again in fall.

LESS SUCCEPTIBLE WEEDS:

- | | | |
|-----------------|---------------|--------------|
| Bindweed | Kochia | Vervain |
| Dock | Lambsquarters | Vetch |
| Eveningprimrose | Mallow | Violet |
| Falsedandelion | Pigweed | Wild carrot |
| Fleabane | Poorjoe | Wild Lettuce |
| Florida pusley | Ragweed | |

This product is NOT effective on perennial veronicas or on weed grasses.

PASTURE, RANGELAND,

Brush Control on Utility Rights-of-Way, along highways and Solid Stands of Oak or Elm: RIVERDALE 20 + 20P LOW VOL is specifically designed for utilities and other industrial users to control woody plants. This product also controls many noxious perennial weeds on uncropped land, such as along highways and drainage ditchbanks. The following is a partial list of the weeds controlled:

- | | | |
|---------------|------------------------------------|----------------|
| Alder | Fir | Red elm |
| Ash | Gooseberry | Red maple |
| Aspen | Hemlock | Salmonberry |
| Birch | Greenbrier | Sand Sagebrush |
| Blackberry | Gum | Serviceberry |
| Black cherry | Honeysuckle | Shinnery oak |
| Blackjack oak | Locust | Snowberry |
| Black locust | Manzanita | Spruce |
| Box elder | Maple | Sumac |
| Brambles | Oak | Sycamore |
| Buckbrush | Osage orange | Tulip poplar |
| Elderberry | Palmetto | Wild cherry |
| Elm | Pine | Wild grape |
| Ceanothus | Poison ivy | Willow |
| Chamise | Poison oak | Winged elm |
| Coffeeberry | Poplar | Yerba santa |
| Currant | Red elder (and many other species) | |

4

NOTE: Local conditions and application regulations vary and may affect use of this herbicide. Consult local agricultural experiment station or extension service weed specialists and state regulatory agencies for recommendations in your area.

NOTE: When stored at temperatures below freezing, it may be necessary to warm contents to 45°F and mix thoroughly before using.

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

PREPARATION OF THE SPRAY: Fill the spray tank with half the required amount of water, then add the recommended amount of Riverdale 20 + 2DP Low Vol herbicide anytime broadleaf weeds are growing actively. Dandelion, plantain, wood sorrel and clovers are best treated in the fall or in spring before flower heads develop. Winter weeds such as chickweeds and henbit should be treated in early spring.

LAWNS: Summer weeds such as oxalis, knotweed and sparges should be sprayed when they are small. In areas with extended growing seasons, such as California, treatment in both spring and fall may be needed to control more resistant species.

Lawns to be over-seeded in the fall should be treated at least 4 weeks before the planned seeding date. Rainfall (1/4 inch or more) or irrigation should follow treatments. Fall-seeded lawns may be treated the following spring.

Spring-seeded lawns may be treated after the grasses have sprouted and been cut at least twice, generally 6 to 10 weeks after seeding, depending on germination and growth rate.

For established lawns, fall treatment fits into a good turf management program. Proper fertilization and mowing should be combined with chemical weed control to thicken the turf after weeds have died and to discourage more weeds from invading.

HOSE ATTACHED SPRAYERS (LAWNS MODELS): Use 4 tablespoons of Riverdale 20 + 2DP Low Vol herbicide diluted with water to the 3 to 5 gallon mark on the sprayer jar for each 500 square feet (25'x20') of lawn area to be sprayed. Adjust water pressure so that spray streams about 10-15 feet with no misting.

AMOUNTS TO USE:

RIVERDALE 20 + 2DP Low Vol

Herbicide	Area Covered	Water Volume
3 oz.	1,000 sq. ft.	1 gal.
1 pt.	5,000 sq. ft.	5 gals.
1 qt.	10,000 sq. ft.	10 gals.
1 gal.	40,000 sq. ft.	40 gals.

RIVERDALE 20 + 2DP may be used as a woody plant herbicide for control of many noxious perennial weeds on non-cropland such as along highways and drainage ditches.

*Riverdale
20 + 2DP
herbicide*

FOLIAGE STEM TREATMENT:

This is the standard method for high-volume sprays along fence rows, highways and rights-of-way. Use as a first-spray" on thick brush composed of mixed species. Apply to both stems and foliage from the time foliage is completely matured until the plants start to go dormant.

All leaves, stems and suckers must be completely wet to the ground line for effective control. Some regrowth may be anticipated on the more resistant species, such as oak, maple and ash. Add 1 to 1½ gallons of Riverdale 2D + 2DP to 100 gallons of water using 200 to 600 gallons of spray mixture per acre, depending upon the height and thickness of the brush. Mix thoroughly before spraying.

BASAL BARK TREATMENT:

Thoroughly wet the base and root collar of all stems until the spray accumulates around the root collar at the ground line. This spray may be applied during any season. Use Riverdale 2D + 2DP for basal bark treatment on scattered brush or as a second spray application on species resistant to first foliage application. Mix 3 to 4 gallons of Riverdale 2D + 2DP in 100 gallons of oil. Apply with a low-volume sprayer or power equipment. Application rate will depend on species present, season applied and volume of spray used. Use a coarse spray to avoid drift.

MODIFIED BASAL TREATMENT

Drench the base of plants, then wet the lower 4/5 of remaining stems and leaves thoroughly to run-off. Apply treatment when brush is in full foliage. This method can be applied where susceptible species have been controlled by prior sprays and more resistant species, such as maple and oak, remain. Soaking the base of the plant and wetting all stems to run-off is absolutely necessary for complete control.

EARLY SEASON SPRAYING: Add 1 to 1½ gallons of Riverdale 2D + 2DP to 10 gallons of diesel oil and thoroughly mix. Add this mixture to 89 gallons of water.

DURING DRY WEATHER OR THE LATTER PART OF SPRAYING SEASON:: Add 1½ gallons of Riverdale 2D + 2DP to 15 gallons of diesel oil and mix thoroughly. Add mixture to 63-1/2 gallons of water and agitate thoroughly before use to insure uniform mixing. DO NOT allow mixture to stand more than 1 hour after mixing.

CUT SURFACE TREATMENT:

STUMPS: This treatment may be used anytime of the year; however, it is more effective when applied as quickly as possible after trees are cut. Spray the entire stump, especially exposed roots and bark. A complete soaking is essential for effective control. Use this procedure after original or capital removal. It is the first step towards a chemical brush control program on newly cleared highways

and rights-of-way. The spray is most effective and profitable on stumps 3 to 4 inches and larger. Mix 3 to 4 gallons of Riverdale 2D + 2DP in 100 gallons of oil. Applications should be made with a low-volume knapsack sprayer using a solid cone-shaped nozzle of medium orifice.

FRILL: Make a frill using an axe to cut overlapping V-shaped notches in a continuous ring, and cut around the trunk near its base. Cut through the bark, but do not remove the chips. This method is recommended for all trees 5 to 6 inches in diameter and larger. Freshly cut frills can be treated anytime of the year. Mix 3 to 4 gallons of Riverdale 2D + 2DP in 100 gallons of oil. Pour in as much of the mixture the frills will hold without wasting the chemical.

STORAGE AND DISPOSAL:

DO NOT contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

PESTICIDE DISPOSAL: Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to Federal, or approved state procedures under Subtitle C of the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL: DO NOT reuse empty container. Triple rinse (or equivalent) and offer for recycling or reconditioning or dispose of in a sanitary landfill, or by other approved State and local procedures.

WARRANTY AND LIMITATION OF DAMAGES:

Seller warrants that this material conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use and Buyer assumes the risk of any use contrary to such directions. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY, INCLUDING ANY OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR OF MERCHANTABILITY, AND NO AGENT OF SELLER IS AUTHORIZED TO DO SO EXCEPT IN WRITING WITH A SPECIFIC REFERENCE TO THIS WARRANTY. In no event shall Seller's liability for any breach of warranty exceed the purchase price of the material as to which a claim is made.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(a). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Richard F. Huntford *RHM*
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (TS-767)

Enclosure