

RIVERDALE 101
WEEDKILLER

DEC 14 1982
Under the Federal
Fungicide, and Herbicide Act
as amended, this product
registered under
EPA Reg. No. 228-1111

FOR PROFESSIONAL TURF MAINTENANCE
OPERATORS ONLY

(Contains 0.33 Pounds Dicamba and
3.17 Pounds 2,4-D Per Gallon)

ACTIVE INGREDIENTS:

Dimethylamine Salt of dicamba (3,6-dichloro-o-anisic acid)*.....	4.17%
Dimethylamine Salts of Related Acids.....	0.67%
Dimethylamine Salt of 2,4-dichlorophenoxy- acetic Acid**.....	40.15%
INERT INGREDIENTS;.....	<u>55.01%</u>
TOTAL.....	100.00%

*Equivalent to 3.46% 3,6-dichloro-o-anisic acid (dicamba)
or 0.33 pounds per gallon.

**Equivalent to 33.3% 2,4-dichlorophenoxyacetic acid or
3.17 pounds per gallon; Isomer Specific by AOAC Method
6 D01-5 (12th Ed.)

KEEP OUT OF REACH OF CHILDREN

DANGER

SEE LEFT PANEL FOR ADDITIONAL CAUTIONS

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

CHICAGO HEIGHTS, ILLINOIS

E.P.A. Reg. No. 228- E.P.A. Est. No. 228-IL-1

Net Contents _____ Gallons

DANGER

Hazard to Humans and Domestic Animals.

STATEMENT OF PRACTICAL TREATMENT

Danger. Corrosive, causes eye damage and skin irritation. Do not get in eyes, on skin or clothing. Wear goggles or face shield and rubber gloves when handling. May be fatal if swallowed. Avoid breathing vapors. If in eyes flush with plenty of water and get medical attention. If on skin wash with plenty of soap and water. Get medical attention if irritation persists. If swallowed, drink one or two glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

Environmental Hazard.

This product is toxic to fish. Do not apply directly to water. Do not contaminate water by the cleaning of equipment or disposal of wastes. Apply this product only as directed on label.

IMPORTANT

Do not apply on or near desirable trees or plants or in locations where chemicals may be washed or moved in contact with their roots.

Avoid drift of spray (coarse sprays are less likely to drift) onto vegetables, flowers, ornamental shrubs, trees, and other desirable crop plants.

Do not contaminate domestic or irrigation waters.

Do not pour solution used to clean sprayer near susceptible plants or where it may contaminate water.

Do not store in direct contact with fertilizer, seeds, insecticides, or fungicides.

Do not use on centipede grass, St. Augustine grass or dichondra nor on lawns or turf where desirable bent-grass or clovers are present. Newly seeded lawns may be treated after they have been mowed the second time. To re-seed small areas in treated lawns wait at least 3 weeks after spraying. A good rain or deep watering should occur during this period.

DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

PESTICIDE DISPOSAL

Pesticide, spray mixture, or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures under the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL

Metal — Triple rinse (or equivalent) and offer for recycling or reconditioning or dispose of in a sanitary land fill, or by other approved State and local procedures

Plastic — Triple rinse (or equivalent) and offer for recycling or reconditioning or dispose of in a sanitary land fill or by incineration if permitted by State and local authorities.

For Use on Lawns, Turf (not pastures) and Golf Fairways, Aprons, Tees and Rough For the control of burdock, chickweed (mouse-ear/common stitchwort), chicory, clovers, dandelions, docks, dog fennel, English daisy, Frenchweed, knotweed (erect/prostrate), ground ivy, hawkweed, healall, henbit, jimsonweed, lambsquarters, mallow, morning glory, mustard, oxalis, pennywort, peppergrass, pigweed, plantain (buckhorn, broadleaf, Rugel's) poison ivy, poison oak, purslane, ragweed, smartweed, sheep sorrel, spotted spurge, thistle (Canada), wild aster, wild carrot, wild garlic, wild onion, wild radish, wood sorrel, yarrow and many other broadleaf weeds, apply 1 to 1.3 fluid ounces of ~~101 Weed Killer~~ **Rivendale**

101 Weed Killer

to 1 to 2 gallons of water per 1,000 sq ft. (equivalent to 11 lbs. dicamba + 108 lbs. 2,4 D per acre to 0.14 lbs. dicamba + 1.4 lbs. 2,4 D per acre). Apply as a foliar spray when weeds are young and actively growing. It may be necessary to make a second application (spring and fall) to kill certain hard-to-kill weeds such as poison ivy, poison oak and English daisy. Weeds that start growing in the fall, such as henbit, may be controlled more effectively with a fall application.

WARRANTY

Rivendale

~~101 Weed Killer~~ warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant, and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.