PIVERDALE



81 SELECTIVE WEEDKILLER

FOR PROFESSIONAL TURF MAINTENANCE

Contains 3.06 pounds 2,4-D and 0.4 pounds Dicamba per gallon

ACTIVE INGREDIENTS:	
Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	
Dimethylamine Salt of Dicamba (3,6-dichloro-o-anisic Acid)**	5.12%
Dimethylamine Salts of related Acids	1.28%
INERT INGREDIENTS:	54.39%
TOTAL I	00.00%
Isomer Specific AOAC Method, Equivalent to: *2,4-Dichlorophenoxyacetic Acid	

KEEP OUT OR REACH OF CHILDREN

DANGER

See Side Panel For Additional Precautionary Statements
And Statement Of Practical Treatment

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near seeds, fertilizers, insecticides or fungicides. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well-ventilated area. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not contaminate water, food or feed by storage or disposal. PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate ground water. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional. Office for Guidance. CONTAINER DISPOSAL: Triple rinse (or equivilent). Then offer for recycling or . reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic containers are also disposable by incineration, of if allowed by state and local authorities.by.. burning. If burned, stay out of smoke.

NE" CONTENTS

EPA REG. NO. 228-169

EPA EST. NO. 228-IL-

1. Fam. 1 6/18/9.

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLIMOIS 60425-1584

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Corrosive, causes eye damage. Do not get in eyes, or on shin or clothing. Wear face shield or goggles and long pants, long-sleeved shirt, shoes and chemical resistant gloves when mixing, loading or applying this product. Wash nondisposable gloves thoroughly with soap and water before reusing. Harmful if swallowed or inhaled. Avoid inhalation of spray mists. Remove contaminated clothing and wash before reuse.

STATEMENT OF PRACTICAL TREATMENT

IF IN F.ES: Immediately flush eyes with plenty of water for 15 minutes and get

medical attention at once.

TF SWALLOWED: Drink one or two glasses of water. Induce vomiting by touching

back of throat with finger. Do not induce vomiting or give

anything by mouth to an unconscious person. Get medical attention

immediately.

IF ON SKIN: Wash promptly with soap and water and rinse thoroughly.

ENVIRONMENTAL HAZARDS.

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). When cleaning equipment, do not pour washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not containate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D, have been associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

USE FRECAUTIONS

CARE MUST BE TAKEN TO AVOID DRIFT OF SPRAY MISTS TO NON-TARGET susceptible plants such as cotton, vegetables, flowers, ornamentals, shrubs, trees, and other desirable plants. Do not spray near these plants. Do not apply on or near desirable trees or plants or in locations where chemicals may be washed or moved in contact with their roots. Do not spray roots of desirable trees and plants. Do not spray on carpetgrass, dichondra, nor on grasses where desirable clovers are present. Except as noted, use a lawn type sprayer with coarse spray as wind drift is less likely. Use the higher rate for hard-to-kill weeds. Do not exceed specified application dosages for any area. Do not apply to newly seeded grasses until well established. Do not spray when air temperature exceeds 90°F. Avoid applying during excessively dry or hot periods unless irrigation is used. For optimum results, (1) turf should not be moved 1 to 2 days before and following application; (2) Do not apply if rainfall is expected within 48 hours; nor should lawns be irrigated (watered) for 48 hours following application. Reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water and rinse thoroughly before reuse for other sprays. Failure to observe the above precautions may result in turf injury.

ORNAMENTAL LAWNS AND TURF

Riverdale 81 Selective Weed Killer is for use on Lawns, Turf (not pastures) and Golf Fairways, Aprons, Tees and Rough. The following weeds will be controlled by Riverdale 81 Selective Weed Killer: Burdock, Canada thistle, Chickweed (mouseear: common and stitchwort), Chicory, Clovers, Dandelions, Docks, Dog fennel, English daisy, Frenchweed, Ground ivy, Hawkweed, Healall, Henbit, Jimsonweed, Knotweed (erect and prostrate), Lambsquarters, Mallow, Morningglory, Mustard, Oxalis, Pennywort, Peppergrass, Pigweed, Plantain (buckhorn, broadleaf, Rugel's), Poison ivy, Poison oak, Purslane, Ragweed, Sheep sorrel, Smartweed, Spotted spurge, Wild aster, Wild carrot, Wild garlic, Wild onion, Wild radish, Wood sorrel, Yarrow, and many other broadleaf weeds.

Apply as a foliar spray when weeds are young and actively growing. It may be necessary to make a second application (Spring and Fall) to kill certain hard-to-kill weeds such as Poison ivy, Poison oak, and English daisy. Weeds that start growing in the Fall, such as Henbit, may be controlled more effectively with a Fall application.

Apply Riverdalc 81 Selective Weed Killer at a rate of 1-1/2 to 2 quarts per acre in 40 to 100 gallons of water; for smaller areas us 1 to 1.4 fluid ounces in 1 to 2 gallons of water per 1,000 square feet.

WARRANTY

Riverdale warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant, and soil conditions, mixtures with other chemicals, and factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.