

Riverdale

ACCEPTED
100
2/11

34-0-10 + Benefin Crabgrass Preventer

Minimum Guaranteed Analysis

Total Nitrogen (N)	34%
Controlled release nitrogen derived from sulfur coated urea	6%
Available Phosphoric Acid (P_2O_5)	0%
Soluble Potash (K_2O)	10%
Sulfur (S)	4%

Herbicide Active Ingredients

N-butyl-N-ethyl-a, a, a-trifluoro-2, 6-dinitro-p- toluidine *	1.175%
Inert ingredients	<u>98.825%</u>
	100.000%

* Balfin™ the trademark for ELANCO Products Company's benefin

KEEP OUT OF REACH OF CHILDREN

CAUTION:

See back of bag for additional precautionary statements.

E.P.A. Reg. No. 228-

E.P.A. Est. No. 228-11-1

50 Lbs. Net Weight Coverage 17,000 sq. ft.

Manufactured by
Riverdale Chemical Company
Chicago Heights, Illinois

34-0-10 + BENEFIN CRABGRASS PREVENTER

Riverdale
34-0-10 + Benefin Crabgrass Preventer is a high quality fertilizer product that combines turf food to feed the lawn and a selective pre-emergence herbicide which controls smooth and hairy crabgrass, annual bluegrass (Poa Annua), green and yellow foxtail, barnyard grass, and goosegrass (silver crabgrass or crow's foot)

Riverdale
34-0-10 + Benefin Crabgrass Preventer can be applied to established turf of bluegrass, ryegrass, fescue, zoysis, and Bermuda and will not result in injury to the turf when applied as directed

DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling

For Crabgrass:

Apply 130# - 175# 34-0-10 + Benefin per acre or 3# - 4# per 1,000 sq ft. At these rates you will be supplying 1# - 1.3# of nitrogen per 1,000 sq ft. and 1.5# - 2# of actual Benefin per acre. Coverage per 50# bag will be 17,000 - 12,750 sq. ft. Do not reseed 12 to 16 weeks after application

For Poa Annua control:

Apply 175#/acre or 4#/1,000 sq. ft. One ton covers 11.7 acres, or one 50# bag covers 12,750 sq ft. This rate supplies 1.3# nitrogen and 2# technical Benefin per 1,000 sq ft. Do not reseed the treated area for 12 to 16 weeks after application

PRECAUTIONS:

Apply 34-0-10 + Benefin Crabgrass Preventer to turf prior to germination of weed grasses since 34-0-10 + Benefin will not control established weeds. 34-0-10 + Benefin should not be applied in the spring to turfgrass planted the previous fall, in Bermudagrass, and areas that have been overseeded with winter grass. A spring application of 34-0-10 + Benefin may thin the overseeded grasses. Because of varying cultural practices 34-0-10 + Benefin should not be used on putting greens. 34-0-10 + Benefin should not be used on Dichondra or Bentgrass. 34-0-10 + Benefin should not be applied to newly sprigged areas of Bermudagrass, St. Augustine or Centipedegrass until these turfs are well established

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing

IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists

ENVIRONMENTAL HAZARDS

Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes

Storage and Disposal:

Do not contaminate water, food or feed by storage or disposal. Do not use empty bag. Place empty bag in trash collection

Notice: The manufacturer makes no warranties, express or implied, concerning this product or its use which extend beyond the description on the label. All statements made concerning this product apply only when used as directed