

OCT 9 1986

Mr. Russell F. Sawyer
Riverdale Chemical Company
220 East 17th Street
Chicago Heights, IL 60411

Dear Mr. Sawyer:

Subject: Label Amendments
Riverdale Weedstroy Triamine
EPA Registration No. 228-178
Your Submission Dated October 2, 1986

The amendment referred to above, submitted in connection with registration under section 3(c)(7)(A) of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:
 - a. In the ingredient substatement specify that the isomer specific method only applies to 2,4-D. Refer to the last accepted label for this product.
 - b. Refer to the child-resistant packaging regulations published in the Federal Register on June 11, 1986.
3. Submit five (5) copies of your final printed labeling before you release the product for shipment.

90561:Mountfort:MF-2:KENCO:10/7/86:10/17/86:bjp:lf

CONCURRENCES

| | | CONCURRENCES | | | | | | |
|-----------|--|--------------|--|--|--|--|--|--|
| SYMBOL ▶ | | | | | | | | |
| SURNAME ▶ | | | | | | | | |
| DATE ▶ | | | | | | | | |

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

Richard F. Mountfort *RFM*
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (TS-767C)

Enclosure

(FRONT PANEL)
RIVERDALE
WEEDESTROY^(R)
TRIAMINETM

ACCEPTED
with comments
OCT 9 1986
U.S. Federal Insecticide, Fungicide, and Rodenticide Act
Registered under Reg. No. 228-178

FOR SELECTIVE BROADLEAF WEED CONTROL
IN ORNAMENTAL LAWNS AND TURF GRASSES

CONTAINS 2,4-D, (MECOPROP AND DICHLORPROP)

only change

ACTIVE INGREDIENTS:

| | |
|--|--------|
| Dimethylamine Salt of 2,4-Dichlorophenoxy Acid* | 16.3% |
| Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid** | 16.4% |
| Dimethylamine Salt of 2-(2,4-Dichlorophenoxy)propionic Acid*** | 16.2% |
| INERT INGREDIENTS | 51.1% |
| TOTAL | 100.0% |

By Isomer Specific A.O.A.C. Method, Equivalent to:

| | | |
|--|-------|--------------|
| *2,4-dichlorophenoxyacetic acid | 13.6% | 1.3 lbs/gal. |
| **2-(2-methyl-4-chlorophenoxy)propionic acid | 13.6% | 1.3 lbs/gal. |
| ***2-(2,4-dichlorophenoxy)propionic acid | 13.6% | 1.3 lbs/gal. |

KEEP OUT OF REACH OF CHILDREN
DANGER
See Side Panel For Additional Precautionary Statements

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Immediately flush eyes with plenty of water for 15 minutes and get medical attention at once.

IF SWALLOWED: Drink one or two glasses of water. Induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention immediately.

IF ON SKIN: Wash promptly with soap and water and rinse thoroughly.

Weedestroy and Triamine are Trademarks of Riverdale Chemical Co. Patent Applied For

40 CFR Part 157 ONLY FOR SALE TO, USE AND STORAGE BY
LAWNCARE AND LANDSCAPING PERSONNEL OR COMMERCIAL APPLICATORS

E.P.A. REG. NO. 228-178 NET CONTENTS GAL. E.P.A. EST. NO. 228-IL-1

MANUFACTURED BY
RIVERDALE CHEMICAL COMPANY
CHICAGO HEIGHTS, ILLINOIS 60411-3699

(LEFT SIDE PANEL)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Corrosive, causes eye damage. Do not get in eyes, or skin, or clothing. Wear goggles or face shield and rubber gloves when handling. If eye exposure occurs, get immediate medical attention. Avoid inhalation or spray mists. Harmful or fatal if swallowed. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Weedestroy Triamine may cause dermal sensitization reactions.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not apply when weather conditions favor drifts from target area. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate domestic or irrigation waters.

DIRECTIONS FOR USE


It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

GENERAL INFORMATION

WEDESTROY TRIAMINE is a stable solution containing a mixture of three phenoxy herbicides. It is non-flammable. It is for use on Lawns, Turf(not pastures) and Golf Fairways, Aprons, Tees and Roughs.

USE PRECAUTIONS

CARE MUST BE TAKEN TO AVOID DRIFT OF SPRAY MISTS TO NON-TARGET susceptible plants such as cotton, vegetables, flowers, ornamentals, shrubs, trees, and other desirable plants. Do not spray near these plants. Do not spray on carpet grass, dichondra, nor on grasses where desirable clovers are present. Use a lawn type sprayer with coarse spray as wind drift is less likely. Avoid fine mists. Do not spray roots of desirable plants. Do not exceed specified application dosages for any area. Do not apply to newly seeded grasses until well established. Do not spray when air temperature is greater than 85°F. Reseed no sooner than three to four weeks after application of this product. Clean and rinse spray equipment using soap or detergent and water and rinse thoroughly before reuse for other sprays. Failure to observe the above precautions may result in injury.



(LEFT SIDE PANEL - CONT')

WEEDS CONTROLLED BY WEEDESTROY TRIAMINE

Black Medic, Buckhorn, Burdock, Buttercup, Canada Thistle, Carpet Weeds, Chickweed, Chicory, Clover, Cocklebur, Dandelion, Dock, Dog Fennel, English Daisy, Florida Parsley, Frenchweed, Ground Ivy, Hawkweed, Heal-All, Heartloaf Drymary, Henbit, Jimson Weed, Knotweed, Lambs-quarter, Mallow, Morning Glory, Mustard, Oxalis, Pennywort, Pepperweed, Pigweed, Plantain, Poison Ivy, Poison Oak, Purslane, Rag Weed, Red Clover, Red Sorrel, Sheep Sorrel, Smartweed, Speedwell, Spurge, Spur Weed, Stitchwort, Toadflax, Veronica, Wild Aster, Wild Carrot, Wild Garlic, Wild Geranium, Wild Lettuce, Wild Onion, Wild Radish, Wood Sorrel, Yarrow and many other broadleaf weeds.

ORNAMENTAL LAWNS AND TURF (Cool Season Grasses)

For normal application, use Weedestroy Triamine at a rate of 3.4 pints per acre in sufficient water for good coverage (40-80 gallons of water per acre or 1-2 gallons of water per 1000 square feet is normal). Use high rate for hard to kill weeds. Spray when air temperature is between 50-85°F. Avoid applying during excessively dry or hot periods unless irrigation is used. Do not irrigate within 24 hours of application. Do not mow one or two days before or after application.

(RIGHT SIDE PANEL)


Controlled Droplet Applicator: Reduced rates of Weedestroy Triamine must be used when grass is stressed from heat, drought, etc. Do not use this method on warm season grasses. Add 2 pints Weedestroy Triamine to the herbicide container then fill with 3 pints of water, keeping container agitated. Spray entire contents over 33,000 square feet (approximately 3/4 acre). Do not overlap between spray patterns.

BENT GRASS TURF

On closely mowed bentgrass, apply Weedestroy Triamine at a maximum rate of 2/3 ounce in 3 gallons water per 1000 square feet (2-3 pints per acre) in May or Mid-August through September when weeds are actively growing. Exercise care to avoid overdosing Bentgrass or injury may result. Do not mow two days before or after application. Do not apply during hot or dry periods. Slight turf yellowing should disappear after about 1 week.

ST. AUGUSTINE AND CENTIPEDE GRASSES

Apply Weedestroy Triamine at a maximum rate of 2-3 pints in up to 150 gallons of water per acre (2/3 - 1 1/3 oz. in 3 gallons of water per 1000 square ft.). Do not spray St. Augustine or Centipede grasses if stressed from heat, drought, etc. Slight turf yellowing should disappear after about 1 week.



STORAGE AND DISPOSAL

STORAGE: Always store pesticides in a secured warehouse or storage building. Do not store near seeds, fertilizers, insecticides or fungicides. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed, labeled container for proper disposal. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and Local authorities. Plastic containers disposed also by incineration, or if allowed by State and Local authorities, by burning. If burned, stay out of smoke.

WARRANTY

Riverdale warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant, and soil conditions, mixtures with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.