

228-178

03/08/2012

1/11



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF  
CHEMICAL SAFETY AND  
POLLUTION PREVENTION

Mr. George Meindl  
Nufarm Americas, Inc.  
150 Harvester Drive, Suite 200  
Burr Ridge, IL 60527

MAR - 8 2012

Subject: Label Amendment to comply with MCPP-p and 2,4-Dp REDs  
Product Name: TRIAMINE  
EPA Reg. No. 228-178  
Revised label submitted: March 8, 2012

Dear Mr. Meindl:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is acceptable.

One copy of the label stamped "Accepted" is enclosed for your records. Products shipped after 18 months from the date on this notice or the next printing of the label, whichever occurs first, must bear the new revised label. Amended labeling will supersede all previously accepted ones. Submit one (1) copy of final printed labeling before you release the product for shipment.

If you have questions or concerns regarding this letter, please contact Beth Benbow at (703) 347-8072 or email at [benbow.bethany@epa.gov](mailto:benbow.bethany@epa.gov).

Sincerely,

A handwritten signature in black ink, appearing to read "Kathryn V. Montague".

Kathryn V. Montague  
Product Manager 23  
Herbicide Branch  
Registration Division (7505P)

2/11

# TRIAMINE®

**A THREE-WAY POST-EMERGENT SELECTIVE BROADLEAF HERBICIDE CONTAINING THE AMINES OF 2,4-D, MECOPROP-p AND DICHLORPROP-p.  
FOR USE ON GOLF COURSES, PARKS, HIGHWAYS, ORNAMENTAL TURF LAWNS AND SIMILAR NON-CROP AREAS. KILLS DANDELIONS, CHICKWEEDS, PLANTAINS, OXALIS, SPURGE AND MANY OTHER BROADLEAF WEEDS, SOME OF WHICH ARE LISTED ON THIS LABEL.**

**ACTIVE INGREDIENTS:**

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	16.3%
Dimethylamine Salt of (+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**‡	8.2%
Dimethylamine Salt of (+)-R-2-(2,4-Dichlorophenoxy)propionic Acid***	8.1%

<b>OTHER INGREDIENTS:</b>	<b>67.4%</b>
<b>TOTAL:</b>	<b>100.0%</b>

Isomer Specific Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid	13.6%, 1.24 lbs./gal.
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid	6.8%, 0.62 lbs./gal.
***(+)-R-2-(2,4-Dichlorophenoxy)propionic Acid	6.8%, 0.62 lbs./gal.

‡CONTAINS THE SINGLE ISOMER FORMS OF MECOPROP-p AND DICHLORPROP-p.


**KEEP OUT OF REACH OF CHILDREN**

**DANGER - PELIGRO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.  
(if you do not understand the label, find someone to explain it to you in detail.)

**SEE INSIDE BOOKLET FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS**

For Chemical Spill, Leak, Fire, or Exposure, Call CHEMTREC (800) 424-9300  
For Medical Emergencies Only, Call (877) 325-1840



EPA REG. NO. 228-178  
EPA EST. NO. 228-IL-1

NET CONTENTS \_\_\_\_\_ Gallons ( \_\_\_\_\_ ) Liters  
[Designation as "NONREFILLABLE" or "REFILLABLE" for containers >5 gals

000228-00178.20120308.revised due to MCPPP-p and 2,4-DP-p RED

Manufactured For  
Nufarm Americas Inc.  
150 Harvester Drive  
Burr Ridge, IL 60527



**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
DANGER**

Corrosive, causes irreversible eye damage. Harmful if swallowed. Harmful if absorbed through skin. Harmful if inhaled. Do not get in eyes, on skin or on clothing. Avoid breathing spray mist or vapor.

**PERSONAL PROTECTIVE EQUIPMENT (PPE):**

Some materials that are chemical-resistant to this product are barrier laminate, butyl rubber, nitrile rubber or viton. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

**Mixers, loaders and all other handlers must wear:**

- Long-sleeved shirt and long pants,
- Shoes and socks,
- Chemical resistant gloves made of waterproof material,
- Goggles, face shield or shielded safety glasses, and
- Chemical-resistant apron when mixing, loading, cleaning up spills or equipment, or otherwise exposed to the concentrate.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry. Discard clothing and other absorbent material that have been drenched or heavily contaminated with the product's concentrate. Do not reuse them.

<b>USER SAFETY RECOMMENDATIONS</b>	
<b>Users Should:</b>	<ul style="list-style-type: none"> <li>• Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.</li> <li>• Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. If pesticide gets on skin, wash immediately with soap and water.</li> <li>• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.</li> </ul>

<b>FIRST AID</b>	
<b>IF IN EYES</b>	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15 to 20 minutes.</li> <li>• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> </ul>
<b>IF SWALLOWED</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> </ul>
<b>IF ON SKIN OR CLOTHING</b>	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<b>IF INHALED</b>	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth- to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
<b>HOT LINE NUMBER</b>	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-877-325-1840 for emergency medical	
<b>NOTE TO PHYSICIAN</b>	
Probable mucosal damage may contraindicate the use of gastric lavage.	

### ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in water adjacent to treated areas. Do not contaminate water when disposing of equipment wash waters or rinsate.

This chemical has properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination. Application around a cistern or well may result in contamination of drinking water or groundwater.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. Do not enter or allow people (or pets) to enter the treated area until sprays have dried.

Aerial application of this product is prohibited. Do not use this product on or near desirable plants, including within the dripline of the roots of desirable trees and shrubs, since injury may result.

### SPRAY DRIFT MANAGEMENT

A variety of factors including weather conditions (e.g., wind direction, wind speed, temperature, relative humidity) and method of application can influence pesticide drift. The applicator must evaluate all factors and make appropriate adjustments when applying this product.

#### Droplet Size

Use only medium or coarser spray nozzles according to ASAE (S572) definition for standard nozzles.

#### Wind Speed

Do not apply at wind speeds greater than 10 mph. Only apply this product if the wind direction favors an on target deposition and there are not sensitive areas (including, but not limited to residential areas, bodies of water, known habitat for nontarget species, nontarget crops) within 250 feet downwind.

#### Temperature Inversions

If applying at wind speeds less than 3 mph, the applicator must determine if: 1) conditions of temperature inversion exist, or 2) stable atmospheric conditions exist at or below nozzle height. Do not make applications into areas of temperature inversions or stable atmospheric conditions.

#### Susceptible Plants

Do not apply under circumstances where spray drift may occur to food, forage, or other plantings that might be damaged or crops thereof rendered unfit for sale, use or consumption. Susceptible crops include, but are not limited to, cotton, okra, flowers, grapes (in growing stage), fruit trees (foliage), soybeans (vegetative stage), ornamentals, sunflowers, tomatoes, beans, and other vegetables, or tobacco. Small amounts of spray drift that might not be visible may injure susceptible broadleaf plants.

#### Other State and Local Requirements

Applicators must follow all state and local pesticide drift requirements regarding application of 2,4-D herbicides. Where states have more stringent regulations, they must be observed.

#### Equipment

All ground application equipment must be properly maintained and calibrated using appropriate carriers or surrogates. Do not apply with a nozzle height greater than 4 feet above the crop canopy.

### PRODUCT INFORMATION

This product is a stable solution containing a mixture of three herbicides. It is nonflammable. It is for use on Ornamental Turf and Non-Cropland areas. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes. Not for use on sod farms.

CARE MUST BE TAKEN TO AVOID DRIFT OF SPRAY MISTS TO NON-TARGET susceptible plants such as cotton, vegetables, flowers, ornamentals, shrubs, trees, and other desirable plants. Do not spray near these plants. Do not spray on carpetgrass, dichondra, nor on grasses where desirable clovers are present. Except as noted, use a lawn type sprayer with coarse spray as wind drift is less likely. Avoid fine mists. Do not spray exposed roots of desirable plants. Use the higher rate for hard-to-kill weeds. Do not exceed specified application dosages for any area. Do not apply to newly seeded grasses until well established. Avoid applying during excessively dry or hot periods unless irrigation is used. Avoid broadcast applications when air temperature exceeds 90 degrees. When using small, spot treatment applications in temperature over 90 degrees, turf injury may occur.

When treating Carpetgrass and St. Augustine grass, avoid broadcast applications when air temperature exceeds 80°F. When air temperatures exceed 80°F, limit application to spot treatment only.

For optimum results, (1) turf should not be mowed 1 to 2 days before and following application; (2) do not apply if rainfall is expected within 48 hours; nor should lawns be irrigated (watered) for 48 hours following application. Reseed no sooner than 3 to 4 weeks after application of this product. Adding oil, wetting agent, or other surfactant to the spray may be used to increase effectiveness on weeds but doing so may reduce selectivity to turf resulting in turf damage. Clean and rinse spray equipment using soap or detergent and water and rinse thoroughly before reuse for other sprays. Failure to observe the above precautions may result in injury.

This product can be mixed with some liquid fertilizers or liquid iron materials. Because liquid fertilizers and liquid iron differ in pH, free ammonia content, density, salt concentration, and percentage of water, a compatibility test is recommended prior to mixing in the application equipment. All regulations, either State or Federal, relating to the application of liquid fertilizers or liquid iron and this product must be strictly followed.

### WEEDS CONTROLLED

Maximum kill of weeds will be obtained by applying in spring and early fall when weeds are actively growing.

Alder	Black medic	Carpetweed	Curly indigo
Alligatorweed	Blessed thistle	Catnip	Dandelion
American lotus	Blue lettuce	Chickweed	Dock
Arrowhead	Box elder	Chicory	Dogbane
Artichoke	Broomweed	Clover	Doofennel
Aster	Buckhorn	Cockle	Duckweed
Austrian fieldcress	Bull thistle	Cocklebur	Elderberry
Beggartick	Bulrush	Coffeebean	English daisy
Biden	Burdock	Coffeeweed	Flea bane (daisy)
Bindweed	Bur raaweed	Common sowthistle	Flixweed
Bitterweed	Buttercup	Creeping jenny	Florida pusley
Bitter winter cress	Canada thistle	Croton	Frenchweed
Galinsoga	Musk thistle	Rush	Vervain
Goatsbeard	Mustard	Russian thistle	Vetch
Goldenrod	Nettle	Sagebrush	Virginia creeper
Ground ivy	Nutarass	St. Johnswort	Water hvacinth
Gumweed	Orange hawkweed	Sheep sorrel	Water lily
Hawkweed	Oxalis	Shepherdspurse	Water plantain
Healall	Parrot feather	Sicklepod	Water primrose
Heartleaf drymary	Parsnip	Smartweed	Water shield
Hemp	Pennycress	Sneezeweed	Wild aster
Henbit	Pennywort	Southern wild rose	Wild carrot
Hoary cress	Pepperarass	Sowthistle	Wild garlic
Honeysuckle	Pepperweed	Spanishneedle	Wild geranium

Horsetail  
 Indigo  
 Indiana mallow  
 Ironweed  
 Jewelweed  
 Jimsonweed  
 Kochia  
 Knotweed  
 Lambsquarter  
 Locoweed  
 Lupine  
 Mallow  
 Marshelder  
 Mexicanweed  
 Morningglory

Pigweed  
 Plantain  
 Poison hemlock  
 Poison ivy  
 Poison oak  
 Pokeweed  
 Poorioe  
 Povertyweed  
 Prickly lettuce  
 Primrose  
 Puncture vine  
 Purslane  
 Ragweed  
 Red clover  
 Red sorrel

Spatterdock  
 Speedwell  
 Spurge  
 Spurweed  
 Stinkweed  
 Stitchwort  
 Sumac  
 Sunflower  
 Sweet clover  
 Tarweed  
 Thistle  
 Toadflax  
 Tumbleweed  
 Velvet leaf  
 Veronica

Wild lettuce  
 Wild onion  
 Wild radish  
 Wild rape  
 Wild strawberry  
 Wild sweet potato  
 Willow  
 Witchweed  
 Woodsorrel  
 Wormseed  
 Yarrow  
 Yellow rocket  
 Virginia buttonweed and  
 other broadleaf weeds.

**ORNAMENTAL TURF**

**(Golf Courses, Cemeteries, Parks, Sports Fields, Turfgrass, Lawns and Other Grass Areas)**

**Cool Season Grasses - such as bluegrass, fescue, and rye:** Apply this product at a rate of 3 to 4 pints per acre in sufficient water for good coverage, 40 to 80 gallons of water per acre (1 to 2 gallons of water per 1,000 square feet is normal). For small lawns, apply 5 to 7 teaspoons in 1 gallon of water per 1,000 square feet.

**Warm Season Grasses - such as St. Augustine, bahia, common Bermuda, centipede and zoysia grasses:** Apply this product at a maximum rate of 2 to 3 pints in up to 150 gallons of water per acre (2/3 to 1-1/3 fluid ounce. in 3 gallons of water per 1,000 square feet). Do not spray St. Augustine, common Bermuda or centipede grasses if stressed from heat, drought, etc. Slight turf yellowing should disappear after about 1 week. Exercise care when applying during growth stages from dormancy to green-up and from green-up to dormancy. Note: Some hybrid Bermuda grasses may be sensitive to this product. Contact your local extension service weed control specialist.

**Bentgrass Turf:** On closely mowed bentgrass, other than golf course greens and tees, apply this product at a maximum rate of 2/3 ounce in 3 gallons water per 1,000 square feet (1-3/4 pints per acre) in May or mid-August through September when weeds are actively growing. Exercise care when applying to avoid overdosing bentgrass or turf injury may result. Slight turf yellowing should disappear after about 1 week.

**Low-Volume Applicator:** When using a Low-Volume Applicator (such as Lawn Doctor's "Turf Tamer") apply this product at a rate of 3 to 4 pints per acre (1.1 fluid ounce to 1.5 fluid ounce. in 1/2 gallon of water per 1,000 square feet).

**Herbi™ Controlled Droplet Applicator:** Add 2 pints of this product to the 5 pint Herbi bottle, then fill with 3 pints of water, or 2-1/2 pints of water and 1/2 pint of a surfactant while agitating the solution. For warm season grasses, apply 1-1/2 pints of this product to 3-1/2 pints of water. While walking at approximately 1 pace (3 feet) per second; spray entire contents over 33,000 square feet (3/4 of an acre). Do not overlap (double coverage) at edge of spray patterns. Reduced rates (use 1/2 of rate shown above) of this product must be applied when grass is stressed from heat, drought, etc.

**Limitations and Restrictions for Ornamental Turf:**

For broadcast treatments, do not apply more than 9 pints of product per acre per application. Do not apply more than 18 pints of product per acre per season, excluding spot treatments. Do not make more than 2 applications per year with a minimum of 30 days between applications.

The maximum application rate for spot treatment is 1.2 lbs ae MCP-P per acre per application. For spot treatments do not apply more than 15 pints of product per acre per application. Do not make more than 2 applications per year with a minimum of 30 days between applications. Broadcast application is prohibited at this rate. A spot treatment is defined as a treatment area not greater than 1,000 square feet per acre.

**NON-CROPLAND  
(Fencerows, Hedgerows, Roadsides, Ditches, Rights-of-Way,  
Utility Power Lines, Railroads, Airports, and Industrial Sites)**

**For the control of broadleaf weeds:** Limited to 2 applications per year. Mix at a rate of 1/3 to 1 gallon of this product per 50 to 300 gallons of water. This mixture will cover 43,500 square feet. Thoroughly saturate all weeds with spray mixture. Apply anytime between the time when plants come into full leaf (spring) to when the plants begin to go dormant. Best results are obtained when weeds are young and actively growing. Do not cut weeds until herbicide has translocated throughout the plant causing root death. For small broadleaf weeds, use the lower rate. Heavy, dense stands require the higher rate with high water volume. For small (spot) applications with small tank sprayers, apply at the rate of 4 ounces of this product per gallon of water and spray to thoroughly wet all foliage. For spot treatments, do not exceed 15 pints of product per spot treatment. A spot treatment is defined as a treatment area not greater than 1,000 square feet per acre.

**For control of woody plants:** Apply to both stems and foliage anytime from the time foliage is completely matured until the time plants start to go dormant. All leaves, stems and suckers must be completely wet to the ground line for effective control. Regrowth may be anticipated on the more resistant species. Add 9 pints of this product to 100 gallons of water per 43,560 sq. ft. depending upon the height and thickness of the brush. Mix thoroughly before spraying. For spot treatment do not apply more than 15 pints of product per application. Do not make more than 2 spot treatment applications per year. Allow a minimum of 30 days between spot treatments. Spot treatment is defined as a treatment area no greater than 1,000 ft<sup>2</sup> per acre.

**Limitations and Restrictions for Non-Cropland:**

For postemergence non-cropland annual and perennial weeds, do not apply more than 9 pints product per acre per application. Do not make more than 2 broadcast applications per year with a minimum of 30 days between applications.

For postemergence non-cropland woody plants, do not apply more than 9 pints product per acre. Do not make more than 1 broadcast application per year. Spot treatments may be made if regrowth occurs. Do not apply more than 15 pints of product per spot treatment application. Do not make more than 2 spot treatments per year with a minimum of 30 days between.

Applications to non-cropland areas are not applicable to treatment of commercial timber or other plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

Use of this product in certain portions of California, Oregon and Washington is subject to the January 22, 2004 Order for injunctive relief in Washington Toxics Coalition, et al. v. EPA, C01-0132C, (W.D. WA). For further information, please refer to <http://www.epa.gov/espp>.

**STORAGE AND DISPOSAL**

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store this product in its original container in a cool, dry, locked place out of direct sunlight and out of the reach of children and pets. Do not use or store near heat or open flame.

**PESTICIDE DISPOSAL:** To avoid wastes, use all material in this container by application according to label directions. If waste cannot be avoided, offer remaining product to a waste disposal facility or pesticide disposal program (often such programs are run by state or local governments or by industry).

**CONTAINER DISPOSAL [HANDLING]:** [Note to Reviewer: The following statement will be included on all Final Printed Labels bearing multiple Container Disposal (Container Handling) statements] **NOTE:** This product is available in multiple containers. Refer to the Net Contents section of this products labeling for the applicable "Nonrefillable" or "Refillable" designation. Follow the container disposal [handling] instructions below that apply to your container type/size."

**Nonrefillable Containers 5 Gallons or Less:** Nonrefillable container. Do not reuse or refill this container.

Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

**Nonrefillable containers larger than 5 gallons:** Nonrefillable container. Do not reuse or refill this container. Offer for recycling if available. Triple rinse or pressure rinse container (or equivalent) promptly after emptying. **Triple rinse as follows:** Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water. Replace and tighten closures. Tip container on its side and roll it back and forth, ensuring at least one complete revolution, for 30 seconds. Stand the container on its end and tip it back and forth several times. Turn the container over onto its other end and tip it back and forth several times. Empty the rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Repeat this procedure two more times. **Pressure rinse as follows:** Empty the remaining contents into application equipment or a mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container, and rinse at about 40 psi for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

**Refillable containers larger than 5 gallons:** Refillable container. Refill this container with pesticide only. Do not reuse this container for any other purpose. Cleaning the container before final disposal is the responsibility of the person disposing of the container. Cleaning before refilling is the responsibility of the refiller. To clean the container before final disposal, empty the remaining contents from this container into application equipment or a mix tank. Fill the container about 10% full with water and, if possible, spray all sides while adding water. Agitate vigorously or recirculate water with the pump for two minutes. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this rinsing procedure two more times.

**Or**

Refillable Container: Refill this container with pesticide only. Do not reuse this container for any other purpose. Close all openings and replace all caps. Contact Nufarm's Customer Service Department at 1-800-345-3330 to arrange for return of the empty refillable container.

**WARRANTY DISCLAIMER**

The directions for use of this product must be followed carefully. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, (1) THE GOODS DELIVERED TO YOU ARE FURNISHED "AS IS" BY MANUFACTURER OR SELLER AND (2) MANUFACTURER AND SELLER MAKE NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND TO BUYER OR USER, EITHER EXPRESS OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE, OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE. UNINTENDED CONSEQUENCES, INCLUDING BUT NOT LIMITED TO INEFFECTIVENESS, MAY RESULT BECAUSE OF SUCH FACTORS AS THE PRESENCE OR ABSENCE OF OTHER MATERIALS USED IN COMBINATION WITH THE GOODS, OR THE MANNER OF USE OR APPLICATION, INCLUDING WEATHER, ALL OF WHICH ARE BEYOND THE CONTROL OF MANUFACTURER OR SELLER AND ASSUMED BY BUYER OR USER. THIS WRITING CONTAINS ALL OF THE REPRESENTATIONS AND AGREEMENTS BETWEEN BUYER, MANUFACTURER AND SELLER, AND NO PERSON OR AGENT OF MANUFACTURER OR SELLER HAS ANY AUTHORITY TO MAKE ANY REPRESENTATION OR WARRANTY OR AGREEMENT RELATING IN ANY WAY TO THESE GOODS.

**LIMITATION OF LIABILITY**

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL MANUFACTURER OR SELLER BE LIABLE FOR SPECIAL, INCIDENTAL, OR CONSEQUENTIAL DAMAGES, OR FOR DAMAGES IN THEIR NATURE OF PENALTIES RELATING TO THE GOODS SOLD, INCLUDING USE, APPLICATION, HANDLING, AND DISPOSAL TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, MANUFACTURER OR SELLER SHALL NOT BE LIABLE TO BUYER OR USER BYWAY OF INDEMNIFICATION TO BUYER OR TO CUSTOMERS OF BUYER, IF ANY, OR FOR ANY DAMAGES OR SUMS OF MONEY, CLAIMS OR DEMANDS WHATSOEVER, RESULTING FROM OR BY REASON OF, OR ARISING OUT OF THE MISUSE, OR FAILURE TO FOLLOW LABEL WARNINGS OR INSTRUCTIONS FOR USE, OF THE GOODS SOLD BY MANUFACTURER OR SELLER TO BUYER. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER, USER, OR ITS CUSTOMERS. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S OR USER'S EXCLUSIVE REMEDY, AND MANUFACTURER'S OR SELLER'S TOTAL LIABILITY SHALL BE FOR DAMAGES NOT EXCEEDING THE COST OF THE PRODUCT.



9/11

**If you do not agree with or do not accept any of directions for use, the warranty disclaimers, or limitations on liability, do not use the product, and return it unopened to the Seller, and the purchase price will be refunded.**

(RV050409RED)

Triamine and The Optical Advantage are Registered Trademarks of Nufarm Americas Inc.

**OPTIONAL MARKETING CLAIMS – FRONT PANEL**

**GET THE OPTICAL ADVANTAGE®**

This product contains a purified form of MCPP known as the optical isomer, which is the portion of the molecule active as an herbicide. The Optical Advantage® is not associated with user eye safety.

11/11

**LABEL HISTORY**

<b>File Name</b>	<b>Revision Mark</b>	<b>Comment</b>
00228.00178.20110929.revised due to 2,4-DP-p RED		
00228.00178.20120308.revised due to MCPP-p and 2,4-DP-p RED		Changes made per Beth Benbow's 20120221 email.