



- c. Under Storage and Disposal, revise the container disposal instructions and add pesticide disposal instructions as described in PR Notice 83-3 for products containing an active ingredient that appears on the RCRA F list (2,4-D).
- d. In the active ingredient statement for MCPP, change ". . .2(2-. . ." to ". . .2-(2 . . ."
- e. In the first line under GENERAL INFORMATION, change 5-3-2 to 12-1-1 consistent with the front page fertilizer analysis.

This registration is issued relying on reported N-nitroso contamination levels below 1 part per million (EPA Accession No. 260049). For future reference, the product specific chemistry data were assigned EPA Accession No. 260050.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 Enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA sec. 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Richard F. Mountfort *LTS*  
 Product Manager (23)  
 Fungicide-Herbicide Branch  
 Registration Division (TS-767C)

Enclosure

# Riverdale

## 3 WAY

### WEED AND FEED

**ACCEPTED**  
*with comments*  
NOV 21 1985 *LTS*  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act,  
as amended, for use as a herbicide,  
registered under:  
EPA Reg. No. 228-180

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION.

**ACTIVE INGREDIENTS:**

*Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid.....	0.156%
**Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid.....	0.157%
***Dimethylamine Salt of 2-(2,4-Dichlorophenoxy)propionic Acid.....	0.155%
Inert Ingredients.....	99.532%
Total	100.000%

**EQUIVALENT TO:**

- \*0.13% of 2,4-Dichlorophenoxyacetic Acid
- \*\*0.13% of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid
- \*\*\*0.13% of 2-(2,4-Dichlorophenoxy)propionic Acid

BY ISOMER SPECIFIC ASSAY

**MINIMUM GUARANTEED ANALYSIS**

Total Nitrogen (N).....	12.50%
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ).....	1.50%
Soluble Potash (K <sub>2</sub> O).....	1.50%

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

- IF SWALLOWED: Get Medical Attention.
- IF ON SKIN: Wash Skin with Soap and Water.
- IF IN EYES: Flush Eyes with Water for 15 Minutes and Get Medical Attention.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

MANUFACTURED BY

**RIVERDALE CHEMICAL COMPANY**  
CHICAGO HEIGHTS, ILLINOIS

EPA Reg. No. 220.

NET WEIGHT : 20 LBS/9.07 KG.  
Treats 2500 Sq. Ft. (25x100)

EPA Est. No. 228-IL-1

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly after handling.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not apply when conditions favor drift from target areas.

Do not contaminate water by cleaning of equipment or disposal of waste.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

STORAGE: Always store product in original container and in locked storage area.

CONTAINER DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash. Do not reuse bag.

5-11-83

200 2 av. F N.C. +

3.- WAY WEED AN FEED (5-3-2) is a stable mixture of three phenoxy herbicides. It is for use on Lawns, Turf (not pastures) and Golf Fairways, Aprons, Tees and Roughs.

### DIRECTIONS

Read the entire label before using this product.

This product may be used on Blue Grass, Fescue, Rye, Bermuda, Bahia, and Zoysia Lawns. DO NOT USE on Dichondra, Carpet Grass and St. Augustine grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" on flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

WHEN TO APPLY: Best results are obtained when weeds are growing rapidly in spring or early fall. IMPORTANT: Apply when grass is moist (after a rain, sprinkling or dew). Do not water lawn after material is applied. Watering will wash off weed killing material. For most effective broadleaf weed killing action, avoid unnecessary disturbance of treated areas for 48 hours after application. Do not re-seed for at least 3 to 4 weeks after application. Do not apply to newly seeded lawn until after 3 mowings.

HOW TO APPLY: This 20 pound bag should be used on 2500 square feet. Apply evenly with a lawn spreader either as one pass application or a two pass application, criss-crossing the area at half the rate each time. Lawns that are heavily overrun by weeds and that are in poor condition, may require more than one application of this product. A suggested second application should be made in approximately 30 days.

The following settings are approximate for applying contents of this package at a suggested rate of 8 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed.

SPREADER SETTINGS:	MAKE	SETTING
	Cyclone	6-7
	Scotts 35	5 1/2
	Scotts	C-2 J-2
	Central	7-8
	K-Mart	7-8
	Montgomery Wards	7-8
	Sears Broadcast	6-7

This product is designed for control of broadleaf weeds in lawns such as:

Black Medic, Buckhorn, Burdock, Chickory, Chickweed, Clover, Dock, Dandelion, English Daisy, Ground Ivy, Heal-All, Knotweed, Lambsquarter, Mallow, Morning Glory, Pigweed, Plantain, Poison Ivy, Poison Oak, Purslane, Ragweed, Red Clover, Red Sorrel, Stitchwort, Speedwell, Spurge, White Clover, Wild Carrot, Wild Garlic, Wild Onion, Wild Lettuc, Yarrow and other Broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing rapidly in spring or early fall. **IMPORTANT:** Apply when grass is moist (after a rain, sprinkling or dew). Do not water lawn after material is applied. Watering will wash off weed killing material. For most effective broadleaf weed killing action, avoid unnecessary disturbance of treated areas for 48 hours after application. Do not re-seed for at least 3 to 4 weeks after application. Do not apply to newly seeded lawn until after 3 mowings.

HOW TO APPLY: This 20 pound bag should be used on 2500 square feet. Apply evenly with a lawn spreader either as one pass application or a two pass application, criss-crossing the area at half the rate each time. Lawns that are heavily overrun by weeds and that are in poor condition, may require more than one application of this product. A suggested second application should be made in approximately 30 days.

The following settings are approximate for applying contents of this package at a suggested rate of 8 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed.

SPREADER SETTINGS:	<u>MAKE</u>	<u>SETTING</u>
	Cyclone	6-7
	Scotts 35	5 1/2
	Scotts	C-2 J-2
	Central	7-8
	K-Mart	7-8
	Montgomery Wards	7-8
	Sears Broadcast	6-7

This product is designed for control of broadleaf weeds in lawns such as:

Black Medic, Buckhorn, Burdock, Chickory, Chickweed, Clover, Dock, Dandelion, English Daisy, Ground Ivy, Heal-All, Knotweed, Lambsquarter, Mallow, Morning Glory, Pigweed, Plantain, Poison Ivy, Poison Oak, Purslane, Ragweed, Red Clover, Red Sorrel, Stitchwort, Speedwell, Spurge, White Clover, Wild Carrot, Wild Garlic, Wild Onion, Wild Lettuc, Yarrow and other Broadleaf weeds.

**WARRANTY**

Riverdale warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant, and soil conditions, mixture with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.