208-181. Please read instructions on reverse before completing form.	2003 Form Approved, OMB No. 2u
United States Environmental Protection A Washington, DC 20460	Registration OPP Identifier Number
Application fo	r Pesticide - Section I
1. Company/Product Number 228-181	2. EPA Product Manager J. I. Miller
4. Company/Product (Name) Riverdale Triamine Lawn Weed Killer	PM# None Re. tricted
5. Name and Address of Applicant (Include ZIP Code) Nufarm Americas, Inc. Riverdale - A Nufarm Company 1333 Burr Ridge Parkway, Suite 125A Burr Ridge, IL 60527-0866 Check if this is a new address	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA File Symbol Product Name
S	ection - II
Amendment - Explain below. Resubmission in response to Agency letter dated Notification - Explain below.	Final printed labels in response to Agency letter dated "Me Too" Application. Other - Explain below. DEC - 5 2003
action and penalties under sections 12 and 14 of FIFRA. Section 1. Material This Product Will Be Packaged In:	CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement
Yes ' No Yes No No If "Yes" No. per If "'	ter Soluble Packaging Yes No No Hetal Plastic Glass Yes" No. per ckage wgt container Other (Specify)
3. Location of Net Contents Information 4. Size(s) Retail Con Label Container	ntainer 5 Location of Label Directions On Label On Labeling accompanying product
6. Manner in Which Label is Affixed to Product Lithograph Paper glued Stenciled	Other
	ection - IV
1. Contact Point (Complete items directly below for identification of inc	
Name Russell F. Sawyer	Regulatory Affairs Manager 708/754-3330 ••
Certification I certify that the statements I have made on this form and all atta I acknowledge that any knowingly false or misleading statement both under applicable low.	
2. Signature 3. Title	Regulatory Affairs Manager
4. Typed Name 5. Date	November 10, 2003
Russell F. Sawyer	* • • • • • • • • • • • • • • • • • • •

RIVERDALE[®]

TRIAMINE LAWN WEED KILLER

FOR SELECTIVE BROADLEAF WEED CONTROL IN ORNAMENTAL LAWNS AND TURF GRASSES And Other Listed Sites SA CONTRACTOR

NOTIFICATION

DEC - 5 2003

Market Committee

'nga an a

THE THE THE THE

高額 一点洗

ar har sala a a a

ONE GALLON COVERS UP TO 80,000 SQUARE FEET (See directions for different grasses.)

GET THE OPTICAL ADVANTAGE

ACTIVE INGREDIENTS:

Dimethylamine Salt of	2,4-Dichlorophenoxyacetic Acid* 4.55% (+)-R-2-(2-Methyl-4-Chlorophenoxy) propionic Acid**‡ 2.30%
Dimethylamine Salt of	(+)-R-2-(2-Methyl-4-Chlorophenoxy) propionic Acid**. 2.30%
Dimethylamine Salt of	(+)-R-2-(2,4-Dichlorophenoxy) propionic Acid*** 4 2.26%
INERT INGREDIENTS:	_90.898
	TOTAL

Isomer Specific Method, Equivalent to:

3.8%, 0.32 lbs./gal. *2,4-Dichlorophenoxyacetic Acid 1.9%, 0.16 lbs./gal. 1.9%, 0.16 lbs./gal. **(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid ***(+)-R-2-(2,4-Dichlorophenoxy)propionic Acid

‡CONTAINS THE SINGLE ISOMER FORM OF MECOPROP-p AND DICHLORPROP-p.

Riverdale, Triamine and The Optical Advantage are Registered Trademarks of Riverdale - A Nufarm Company

KEEP OUT OF REACH OF CHILDREN

WARNING
SEE SIDE PANELS FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

NET CONTENTS

EPA REG. NO. 228-181

To Buston Louis Brook 21 or very EPA EST. NO. 228-IL-1

LANCE CONTRACTOR CONTRACTOR AS IN received the received

MANUFACTURED BY NUFARM AMERICAS INC., BURR RIDGE, ILLINOIS 60527-0866

a for yellor and the second and a second to the state of th to this of the market and the collection of

THE PROPERTY OF THE PARTY OF TH

SOUND THE CHESTON BUREAU WAY OF THE

Revised 2/14/03 Changed First Aid Statement per PR 2001-1 and deleted California use limitation via Amendment. to be the state of the state of

6/13/03 Corrected Ornamental Sites per EPA's Request 5/28/03 Meeting.

A/O 7/31/03 Corrected per EPA's letter dated 7/31/03.

11/10/03 Revised Residential label only.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

TOR ENGINEERS

Causes substantial but temporary eye injury. Harmful if swallowed, inhaled or absorbed through skin. Do not get in eyes or on clothing. Avoid breathing spray mist and contact with skin. Wear long pants, long-sleeved shirt, socks, shoes, chemical resistant gloves and eye protection when mixing, loading or applying this product or repairing or cleaning equipment used with this product. It is recommended that safety glasses include front, brow and temple protection. For aerial applicators in an enclosed cockpit and applicators applying this product from a tractor that has a completely enclosed cab, eye protection is not required. Wash hands, face and arms with soap and water as soon as possible after mixing, loading or applying this product. Wash hands, face and arms with soap and water before eating, smoking or drinking. Wash hands, face and arms before using toilet. After work, wash nondisposable gloves thoroughly with soap and water before removing, remove all clothing and shower using soap and water. Do not reuse clothing worn during the previous day's mixing and loading or application of this product without cleaning first. Clothing must be kept and washed separately from other household laundry. Remove saturated must be kept and washed separately from other household laundry. Remove saturated clothing as soon as possible and shower. When applying to turf the maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried.

If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron. If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable

transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

FIRST AID STATEMENT

EST. Have the product container or label with you when calling a poison control center or · Jana Control doctor, or going for treatment.

Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. IF IN EYES:

Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Call a poison control center or doctor immediately for treatment advice. IF SWALLOWED:

Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do

not give anything by mouth to an unconscious person.

IF ON SKIN Take off contaminated clothing. Rinse skin immediately with plenty of OR CLOTHING:

water for 15 to 20 minutes. Call a poison control center or doctor for

treatment advice.

Move person to fresh air. If person is not breathing, call 911 or an IF INHALED: ambulance, then give artificial respiration, preferably mouth-to-mouth if

possible. Call a poison control center or doctor for further treatment

· 克里克拉爾 一种护道强的 5 186

Lalthar Inco

advice.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters nor pour the washwater on the ground; spray or drain over a large area away from wells, other water sources and desirable plants. Do not apply this product through any type or irrigation system. not contaminate water used for irrigation or domestic purposes." If spills occur, collect the material and dispose of by following disposal instructions on this label. 医外侧侧 侧额的 员

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D, MCPP, and 2,4-DP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

GENERAL INFORMATION

TRIAMINE LAWN WEED KILLER is a stable solution containing a mixture of three herbicides. It is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs), also along Highways, Rights-of-Way, Drainage Ditchbanks and other similar turf areas.

USE PRECAUTIONS

IMPORTANT: Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray exposed roots of desirable plants. Do not spray on Dichondra, Lippia, nor on grasses where desirable Clovers are present. Do not apply to newly seeded grasses until they have been mowed 3 times. Do not reseed for 3 to 4 weeks after use. If lawn needs watering, water thoroughly before application. Do not water for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours. Avoid fine mists. lawn type sprayer with a coarse spray as wind drift is less likely. Spray entire lawn, wetting weeds and lawn grasses. Do not exceed recommended rates because damage to turf may occur. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Do not apply when temperature exceeds 90°F. Make a new dilution for each use. Shake well before using.

WEEDS CONTROLLED BY TRIAMINE® LAWN WEED KILLER

Alder, Artichoke, Aster, Austrian fieldcress, Beggartick, Biden, Bindweed, Bitterweed, Bitter wintercress, Black medic, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckhorn, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common sowthistle, Creeping jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hawkweed, Healall, Heartleaf drymary, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Indigo, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarter, Locoweed, Lupine, Mallow, Marshelder, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Nutgrass, Orange hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, St. Johnswort, Sheep sorrel, Shepherdspurse, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spurge, Spurweed, Stinkweed, Stitchwort, Sumac, Sunflower, Sweet clover, Tarweed, Thistle, Toadflax, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Woodsorrel, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds.

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. Some hard-to-kill (or control) broadleaf weeds may require a second broadcast treatment in 3 to 4 weeks. In the event further application is necessary, use as a spot treatment.

ORNAMENTAL LAWNS AND TURF

(ART 第二) (AR W. 1) Cool Season Grasses - such as Bluegrass, Fescue, and Rye: To kill Dandelion, Wild onion, Pennywort, Plantain, Healall and similar weeds: use 8 to 13 ounces in 6 gallons of water and apply with pump or hose-end sprayer or sprinkling can to 2,500 square feet of lawn. For smaller areas - use 3 teaspoons in 2 quarts of water applied to 100 square feet. For larger areas - use 1 to 1-3/4 gallons in 50 gallons of water for 43,500 square feet.

To kill undesirable weeds such as Thistle, Bindweed, Dock and Ragweed, use 8 ounces in 6 gallons of water (or 3 teaspoonfuls in 2 quarts of water), and apply directly to plants, wetting them thoroughly. For larger areas, use 1 to 1-1/4 gallons in 50 gallons of water.

To kill undesirable plants such as Poison ivy and Poison oak, use 2 ounces in 2 quarts of water or 2 gallons in 12-1/2 gallons of water and drench plants when they are fully leaved. Repeat in about 4 weeks if necessary. Certain hard-to-kill(or control) weeds

such as Poison ivy, Poison oak and English daisy may require a Fall application.

Warm Season Grasses - St. Augustine, Bahia, common Bermuda, Centipede, Zoysia, also for BENT: Use a one-half rate which is 4 to 6-1/2 ounces in 6 gallons of water to treat 2,500 square feet (3 teaspoonfuls in 4 quarts of water applied to 200 square feet). Make two applications 10 days apart. Do not spray grasses if stressed from heat, drought, etc. Slight turf yellowing should disappear after about I week. Note: Some hybrid Bermuda grasses may be sensitive to this product. Contact your local de l'accident de la company de Extension Service weed control specialist.

Roadsides (including aprons and guard rails), rights-of-way, drainage ditchbanks and other similar non-crop areas: For control of broadleaf weeds, mix at a rate of 1-1/3 to 4 gallons of Triamine Lawn Weed Killer per 50 to 300 gallons of water. This mixture will cover one acre. Thoroughly saturate all weeds with spray mixture. Apply when plants come into full leaf (Spring) to the time plants begin to go dormant. Best results are obtained when weeds are young and actively growing. Do not cut weeds until herbicide has translocated throughout the plant causing root death." For small broadleaf weeds, use the lower rate. Heavy, dense stands require the higher rate with high water volume.

For Spot Treatment: Put 3 teaspoons Triamine Lawn Weed Killer into an empty quart container with trigger sprayer or applicator cap. Then add 30 oz. of tap water. Trigger Sprayer - Adjust spray nozzle to give coarse spray. Aim at center of weed and spray to wet. Applicator Cap - Hold applicator end about 6 inches from surface to be treated and thoroughly wet. To obtain wider coverage, raise container. One application should be sufficient. Effects begin to show after a few days and weeds gradually die. A repeat application may be required in 3 to 4 weeks for hard-to-kill (or control) weeds. LISMO COMPONE, CRANCE CO.

STORAGE AND DISPOSAL
STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Store at temperatures above 32F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well ventilated area. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not store near open containers of fertilizer, seed or other pesticides. Do not contaminate water, food or feed by storage or disposal. THE WAR WELL PLANE TO THE WAR IN

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate ground water. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and local authorities. Plastic containers are also disposable by incineration, or if allowed by State and local authorities by burning. If burned, stay out of smoke.

Riverdale Triamine Lawn Weed Killer

(Commercial)

Page 5 6/7

WARRANTY

Seller warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant and soil conditions, mixtures with other chemicals, and factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

(RV 073103A)



Regulatory Affairs Department 1333 Burr Ridge Parkway, Suite 125A

Burr Ridge IL 60527-0866 Office Ph: 708 754 3330 Facsimile: 708 754 0314

UPS - Overnight November 10, 2003

Document Processing Desk (NOTIF)
Registration Division (7505C)
Office of Pesticide Programs
U.S. Environmental Protection Agency
Crystal Mall Building, #2
1921 Jefferson Davis
Arlington, VA 22202

Dear Sir/Madam:

Subject: Riverdale Triamine Lawn Weed Killer

EPA Registration No. 228-181 Application For Pesticide: Other OPP Identifier No.: 292582

We wish to notify the Agency we have added another sprayer instruction and alternate rates to the above subject label.

We have also enclosed an extra copy of the label highlighted to show just the changes contained in this submission.

If there is anything else we should do at this time please contact us immediately.

Thank you for your continuing assistance.

Sincerely yours,

RIVERDALE - A NUFARM COMPANY

By Russell F. Sawyer

Regulatory Affairs Manager

RFS:jb

Enclosure