(Non-Commercial Label)

RIVERDALE ®

TRIAMINE®

SPOT WEED KILLER

READY TO USE

ACCEPTED JUN 30 2003

Under the Federal Insecticide, Fungicide, and Rodenticide Act. as amended, for the posticide

CONTROLS (OR KILLS) DANDELIONS, CLOVER, CHICKWEED AND MANY OTHER LISTED WEEDS ON LAWNS AND ORNAMENTAL TURF

GET THE OPTICAL ADVANTAGE®

ACTIVE INGREDIENTS: Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*
INERT INGREDIENTS:
Isomer Specific Method, Equivalent to: *2,4-Dichlorophenoxyacetic Acid

Riverdale, Triamine and The Optical Advantage are Registered Trademarks of Riverdale - A Nufarm Company

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANELS FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 228-183

NET CONTENTS 32 FL. OZ. (1 QT.)

EPA EST. NO. 228-IL-1

MANUFACTURED BY NUFARM AMERICAS, INC., BURR RIDGE, ILLINOIS 60527-9866

Page 2

(Left Side Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Avoid inhalation of spray mist and contact with eyes or clothing. Harmful if swallowed or absorbed through skin. Causes moderate eye irritation. Wear long pants, long-sleeved shirt, socks, shoes and chemical resistant gloves when handling or applying this product. After using this product, wash nondisposable gloves thoroughly with soap and water before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower. Do not allow people (other than applicator) or pet on treatment area during application. Do not enter treatment areas until spray has dried.

FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES:

Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters nor pour the washwater on the ground; spray or drain over a large area away from wells and other water sources and desirable plants. Do not contaminate water used for irrigation or domestic purposes. If spills occur, collect the material and dispose of by following disposal instructions on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

USE PRECAUTIONS

IMPORTANT: Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray exposed roots of desirable plants. Do not spray on Dichondra, Lippia, nor on grasses where desirable Clovers are present. Do not apply to newly seeded grasses until they have been moved 3 times. Do not cut grass 1 to 2 days beter and after application. Do not reseed for 3 to 4 weeks after use. If lawn needs watering, water thoroughly before application. Do not water for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours. Avoid fine mists. Do not exceed recommended rates because damage to turn may occur. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Do not apply when temperature exceeds 90°F. Shake well before using.

WEEDS CONTROLLED (OR KILLED) BY TRIAMINE SPOT WEED KILLER

Alders, Artichoke, Aster, Austrian fieldcress, Beggartick, Biden, Bindweed, Bitterweed, Bitter winter cress, Black medic, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckhorn, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common

sowthistle, Creeping jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hawkweed, Healall, Heartleaf drymary, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarters, Locoweed, Lupines, Mallow, Marshelder, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Nutgrass, Orange hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, St. Johnswort, Sheep sorrel, Shepherdspurse, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spurge, Spurweed, Stinkweed, Stitchwort, Sumac, Sunflower, Sweet clover, Tarweed, Thistle, Toadflax, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Woodsorrel, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds.

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. One application should be sufficient. Some hard-to-control(or kill) broadleaf weeds may require a second (or an additional) treatment in 3 to 4 weeks.

HOW TO USE: Trigger Spray - Adjust spray nozzle to give coarse spray. Aim at center of weed and spray to wet. Applicator Cap - Hold applicator end about 6 inches from surface to be treated and thoroughly wet. To obtain wider coverage raise container.

TO PREPARE REFILL: Add 3 teaspoons (1/2 oz.) Riverdale Triamine Lawn Weed Killer to this empty quart container. Then add 30 oz. tap water and use according to directions in this label.

STORAGE AND DISPOSAL

STORAGE: Keep from freezing. To be stored in original container and placed in areas in-accessible to children.

DISPOSAL: If empty - Do not reuse this container except as specified in directions to prepare refill. Place in trash or offer for recycling if available. **If partly filled** - If product cannot be used as directed, or offered to others for their use, call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

NOTICE: Buyer assumes all risks of use, storage and handling of this product not in trict accordance with directions given herewith. (RV 022103)

ALTERNATE INSTRUCTIONS FOR "PUMP & SHOOT" SPRAY BOTTLE

Shake before using. Remove cap and fill Pump & Shoot spray bottle with product to within about one inch of top. Replace sprayer cap. Pump the spray cap slowly 8 to 10 times by alternately pulling the cap up and pushing it down. Once the spray bottle is pumped up, follow label instructions for application directions. Once pumped up, the spray bottle will shoot a straight stream up to 20 feet. Spray water through the sprayer after each use. The sprayer should be rinsed thoroughly and pumped empty for storage.

Optional CLI languaget:

Dimethylamine Salt of 2,4-D*
Dimethylamine Salt of Mecoprop-p**
Dimethylamine Salt of Dichlorprop-p***

*2,4-D Acid

**Mecoprop-p Acid

***Dichlorprop-p Acid

† Note: Some states may still require the full chemical name(s).

Optional Marketing Hype:

KILLS 140+ WEED TYPES STARTS WORKING OVERNIGHT Kills weeds, not the lawn Kills roots Kills unwanted weeds in your lawn