

# Riverdale

## WEEDSTROY TRI-ESTER™

**A THREE-WAY POST-EMERGENT SELECTIVE BROADLEAF HERBICIDE CONTAINING THE ESTERS OF 2,4-D, MECOPROP AND DICHLOFPROP. FOR USE ON GOLF COURSES, PARKS, HIGHWAYS, SOG FARMS, ORNAMENTAL TURF LAWNS AND SIMILAR NON-CROP AREAS. KILLS DANDELIONS, CHICKWEEDS, PLANTAINS, OXALIS, SPURGE AND MANY OTHER BROADLEAF WEEDS, SOME OF WHICH ARE LISTED ON THIS LABEL.**

**ONE GALLON COVERS UP TO 4 ACRES**

**NOT REVIEWED**  
In Accordance with PR Notice 82-2,  
Based on Draft Labeling Data

ACTIVE INGREDIENTS	
Isocetyl ester of 2,4-dichlorophenoxyacetic Acid*	23.90%
Isocetyl ester of 2-(2-methyl-4-chlorophenoxy) propanoic Acid**	24.41%
Isocetyl ester of 2-(2,4-3-chlorophenoxy) propanoic Acid***	3.54%
INERT INGREDIENTS	
	18.00%
<b>TOTAL</b>	<b>100.00%</b>

By Isomer Specific Assay ADAC Method, Equivalent to:  
\*2,4-dichlorophenoxyacetic acid 15.9%, 1.4 lbs/gal  
\*\*2-(2-methyl-4-chlorophenoxy) propanoic acid 15.9%, 1.4 lbs/gal  
\*\*\*2,4-dichlorophenoxy propanoic acid 22.7%, 2.0 lbs/gal

Weedstroy™ and Tri-Ester™ are Trademarks of Riverdale Chemical Co. Patent Applied For

**KEEP OUT OF REACH OF CHILDREN**  
**CAUTION**  
**SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS**

EPA REG. NO. 230-106 NET CONTENTS GAL. EPA EST. NO. 230-B-1

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION**  
Harmful if inhaled, absorbed or absorbed through skin. Avoid breathing vapor or spray mist. Avoid contact with skin, eyes or clothes. Do not contaminate clothing and wash before reuse.  
**STATEMENT OF PRACTICAL TREATMENT**  
If SWALLOWED Call a physician or Poison Control Center. DO NOT induce vomiting or give anything by mouth to an unconscious person.  
If ON SKIN Wash thoroughly with plenty of soap and water. Get Medical Attention.  
**ENVIRONMENTAL HAZARDS**  
Do not apply directly to water or wetlands. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment or disposal of waste. Do not contaminate domestic or irrigation waters.  
**PHYSICAL OR CHEMICAL HAZARDS**  
Do not use in close proximity to open flames.

**DIRECTIONS FOR USE**  
It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire label before using this product.

**GENERAL INFORMATION**  
WEEDSTROY TRI-ESTER is a stable solution containing a mixture of three phenoxy herbicides. It is for use on Lawns, Ornamental Turf (not pastures) and Golf Course Greens, Tees and Roughs, Parks, Sod Farms, Cemeteries, along Highways, Rights-of-Way, Drainage Ditchbanks and other similar non-crop areas.

**CARE MUST BE USED TO AVOID DRIFT OF SPRAY MISTS TO NON-TARGET** susceptible plants such as cotton, vegetables, flowers, ornamentals, shrubs, trees, and other desirable plants. Do not spray near these plants. Do not spray on crop grass, dandelion, nor on grounds where dandelion flowers are present. Use a low speed sprayer with coarse spray as wind drift is less likely. Avoid the rain. Do not spray roads or desirable plants. Do not exceed specified application rates for any site. Do not apply to newly seeded grounds until well established. Do not spray when air temperature is greater than 80°F. Follow no water than three to four weeks after application of this product. Clean and rinse spray equipment only up and down and water thoroughly before reuse for other sprays. Failure to do so will cause serious injury.

**WEEDS CONTROLLED BY WEEDSTROY TRI-ESTER**  
Burdock, Black Medic, Buckhorn, Bucktail, Cleome, Chickweed, Clover, Cock  
Sandwich, English Daisy, Ground Ivy, Heart-As, Henbit, Knotweed, Lamb's-ear  
Lamb's-ear, Madia, Madia, Morning Glory, Poppygrass, Pigweed, Plantain, Pursue  
by, Purple Oak, Purslane, Ragwort, Red Clover, Red (stem) Sorrel, Oxalis (wild  
Sweet, Shepherd's Purse, Spurge, Spurge, Spurge, White Clover, Wild Cow  
Wild Garlic, Wild Green, Wild Lettuce, Yarrow and many other broadleaf weeds.

**PREPARATION OF THE SPRAY**  
Add one-half the labeled amount of water to the concentrate. Then add the remainder.

### ORNAMENTAL LAWNS AND TURF

**Cool Season (Green Other Than Bent Grass)**  
Maximum kill of weeds will be obtained from spring or early fall application when weeds are actively growing. For normal applications, use WEEDSTROY TRI-ESTER at a rate of 2-3 pints in 20 to 200 gallons of water per acre or 1/4 to 1 1/2 ounces in 1 to 6 gallons of water per 1000 square feet as needed. Spray when air temperature is between 50-80°F. Avoid spraying during excessively dry or hot periods unless irrigation is used. Do not reapply within 24 hours of application. Do not use on Dichondra, carpet grass, or St. Augustine. Do not mow site of two days before or after application.

**Controlled Drift Application**  
Reduced rates of WEEDSTROY TRI-ESTER must be used when grass is stressed from frost, drought, etc. Do not use on areas stressed severely. Add 1/4 pint WEEDSTROY TRI-ESTER to the herbicide container than 1/4 with 3 parts of water, keeping container agitated. Spray when conditions over 35,000 square feet (approx. 1/2 acre). Do not overlap between spray patterns.

### ROADS, RIGHTS-OF-WAY, DRAINAGE DITCH BANKS AND OTHER SIMILAR NON-CROP AREAS

A control of broadleaf weeds, use at a rate of 1/4 to 1 gallon of WEEDSTROY TRI-ESTER per 100 to 300 gallons of water. This mixture will cover one acre. Thoroughly saturate all weeds with spray mixture. Apply when plants come into full leaf (spray to the time plants begin to go dormant). Best results are obtained when weeds are young and actively growing. Do not cut weeds until the herbicide has translocated throughout the plant causing root death. For small broadleaf weeds, use the lower rate. Heavy dense stands require the higher rate with high water volume. For small applications with small tank sprayers use at the rate of 1 ounce to 1/2 ounce WEEDSTROY TRI-ESTER per gallon of water.

### STORAGE AND DISPOSAL

**STORAGE:** Always store pesticides in a secure warehouse or storage building. Do not store near acids, fertilizers, explosives or combustibles. Containers should be opened in well-ventilated areas. Keep container tightly closed when not in use. Do not stack cardboard cases more than two pallets high. Do not contaminate water, food or feed by storage or disposal.  
**PERSONAL PROTECTION:** Pesticides contain are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with pebbles or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixture, or dilute is a violation of Federal law if these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the nearest State representative of the national EPA Regional Office for guidance.  
**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then allow for recycling or recondition or purchase and dispose of in a sanitary landfill, or by other procedures approved by State and Local authorities. Plastic containers dispose also by incineration, or if allowed by State and Local authorities by burning if burned, stay out of water.

### WARRANTY

Riverdale warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, the herbicide is warranted to be an effective weed killer. Shows ground, method of application, weather, soil, and soil conditions, mixture with other chemicals, and other

PM 23 228-125