


929


**Dexol Industries  
Sprayer Instructions**

**UAP & FORD's  
Sprayer Instructions**



**ASSEMBLY INSTRUCTIONS**


1. Attach siphonage tube to hose-end sprayer attachment. 2. Remove cap from bottle and insert sprayer attachment. Securely tighten sprayer attachment on bottle. 3. Screw hose into hose-end sprayer attachment.




**SPRAYING INSTRUCTIONS**

1. Turn on water. **TO START FLOW OF PRODUCT:** Place finger over hole on top right side of sprayer attachment. Hold about 3 feet above area to be treated moving at a steady pace to ensure consistent coverage. 2. **TO STOP FLOW OF PRODUCT:** Remove finger from hose to break siphon action. **NOTE:** To ensure adequate coverage, use moderate to high water pressure.

**HOW TO USE SPRAYER**




Connect spray nozzle to hose. Rubber washer should be checked for proper position. Determine area to be treated. For example, 500 sq. ft. = an area measuring 10 ft. x 50 ft. When applying, check area being treated against measuring guide on side panel. Spray pattern can be tested (without discharging product) by not placing finger over hole on right side of nozzle.



Remove bottle cap and screw bottle into nozzle (save bottle cap for storage after use). Check siphon tube for proper position. Do not squeeze bottle while attaching nozzle, or during application as contents may discharge.

To store unused product: Turn off water supply at tap. Remove bottle from sprayer and replace original bottle cap. Remove nozzle from hose and rinse in clear water.



Hold sprayer attached to hose in one hand and turn on the water at the tap. Clear water will discharge when water is turned on at tap. Let clear water run until all air is discharged from hose. When ready to treat, place finger over hole on right side of nozzle and this product will mix automatically at the proper ratio. To stop flow of product, remove finger from hole to break siphon action.


While holding sprayer at waist level, point in direction away from face and body. Walk back and forth at a brisk, steady pace. Overlap slightly for uniform coverage. Marks on right side panel indicate square feet coverage. For certain areas of lawn, it may be more convenient to stand still and direct spray across treated area in a broad sweeping motion.

**NOTE:**

Read and follow directions and precautions for each E-Z Flo product.

Do not interchange nozzles from one E-Z Flo product to another.

E-Z Flo is a T.M. of United Agri Products, Inc.



# Riverdale

C H E M I C A L C O M P A N Y

PM 23 028-186

TELEPHONE (800) 345-3330 1210  
(708) 754-3330  
FAX (708) 754-0314

OFFICE: 425 WEST 194th STREET, GLENWOOD, IL 60425-1584

PLANT: 220 EAST 17th STREET, CHICAGO HEIGHTS, IL 60411-3699

April 30, 1993

Ms. Joanne I. Miller  
Product Manager (23)  
Registration Division - H7505C  
U.S. Environmental Protection Agency  
Crystal Mall Bldg. #2  
1921 Jefferson Davis Highway  
Arlington, VA 22202

Dear Ms. Miller:


Subject: 2,4-D Exposure Reduction Program Label Revisions Due 3/1/93  
Label Corrections

Following the submission of approximately sixty (60) labels, we conducted an audit and found, for more reasons than we care to go in to, it became necessary to correct the majority of the labels previously submitted.

We have made two copies of each corrected label. We respectfully request that they be entered in to our registration file as replacements. We apologize for any inconvenience this may cause the Agency.


Sincerely yours,

RIVERDALE CHEMICAL COMPANY

  
By Russell F. Sawyer  
Regulatory Affairs Manager

RFS:ls

Enc.



**CPDA** Chemical Producers  
and Distributors  
Association

  
PLCAA

RIVERDALE

1D + 1DP LOW VOL

A WOODY PLANT HERBICIDE FOR CONTROLLING MIXED BRUSH ON UTILITY RIGHTS-OF-WAY, HIGHWAYS, DRAINAGE DITCHBANKS, FENCE ROWS AND SOLID STANDS OF OAK OR ELM. CONTROLS NOXIOUS PERENNIAL WEEDS ON NON-CROPLAND.

ALSO CONTROLS NOXIOUS PERENNIAL WEEDS ON UNCROPPED LAND, LAWNS, GOLF COURSES, ATHLETIC FIELDS, PARKS, AND OTHER ORNAMENTAL TURF GRASS AREAS.

ACTIVE INGREDIENTS:

|  |        |
|--|--------|
| Isooctyl (2-ethylhexyl) Ester of 2,4-Dichlorophenoxyacetic Acid*         | 16.05% |
| Isooctyl (2-ethylhexyl) Ester of 2-(2,4-Dichlorophenoxy)propionic Acid** | 16.10% |

|                       |         |
|-----------------------|---------|
| INERT INGREDIENTS:*** | 67.85%  |
| TOTAL                 | 100.00% |

Isomer Specific by AOAC Method, Equivalent to:

|   |       |               |
|---|-------|---------------|
| *2,4-Dichlorophenoxyacetic Acid         | 10.6% | .92 lbs./gal. |
| **2-(2,4-Dichlorophenoxy)propionic Acid | 10.9% | .94 lbs./gal. |

\*\*\*This product contains Petroleum Distillates

KEEP OUT OF REACH OF CHILDREN

CAUTION - CAUCION

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

NET CONTENTS GAL.


EPA REG. NO. 228-186

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584




NOTE: Spanish language is optional

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Avoid inhalation of vapors or spray mist and contact with eyes, skin, or clothing. Wash



3210  
 nondisposable gloves thoroughly with soap and water before removing. Remove saturated clothing as soon as possible and shower.

If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron. If this container is five gallons or more in capacity, then a mechanical system (probe and pump) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

Please see additional Precautionary Statements in Directions For Use instructions.

#### STATEMENT OF PRACTICAL TREATMENT

- IF SWALLOWED: Do not induce vomiting. If vomiting occurs spontaneously, keep airway clean. Never give anything by mouth to an unconscious person. Get medical attention.
- IF IN EYES: Immediately flush eyes with plenty of water for at least 15 minutes. Get medical attention.
- IF ON SKIN: Wash exposed skin gently with plenty of soap and water.

#### ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or run-off may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. When cleaning equipment, do not pour washwaters on the ground, spray or drain over a large area away from wells and other water sources. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system. Do not contaminate irrigation ditches or water used for domestic purposes. If spills occur, collect the material and dispose of by following disposal instructions on this label. Use care to avoid spray contact or drift to 2,4-D or 2,4-DP susceptible plants such as okra, cotton, tomatoes, flowers, grapes, fruit trees, vegetables and ornamentals. Do not permit spray mist containing this product to drift onto them since even very small quantities of the spray, which may not be visible, can cause severe injury during both growing and dormant periods. Do not spray when the wind is blowing towards susceptible crops or ornamental plants. Use coarse sprays to minimize drift. Spray drift can be lessened by keeping the spray boom as low as possible; by applying 20 gallons or more of spray per acre; by using 20 pounds spraying pressure with flat fan or flooding flat fan nozzle tips; by spraying when wind velocity is low; and, by stopping all spraying when wind exceeds 6 to 7 miles per hour. Do not apply with hollow cone-type insecticide or other nozzles that produce a fine droplet spray. Do not use in greenhouses. Do not use the same spray equipment for applying other materials to 2,4-D susceptible crops as injury may result.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D and 2,4-DP have been associated with mixing/loading and

disposal sites. Caution should be exercised when handling these pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

#### BRUSH CONTROL

#### ADDITIONAL PRECAUTIONARY STATEMENTS FOR AGRICULTURAL CROP USES (i.e. Sod Farms and Forestry works engaged in the commercial production of wood fiber or timber products including forestry site preparation)

When mixing, loading or applying this product or repairing or cleaning equipment used with this product, wear eye protection (face shield or safety glasses), chemical resistant gloves, long-sleeved shirt, long pants, socks and shoes. It is recommended that safety-glasses include front, brow and temple protection. For aerial applicators in an enclosed cockpit and applicators applying this product from a tractor that has a completely enclosed cab, eye protection is not required. Wash hands, face and arms with soap and water as soon as possible after mixing, loading or applying this product. Wash hands, face and arms with soap and water before eating, smoking or drinking. Wash hands, face and arms before using toilet. After work, remove all clothing and shower using soap and water. Do not reuse clothing worn during the previous day's mixing and loading or application of this product without cleaning first. Clothing must be kept and washed separately from other household laundry. There isn't any restriction on the annual maximum number of broadcast applications for Brush Control nor Sod Farms. Do not enter or allow worker entry into treated areas during the restricted-entry interval of 12 hours. No re-entry restriction for non-agricultural uses for Brush Control.

#### BRUSH CONTROL

RIVERDALE 1D + 1DP LOW VOL is specifically designed for utilities and other industrial users to control woody plants. This product also controls many noxious perennial weeds on uncropped land, such as along highways, drainage ditchbanks and fence rows. The following is a partial list of the weeds controlled: Alder, Ash\*, Aspen, Birch,

Blackberry, Black cherry, Black jack oak, Black locust, Box elder, Brambles, Buckbrush, Ceanothus, Chamise, Coffeeberry, Currant, Dewberry, Dogwood, Elderberry, Elms, Fir, Gooseberry, Greenbrier, Gum, Hazel, Hemlock, Hickory, Honeysuckle, Kudzuvine, Locust, Manzanita, Maples\*, Multifloral rose, Oaks, Osage orange, Palmetto, Persimmon\*, Pine, Poison ivy, Poison oak, Poplar, Raspberry, Red elder, Salmonberry, Sand sagebrush, Sassafras, Serviceberry, Shinnery oak, Snowberry, Spicebush, Spruce, Sumac, Sweetgum, Sycamore, Tulip poplar, Virginia creeper, Wild cherry, Wild grape, Wild rose, Willow,

5210

Winged elm, Yerba santa, and many other species.

\*Best controlled by stump or basal treatment.

**TO PREPARE SPRAY:** Add one-half the required amount of oil (kerosene, diesel or fuel oil) or water to the spray tank, then add this product with agitation and finally the balance of the water or oil with continued agitation. This material forms an emulsion in water not a solution. This tends to separate on standing. Provide agitation to prevent such separation and ensure a uniform mixture. If this material is to be used in straight oil mixtures, do not let water get into it or the finished mixture.

Note: Water and a surfactant (spreader sticker) may be substituted in amounts equal to recommended amounts of oil.

**FOLIAGE STEM TREATMENT:** This is the standard method for high-volume sprays along fence rows, highways and rights-of-way. Use as a spray on thick brush composed of mixed species. Apply to both stems and foliage from the time foliage is completely matured until the plants start to go dormant. All leaves, stems and suckers must be completely wet to the ground line for effective control. Some regrowth may be anticipated on the more resistant species, such as oak, maple, ash and persimmon. Add 1-3/4 to 3 gallons of RIVERDALE 1D + 1DP to 100 gallons of water using 200-600 gallons of spray mixture per acre (optional: 2/3 to 1 pint of Riverdale 1D + 1DP to 4 gallons of water using 8-1/2 to 25 gallons of spray mixture for 1800 sq. ft.), depending upon the height and thickness of the brush. Mix thoroughly before spraying.

**SPOT TREATMENT:** For spot spraying with backpack sprayer mix 2 cups (16 oz.) RIVERDALE 1D + 1DP with 5 gallons of water. Wet brush stems and foliage thoroughly.

**BASAL BARK TREATMENT:** Thoroughly wet the base and root collar of all stems until the spray accumulates around the root collar at the ground line. This spray may be applied during any season. Use RIVERDALE 1D + 1DP for basal bark treatment on scattered brush or as a second spray application on species resistant to first foliage application. Mix 6-8 gallons of RIVERDALE 1D + 1DP in 100 gallons of oil. Apply with a low volume sprayer or power equipment. Application rate will depend on species present, season applied and volume of spray used. Use a coarse spray to avoid drift.

**MODIFIED BASAL TREATMENT:** Drench the base of plants, then wet the lower 4/5 of remaining stems and leaves thoroughly to runoff. Apply treatment when brush is in full foliage. This method can be applied where susceptible species have been controlled by prior sprays and more resistant species, such as maple and oak, remain. Soaking the base of the plant and wetting all stems to runoff is absolutely necessary for complete control.

**EARLY SEASON SPRAYING:** Add 2-3 gallons of RIVERDALE 1D + 1DP to 10 gallons of diesel oil and thoroughly mix. Add this mixture to 89 gallons of water.

**DURING DRY WEATHER OR THE LATTER PART OF SPRAYING SEASON:** Add 3 gallons of RIVERDALE 1D + 1DP to 15 gallons of diesel oil and mix thoroughly. Add mixture to 83 1/2 gallons of water and agitate thoroughly before use to ensure uniform mixing. DO NOT allow mixture to stand more than 1 hour after mixing.

**CUT SURFACE TREATMENT - STUMPS:** This treatment may be used anytime of the year; however, it is more effective when applied as quickly as possible after trees are cut. Spray the entire stump, especially exposed roots and bark. A complete soaking is essential for effective control. Use this procedure after original or capital removal. It is the first step towards a chemical brush control program on newly cleared highways

and rights-of-way. The spray is most effective and profitable on stumps 3-4 inches and larger. Mix 6 to 9-1/2 gallons of RIVERDALE 1D + 1DP in 100 gallons of oil. Application should be made with a low volume knapsack sprayer using a solid cone-shaped nozzle medium orifice or a sprinkler can.

**CONCENTRATE STUMP TREATMENT:** For small (up to 3 inch diameter) stems, cut them as close to the ground as possible and apply undiluted RIVERDALE 1D + 1DP directly from the can to the surface of the freshly cut stump.

**FRILL:** Make a frill using an axe to cut overlapping V-shaped notches in a continuous ring, and cut around the trunk near its base. Cut through the bark, but do not remove the chips. This method is recommended for all trees 5-6 inches in diameter and larger. Freshly cut frills can be treated anytime of the year. Mix 6 to 9-1/2 gallons of RIVERDALE 1D + 1DP in 100 gallons of oil (optional: 1 pint of Riverdale 1D + 1DP in 2 gallons of oil). Pour in as much of the mixture as the frills will hold without wasting the chemical.

**POWERED KNAPSACK BASAL SPRAY:** Mix 16-28 gallons of RIVERDALE 1D + 1DP with fuel oil or kerosene to make 100 gallons of spray solution. Apply with a portable knapsack mistblower to lower brush stems. Apply spray to all sides of stems; good root collar coverage is essential. Run knapsack mistblower at 1/4-1/3 throttle for best spray delivery and coverage. For maximum drift control use a basal nozzle attachment. Do not raise spray nozzle above horizontal position.

**FENCE ROW APPLICATION:** To control mixed brush, perennial and annual broadleaf weeds, use one of the application methods such as the foliage stem method described on this label. Some regrowth may be expected on resistant species such as ash, maple, oak and persimmon.

**LOW VOLUME STEM FOLIAGE SPRAY - AERIAL OR GROUND:** Apply the spray only through equipment designed to provide effective drift control.

**MIXED BRUSH - UTILITY RIGHTS-OF-WAY:** For aerial application to solid stands of brush use 4-8 quarts in 3-12 gallons volume per acre. 1-4 quarts of fuel oil may be included in this mixture.

**SOLID STANDS OF OAK OR ELM:** Apply with fixed-wing or helicopter aircraft. Apply in the Spring after hardwoods have just developed full sized leaves. The spray season normally runs from early May to mid-June in Texas and California, and from early May to early July in Oklahoma and northward. Spray when wind velocities are less than 5 miles per hour.

| Brush                              | 1D + 1DP     | Oil      | Add Water to Make Total |
|------------------------------------|--------------|----------|-------------------------|
| post, blackjack oak and winged elm | 1-1/3 gallon | 1 gallon | 5 gallons               |
| sand shinnery oak                  | 1 - 2 quart  | 1 gallon | 3 gallons               |

For maximum control, use higher rate and repeat spray the second year using 2 quarts RIVERDALE 1D + 1DP per acre.

**FORESTRY SITE PREPARATION:** To reduce competition from mixed trees and brush before planting forest trees. Apply 2 gallons RIVERDALE 1D + 1DP for easy-to-control species

7 of 10  
such as aspen, hazel alder, sassafrass, sumac, black locust, willow, and other similar species. Apply 4 gallons RIVERDALE 1D + 1DP to control more difficult species such as oaks, sweet gum, black gum, hickory and tulip-poplar.

Mix with water to make 10 gallons total solution when applying as an aerial spray or 20 gallons total solution when applying as a ground spray. Do not apply to established plantations as this spray mix will injure planted conifers.

**PINE RELEASE:** To control hardwood brush and release Northern conifers such as red pine, Jack pine, white pine, and white spruce.

Apply 4-8 quarts RIVERDALE 1D + 1DP as a broadcast spray in 10 gallons of water per acre. Make applications in midsummer after pine height growth is complete and the conifer buds are set. RIVERDALE 1D + 1DP will control aspen, birch, cherry, alder, hazel, oaks, and similar species. RIVERDALE 1D + 1DP will not, at these rates, provide satisfactory control of red maple, sugar maple or ash.

#### RIVERDALE 1D + 1DP AND BANVEL HERBICIDE<sup>(R)</sup> MIXTURES

**HIGH VOLUME STEM FOLIAGE SPRAY** - Mix 2 gallons to 3 gallons RIVERDALE 1D + 1DP with 1-2 quarts Banvel 4WS per 100 gallons total spray mix. Use the low rate for easy to kill brush such as alders, aspen, cherry, sycamore, tulip-poplar, and willows. Use the high rate to control species such as oak, ash, elm, pines, spruce, and fir.

**LOW VOLUME STEM FOLIAGE SPRAY:** Aerial or Ground - Mix 4-6 gallons of RIVERDALE 1D + 1DP with 1-1½ gallons Banvel herbicide in 15-50 gallons of water per acre to control mixed brush including conifers and ash.

Observe all restrictions, precautions and limitations on the labeling of each product used in tank mixture.

### TURF CONTROL

---

#### ADDITIONAL PRECAUTIONARY STATEMENTS FOR TURF CONTROL

When using this product, wear long-sleeved shirt, long pants, socks, shoes and chemical resistant gloves. Commercial mixers/loaders must also wear these clothes, except when the product is to be applied to golf courses. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

---

Lawns and other ornamental turf grass areas - RIVERDALE 1D + 1DP LOW VOL is recommended for professional weed control of broadleaf weeds in lawns, ornamental turf, and similar turf areas. This treatment may injure bentgrass, St. Augustinegrass, centipede grass, carpetgrass and newly seeded lawns. If necessary to control weeds in such turf, use half the recommended rate on chart and repeat application in 2 to 3 weeks. Do not apply to Lippia, Dandelion and where desirable clover are present nor on bentgrass greens and tees as severe injury may result. Do not use this product to control weeds in flower and vegetable beds, shrubs and ornamental plantings. Spray when air is calm to avoid spray drift that might injure nearby desirable plants. A

separate sprayer should be kept for killing weeds as 2,4-D and related chemicals are difficult to clean from equipment.

The following is partial list of weeds controlled by RIVERDALE 1D + 1DP

Bindweed, Black medic, Brambles, Buckhorn plantain, Bull thistle, Burdock, Buttercups, Canada thistle, Carolina geranium, Chickweeds (common and mouseear), Chicory, Clovers, Cocklebur, Cudweed, Dandelion, Dock, Evening primrose, False dandelion, Fleabane, Florida pusley, Frenchweed, Ground ivy, Hawkweed, Healall, Henbit, Honeysuckle, Jimsonweed, Knotweed, Kochia, Kudzu, Lambsquarters, Little startwort, Mallow, Morningglory, Mustards, Oxalis (yellow woodsorrel), Pennywort, Peppergrass, Pigweed, Plantains (narrow or buckhorn; broadleaf), Poison ivy, Poison oak, Poorjoe, Povertyweed, Purslane, Ragweed, Sheep sorrel, Shepherdspurse, Smartweeds, Soliva, Sowthistle, Speedwell (annual), Spurge, Sumac, Vervain, Vetch, Violet, Wild aster, Wild blackberry, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild raspberry, Yarrow.

This product is not effective on perennial Veronicas or on weed grasses. Resistant weeds such as Oxalis (yellow woodsorrel), Bindweed and Yarrow should be retreated whenever new growth appears. Control of difficult weeds such as Sheep (red) sorrel, Soliva, and Violets may be only partial.

**PREPARATION OF THE SPRAY:** Fill the spray tank with half of the required amount of water, then add the recommended amount of RIVERDALE 1D + 1DP herbicide with agitation and finally the balance of water with agitation. Apply any time broadleaf weeds are growing actively. Dandelion, plantain, wood sorrel and clovers are best treated in the Fall or in Spring before flower heads develop. Winter weeds such as chickweeds and henbit should be treated in early Spring.

Summer weeds such as oxalis, knotweed and sparges should be sprayed when they are small. Resistant weeds such as bindweed and violets may require 1½ gallons per 40,000 sq. ft. and should be retreated whenever new growth appears. In areas with extended growing seasons, such as California, treatment in both Spring and Fall may be needed to control more resistant species.

On turf to be over-seeded in the Fall, treatment should be scheduled at least 4 weeks before the planned seeding date. Rainfall (¼ inch or more) or irrigations should occur prior to seeding. Fall-seeded lawns may be treated the following Spring.

Spring-seeded lawns may be treated after the grasses have sprouted and been cut at least twice, generally 6-10 weeks after seeding, depending on germination and growth rate.

For established lawns, Fall treatment fits into a good turf management program. Proper fertilization and mowing should be combined with chemical weed control to thicken the turf after the weeds have died and to discourage more weeds from invading.

**HOSE ATTACHED SPRAYER (LAWN MODELS):** Use 3 teaspoons of RIVERDALE 1D + 1DP herbicide diluted with water to the 3-5 gallon mark on the sprayer jar for each 500 square feet (25' x 20') of lawn area to be sprayed. Adjust water pressure so that spray streams about 10-15 feet with no misting.

## AMOUNTS TO USE

| Riverdale 1D + 1DP        | Area Covered   | Volume   |
|---------------------------|----------------|----------|
| 3 oz.                     | 1,000 sq. ft.  | 1 gal.   |
| 1 pt.                     | 5,000 sq. ft.  | 5 gals.  |
| 1 qt.                     | 10,000 sq. ft. | 10 gals. |
| 1 gal.                    | 40,000 sq. ft. | 40 gals. |
| 1½ gal. (resistant weeds) | 40,000 sq. ft. | 40 gals. |

## Consumer Statement (for containers 1 gallon or less)

**STORAGE AND DISPOSAL**

**STORAGE:** To be stored in original container and placed in an area inaccessible to children.

**PRODUCT DISPOSAL:** Partially filled container may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Do not reuse empty container. Rinse thoroughly before discarding in trash.

## Commercial Applicator Statement (for containers over 1 gallon)

**STORAGE AND DISPOSAL**

**STORAGE:** Always use original container to store pesticides in a secured warehouse or storage building. Do not store near seeds, fertilizers, insecticides or fungicides. Containers should be opened in well-ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE DISPOSAL:** Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or recondition or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic containers disposable also by incineration, or if allowed by state and local authorities by burning. If burned, stay out of smoke.