

(NOTE: Home Owner Use Label)

228-187

PM 12

133

(Front Panel)

RIVERDALE

FRUIT AND BERRY SPRAY

DESTROYS INSECTS AND CONTROLS DISEASES  
COMMON TO FRUIT

ACTIVE INGREDIENTS:

Captan* .....	15.6%
Related Derivatives .....	0.4%
Malathion** .....	8.0%
Methoxychlor Technical*** .....	16.0%
INERT INGREDIENTS: .....	60.0%
	TOTAL .....
	100.0%

\*N-Trichloromethylthio-4-cyclohexene-1,2-dicarboximide

\*\*O,O-Dimethyl Dithiophosphate of Diethyl Mercaptosuccinate

\*\*\*Equivalent to 14.08% of 2,2-bis(p-methoxyphenyl)-1,1-trichloroethane and  
1.92% other isomers and related compounds

KEEP OUT OF REACH OF CHILDREN

DANGER

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS

EPA Reg. No. 228-187

NET WEIGHT 11 OUNCES

EPA Est. No. 228-IL-1

*Per sell Cert. #77*  
 DEC 9 1987  
 228-187

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, IL 60425-1584


(Back Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Causes irreversible eye damage. Do not get in eyes. Harmful if  
swallowed or inhaled. Avoid breathing of spray mist. Avoid  
contact with skin. May cause allergic skin reactions. Wash  
thoroughly after using.



STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with plenty of water. Call a physician.

IF SWALLOWED: Promptly drink a large quantity of milk, egg  
whites, gelatin solution, or if these are not  
available, drink large quantities of water.  
Avoid alcohol.

IF INHALED: Remove victim to fresh air. If not breathing  
give artificial respiration, preferably mouth-  
to-mouth. Get medical attention.

(Back Panel cont'd)

ENVIRONMENTAL HAZARDS

This product is highly toxic to bees exposed to direct treatment or residues on crops. This product is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FRUIT AND BERRY SPRAY is an effective Insecticide-Fungicide formula to control the insects and diseases of Fruits listed below.

USE PRECAUTIONS

Wear goggles or face shield when handling. Remove contaminated clothing and separately launder clothing before reuse. Wear chemical resistant gloves when using indoors or outdoors. When using wear long pants and long-sleeved shirt and apply with the wind to your back. Wash non-disposable gloves thoroughly with soap and water before removing. Wash fruit before marketing or eating.

NOTE: Spray thoroughly, covering both sides of all leaves, branches and the trunk on a regular program.

TO MIX

4 1/2 level tablespoonfuls make 1 gallon of spray  
11 oz. size makes 11 gallons of spray

APPLES AND PEARS:

Make first application when leaves begin to show. Continue at 7 day intervals until tree begins to bloom. Additional sprays should be made at Petal Fall-7 days later, and then repeated at two week intervals until 7 days before harvest.

Aphids, Mites, Japanese Beetles, Spittle Bugs, Cankerworms, Red Spiders, Leaf hoppers. Scab, Botrytis Rot, Frogeye Leaf Spot.

PEACHES, CHERRIES, APRICOTS AND PLUMS:

Apply first spray when blossoms first begin to show pink; spray again when petals fall and thereafter every 7-10 days until 7 days before harvest for cherries and plums and 21 days before harvest for peaches and apricots.

Aphids, Mites, Japanese Beetles, Spittle Bugs, Cankerworms, Red Spiders, Leafhoppers, Brown Rot, Leaf Spot.

(If a later application is needed for Fruit Rot, use Captan 50% Wettable Powder alone. Remove residue at harvest by washing or other effective means.)


STRAWBERRIES:

Spray plants at 7-10 day intervals beginning when the plants start to grow, until 14 days before harvest.

Aphids, Mites, Japanese Beetles, Spittle Bugs, Red Spiders, Leaf-hoppers, Leaf Spot and Fruit Rot.

(If a later application is needed for Fruit Rot, use Captan 50% Wettable Powder alone. Remove residue at harvest by washing or other effective means.)

X  
will  
be  
corrected  
on  
printed  
label  
E. K. Miller  
11-18-87



STORAGE AND DISPOSAL

STORAGE: Store in original container and place in a cool, dry location in an area inaccessible to children. Keep container tightly sealed when not in use. Avoid contamination of feed and foodstuffs.

PESTICIDE DISPOSAL: Partially filled containers may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash. ✓

CONTAINER DISPOSAL: Do not reuse container. Discard container in trash. ✓

NOTICE

Riverdale Chemical Company warrants that this product conforms to the chemical description on this label. Seller makes no warranty, express or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

