

228-189

ACCEPTED

01/13/2005
(Residential Label)

1/5

JAN 13 2005
Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

228-189

RIVERDALE[®]
TRIAMINE[®] LIQUID

WEED AND FEED

READY-TO-USE

FOR USE WITH A HOSE-END SPRAYER ON ORNAMENTAL TURF (GRASS) LAWNS
NO MIXING

Just Connect Hose, Turn On Water, and Spray

TREATS (COVERS) 5,000 TO 7,500 SQ. FT.
(Read directions for different grasses.)

CONTAINS 2,4-D, MECOPROP-p, DICHLORPROP-p AND A X-X-X FERTILIZER

GET THE OPTICAL ADVANTAGE[®]

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	3.26%
Dimethylamine Salt of (+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**†	1.63%
Dimethylamine Salt of (+)-R-2-(2,4-Dichlorophenoxy)propionic Acid***‡	1.60%
INERT INGREDIENTS:	93.51%
TOTAL	100.00%

Equivalent to:

- *2,4-Dichlorophenoxyacetic Acid 2.70%, 0.26 lb(s)/gal.
 - ** (+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid † 1.35%, 0.13 lb(s)/gal.
 - *** (+)-R-2-(2,4-Dichlorophenoxy)propionic Acid 1.33%, 0.13 lb(s)/gal.
- CONTAINS THE SINGLE ISOMER FORMS OF MECOPROP-P AND DICHLORPROP-p

GUARANTEED ANALYSIS:[°]

Total Nitrogen (N) X%

X% Urea Nitrogen

Fertilizer guaranteed by (Name of Fertilizer registrant).

Riverdale, Triamine & The Optical Advantage are Registered Trademarks of Nufarm Americas Inc.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE AND BACK PANELS FOR FIRST AID
AND ADDITIONAL PRECAUTIONARY STATEMENTS

NET CONTENTS 1 QUART (32 FL. OZ.)

NET WEIGHT: 2.3 LBS.

EPA Reg. No. 228-189

EPA EST. No. 228-IL-1

MANUFACTURED BY NUFARM AMERICAS INC., BURR RIDGE, ILLINOIS 60527-0866

- Revised 1/31/03 Changed First Aid Statement per PR 2001-1 and Disposal Statement per PR 2001-6 via Amendment.
- 3/15/04 Corrected the Disposal Statement via Notification.
- 4/28/04 Converted Mecoprop and Dichlorprop to Optical Isomer via Amendment.
- 6/28/04 Corrected per EPA's letter dated 5/17/04.
- 10/21/04 Corrected by Linda Kutney fax . Front panel changes only.

(Back Panel)

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear long pants, long-sleeved shirt, socks, shoes and chemical resistant gloves when handling or applying this product. After using this product, wash non-disposable gloves thoroughly with soap and water before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

(OPTIONAL: In case of emergency involving this product call collect day or night, XXX-XXX-XXXX.)

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters, nor pour the washwaters on the ground; spray or drain over a large area away from wells, other water sources and desirable plants. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes. If spills occur, collect the material and dispose of by following disposal instructions on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Spray anytime during the growing season when weeds are actively growing (except as noted on S-Southern or Warm Season grasses)

5,000 Square Feet Coverage for (N-Northern or Cool Season Grasses) such as Blue Grass, Fescue, and Rye grasses.

7,500 Square Feet Coverage for (S-Southern or Warm Season Grasses) Bahia, common Bermuda, St. Augustine, Centipede, Zoysia, and Bent grasses. Some temporary injury may occur with these grasses. In Spring, do not apply until grass has reached full green-up. In Fall, exercise care when grass growth is going into dormancy. When applying to Bent grass, height of grass should be 1-1/2 inches or more. Note: Some hybrid Bermuda grasses may be sensitive to this product. Contact your local Extension Service weed control specialist. (Optional Statement: Do not use on Floratam or other improved varieties of St. Augustine grass in Florida.)

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed this product will supplement your regular fertilizer program by providing *(No. of lbs.)* pounds of nitrogen per 1,000 square feet (optional P₂O₅ and K₂O levels) and Triamine to control listed weeds. Please contact your local extension office for questions regarding a regular fertilizer program.

*Based on Inert Fertilizer claimed.

USE PRECAUTIONS

IMPORTANT: Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray exposed roots of desirable plants. Do not spray on Dichondra, Lippia, nor on grasses where desirable Clovers are present. Do not apply to newly seeded grasses until they have been mowed 3 times. Do not cut grass 1 to 2 days before and after application. Do not reseed or plant stolons for 3 to 4 weeks after use. If lawn needs watering, water thoroughly before application. Do not water for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours. (Optional statements for adjustable sprayers - Avoid fine mists. Use a lawn type sprayer with a coarse spray as wind drift is less likely.) Spray entire lawn, wetting weeds and lawn grasses. Do not exceed recommended rates because damage to turf may occur. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Do not apply when temperature exceeds 85°F. Shake well before using.

WEEDS CONTROLLED BY TRIAMINE® LIQUID WEED AND FEED

Alder, Artichoke, Aster, Austrian fieldcress, Beggartick, Bident, Bindweed, Bitterweed, Bitter wintercress, Black medic, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckhorn, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chickweed, Chicory, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common sowthistle, Creeping jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hawkweed, Healall, Heartleaf drymary, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarter, Locoweed, Lupine, Mallow, Marshelder, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Orange hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, St. Johnswort, Sheep sorrel, Shepherdspurse, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spurge, Spurweed, Stinkweed, Stitchwort, Sumac, Sunflower, Sweet clover, Tarweed, Thistle, Toadflax, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Woodsorrel, Wormseed, Yarrow, Yellow rocket, and (many) other broadleaf weeds.

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. Some hard-to-control broadleaf weeds may require a second (or an additional) treatment in 3 to 4 weeks.

(HOSE-END SPRAYER INSTRUCTIONS)

HOW TO APPLY: Use any hose-end sprayer that is designed for use on lawns and makes a minimum of 15 gallons of finished spray. Such sprayers are Ortho's Orthomatic Lawn Sprayer, Ortho's and Black Leaf's Lawn Sprayers (dilution ratio of 60 to 1). The

following are adjustable/dial sprayers:

<u>MANUFACTURER</u>	<u>SETTING</u>
Super K-Gro	20
Melnor #94 & #95	11
Melnor #96 & #97	13
Gilmour #362	4
Gilmour #486	12

SPECIAL NOTE: Some adjustable sprayers do not hold 32 ounces. The above settings are based on using 32 ounces per 5,000 or 7,500 square feet. The entire contents of this bottle must be used to make the proper recommended application rate.

Determine area to be sprayed. Apply at a rate of 32 ounces (1 quart) per 5,000 square feet on N-Northern or Cool Season grasses or 7,500 square feet for application on S-Southern or Warm Season Grasses. Fill sprayer jar with appropriate amount of this product. Do not dilute this product with water when filling sprayer jar. The water will automatically mix with this product during spraying. Attach sprayer to garden hose while holding sprayer at waist level. Point in direction away from face and body. Walk back and forth at a steady pace. Overlap slightly for uniform coverage.

(DISPOSABLE SPRAYER AND BOTTLE INSTRUCTIONS)

HOW TO APPLY: Connect spray nozzle to hose. Remove bottle cap and screw bottle into nozzle. While holding sprayer at waist level, point in direction away from face and body. Turn on the water at the tap. Place finger over hole on right side of nozzle (or turn on/off lever to the "ON" position) and this product will mix automatically at the proper ratio. Walk back and forth at steady pace. Overlap slightly for uniform coverage.

(Left Side Panel)

COVERAGE MEASURING GUIDE

(Marks on left side panel of bottle indicate square feet covered.)
(Not to scale)

1000	Square Feet	1500
<hr/>		
2000	Square Feet	3000
<hr/>		
3000	Square Feet	4500
<hr/>		
4000	Square Feet	6000
<hr/>		
5000	Square Feet	7500
<hr/>		
Cool Season (N)		Warm Season (S)

(Right Side Panel)

STORAGE AND DISPOSAL

STORAGE: Keep from freezing. To be stored in original container and placed in an area inaccessible to children. Do not store near open containers of feed or foodstuffs.

DISPOSAL: If empty - Do not reuse this container, except as a hose-end sprayer bottle only when using Riverdale Triamine Liquid Weed and Feed refill size. Place in trash or offer for recycling if available. **If partly filled** - If product cannot be used as directed, call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

NOTICE

Buyer assumes all responsibility for safety and use not in strict accordance with directions and instructions for use. (Optional statement - If these terms are not acceptable, return at once unopened.) (RV 042804A)

The following note will not be part of the actual label copy.

^oNote: Guaranteed (fertilizer) analysis and weight will vary among various commercial fertilizer manufacturers.

(Additional copy for refill size)

For refill only: Use this bottle to refill the quart hose-end sprayer of Riverdale Triamine Liquid Weed and Feed. Follow directions for use on hose-end sprayer bottle (Optional statement - and directions pamphlet enclosed with 1 quart bottle.)

(GWF SPRAYER INFORMATION)

(Optional statements - For Residential Use Only
- For more information about Seller's products call 1-800-XXX-XXXX)

Nozzle Instructions

HOW TO APPLY

- No Mix, No Measure
- Make sure ON/OFF valve is in OFF position.
- Shake well before applying.
- Connect spray nozzle to hose.
- Turn on the water at the tap.
- When ready to treat, point in direction away from face and body and pull the tab toward the hose end connection. Turn the ON/OFF valve to the ON position. The water mixes automatically with this product at the proper ratio.
- Walk and spray at a steady pace. Marks on side of bottle indicate the number of square feet covered in 1,000's.