

Reg # 228-190

PM-23

194

EEB 08 1991

Mr. Russell F. Sawyer
 Riverdale Chemical Company
 425 West 194th Street
 Glenwood, IL 60425-1584

Dear Mr. Sawyer:

Subject: Label Amendment- Add specific weed names
 Revise the precautionary statements
 Riverdale Triamine Jet-Spray Spot Weed Killer
 EPA Registration Number 228-190
 Your Submission Dated July 11, 1991

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable provided you add the following:

1. The Caution statement, "Harmful if swallowed or absorbed through skin. Wear goggles, long pants, long-sleeved shirt, shoes and chemical resistant gloves when handling or applying this product. Wash nondisposable gloves thoroughly with soap and water before removing. If on skin, wash with plenty of soap and water. Do not allow persons and pets on treated areas until sprays have dried."

- Please note, this Caution statement is the same as on the labeling submitted with your Certification of Compliance dated October 7, 1990.

2. The method number used in your most recent Isomer Specific AOAC Method testing.

A stamped copy is enclosed for your records. Please submit three (3) copies of your final printed labeling incorporating this amendment and the above changes.

Sincerely yours,

BEST AVAILABLE COPY

Joanne I. Miller
 Acting Product Manager(23)

CONCURRENCES Fungicide-Herbicide Branch

SYMBOL	H7505C				Registration Division (H7505C)	
SURNAME	ROBBINS					
DATE	2/6/91					
Enclosure						

ACCEPTED
with COMMENTS
in EPA Letter Dated:

Feb. 8, 1991

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
228-190

(Front Panel)

RIVERDALE
TRIAMINE™
JET-SPRAY
SPOT WEED KILLER
READY TO USE

CONTROLS DANDELIONS, CLOVER, CHICKWEED AND
MANY OTHER LISTED WEEDS ON LAWNS AND ORNAMENTAL TURF

ACTIVE INGREDIENTS:†

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*326%	<i>2,4-D PROP 2,4-DP</i>
Dimethylamine Salt of 2-(2-Methyl-4-chlorophenoxy)propionic Acid**328%	
Dimethylamine Salt of 2-(2,4-Dichlorophenoxy)propionic Acid***324%	
INERT INGREDIENTS:	99.022%	
TOTAL	100.000%	

Isomer Specific AOAC Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid27%	.023 lbs/Gallon
**2-(2-Methyl-4-chlorophenoxy)propionic Acid27%	.023 lbs/Gallon
***2-(2,4-Dichlorophenoxy)propionic Acid27%	.023 lbs/Gallon

†TRIAMINE IS A TRADEMARK OF RIVERDALE CHEMICAL COMPANY
Patent Pending

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET WEIGHT 1 lb. 7 oz.

EPA REG. NO. 228-190

EPA EST. NO. 33994-IL-2

Manufactured By
RIVERDALE CHEMICAL COMPANY
Glenwood, Illinois 60425-1584

(Left Side Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. If on skin, wash with plenty of soap and water.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes).

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Exposure to temperatures above 130°F may cause bursting.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

USE PRECAUTIONS

IMPORTANT: Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray exposed roots of desirable plants. Do not spray on Dichondra, Lippia, nor on grasses where desirable Clovers are present. Do not apply to newly seeded grasses until they have been mowed 3 times. Do not cut grass 1 to 2 days before and after application. Do not reseed for 3 to 4 weeks after use. If lawn needs watering, water thoroughly before application. Do not water for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected with 48 hours. Do not exceed recommended rates because damage to turf may occur. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Do not apply when temperature exceeds 90°F. Shake well before using.

checked on 12/1/78 Label

WEEDS CONTROLLED BY TRIAMINE SPOT WEED KILLER

- Alders, Alligatorweed, American lotus, Arrowhead, Artichoke, Aster, Austrian fieldcress, Beggarticks, Bidden, Bindweed, Bitterweed, Bitter winter cress, Black medic, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckhorn, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common sowthistle, Creeping jenny, Cotton, Curly indigo, Dandelion, Dock, Dogbane, Dogfennel, Duckweed, Elderberry, English daisy, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hawkweed, Healall, Heartloaf drymary, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Indigo, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarters, Locoweed, Lupines, Mallow, Marshelder, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Nutgrass, Orange hawkweed, Oxalis, Parrot feather, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison

ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, Sagebrush, St. Johnswort, Sheep sorrel, Shepherdspurse, Sicklepod, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedles, Spatterdock, Speedwell, Spurge, Spurweed, Stinkweed, Stitchwort, Sumac, Sunflower, Sweet clover, Tarweed, Thistle, Toadflax, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia creeper, Water hyacinth, Water lily, Water plantain, Water primrose, Water shield, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Woodsorrel, Wormseed, Yarrow, Yellow rocket, Virginia buttonweed and other broadleaf weeds.

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. Some hard-to-control broadleaf weeds may require a second (or an additional) treatment in 3-4 weeks.

This product contains a foam which marks the spot where the weed killer has been applied.

DIRECTIONS

1. Shake can just before using and occasionally during use.
2. Invert can, hold at arm's length, and point can at weed as you press button. Do not press button when can is in upright position.
3. Stand close to weed (1 to 2 feet) with can aimed at weed and away from face.
4. Push down briefly on button. Spray to cover center of weed, do not over-spray. The spray's foaming action will make it easy for you to see coverage.
5. One application will kill most weeds listed. If certain tough weeds are not dead within 3 weeks, repeat application.

STORAGE AND DISPOSAL

STORAGE: Keep from freezing. To be stored in original container and placed in an area inaccessible to children. Do not store near heat or open flame.

PESTICIDE DISPOSAL: Product remaining in original container should be disposed of by securely wrapping original container in several layers of newspaper and discarding in trash.

CONTAINER DISPOSAL: Do not incinerate or puncture. Replace cap and discard container in trash.

NOTICE: Buyer assumes all risks of use, storage and handling this product not in strict accordance with directions given herewith.

