

Reg # 228-190

PM-231

TELEPHONE (800) 345-3330
(708) 754-3330
FAX (708) 754-0314



C H E M I C A L C O M P A N Y

194

OFFICE: 425 WEST 194th STREET, GLENWOOD, IL 60425-1584
PLANT: 220 EAST 17th STREET, CHICAGO HEIGHTS, IL 60411-3699

February 25, 1993

CERTIFICATION


I, being an authorized representative of Riverdale Chemical Company certify that all containers of Riverdale TriamineTM Jet-Spray Spot Weed Killer, EPA Reg. No. 228-190 produced by June 15, 1994, will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products. I further certify this company by January 1, 1995, will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products.

Sincerely,

RIVERDALE CHEMICAL COMPANY

James K. Champion
James K. Champion
President

JKC:ls



(Front Panel)

RIVERDALE

TRIAMINE™

JET-SPRAY

SPOT WEED KILLER

READY TO USE

CONTROLS DANDELIONS, CLOVER, CHICKWEED AND
MANY OTHER LISTED WEEDS ON LAWNS AND ORNAMENTAL TURF

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*326%
Dimethylamine Salt of 2-(2-Methyl-4-chlorophenoxy)propionic Acid**328%
Dimethylamine Salt of 2-(2,4-Dichlorophenoxy)propionic Acid***324%

INERT INGREDIENTS:	99.022%
	TOTAL	100.000%

By Isomer Specific AOAC Method No. 6.321, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid27%	.023 lbs/Gallon
**2-(2-Methyl-4-chlorophenoxy)propionic Acid27%	.023 lbs/Gallon
***2-(2,4-Dichlorophenoxy)propionic Acid27%	.023 lbs/Gallon

†TRIAMINE IS A TRADEMARK OF RIVERDALE CHEMICAL COMPANY
Patent Pending

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS
AND STATEMENT OF PRACTICAL TREATMENT

NET WEIGHT 1 lb. 7 oz.

EPA REG. NO. 228-190

EPA EST. NO. 36994-IL-1

Manufactured By

RIVERDALE CHEMICAL COMPANY

Glenwood, Illinois 60425-1584

(Left Side Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if swallowed or absorbed through skin. Wear goggles, long pants, long-sleeved shirt, socks, shoes and chemical resistant gloves when handling or applying this product. Wash nondisposable gloves thoroughly with soap and water before removing. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes).

PHYSICAL OR CHEMICAL HAZARDS

Contents under pressure. Exposure to temperatures above 130°F may cause bursting.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

USE PRECAUTIONS

IMPORTANT: Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray exposed roots of desirable plants. Do not spray on Dichondra, Lippia, nor on grasses where desirable Clovers are present. Do not apply to newly seeded grasses until they have been mowed 3 times. Do not cut grass 1 to 2 days before and after application. Do not reseed for 3 to 4 weeks after use. If lawn needs watering, water thoroughly before application. Do not water for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours. Do not exceed recommended rates because damage to turf may occur. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Do not apply when temperature exceeds 90°F. Shake well before using.

WEEDS CONTROLLED BY TRIAMINE™ JET-SPRAY SPOT WEED KILLER

Alders, Artichoke, Aster, Austrian fieldcress, Beggarticks, Bident, Bindweed, Bitterweed, Bitter winter cress, Black medic, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckhorn, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common sowthistle, Creeping jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed,

494

Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hawkweed, Healall, Heartloaf drymary, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Indigo, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarters, Locoweed, Lupines, Mallow, Marshelder, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Nutgrass, Orange hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, Sagebrush, St. Johnswort, Sheep sorrel, Shepherdspurse, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedles, Spatterdock, Speedwell, Spurge, Spurweed, Stinkweed, Stitchwort, Sumac, Sunflower, Sweet clover, Tarweed, Thistle, Toadflax, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Woodsorrel, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds.

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. Some hard-to-control broadleaf weeds may require a second (or an additional) treatment in 3-4 weeks.

This product contains a foam which marks the spot where the weed killer has been applied.

DIRECTIONS

1. Shake can just before using and occasionally during use.
2. Invert can, hold at arm's length, and point can at weed as you press button. Do not press button when can is in upright position.
3. Stand close to weed (1 to 2 feet) with can aimed at weed and away from face.
4. Push down briefly on button. Spray to cover center of weed, do not over-spray. The spray's foaming action will make it easy for you to see coverage.
5. One application will kill most weeds listed. If certain tough weeds are not dead within 3 weeks, repeat application.

STORAGE AND DISPOSAL

STORAGE: Keep from freezing. To be stored in original container and placed in an area inaccessible to children. Do not store near heat or open flame.

PESTICIDE DISPOSAL: Product remaining in original container should be disposed of by securely wrapping original container in several layers of newspaper and discarding in trash.

CONTAINER DISPOSAL: Do not incinerate or puncture. Replace cap and discard container in trash.

NOTICE: Buyer assumes all risks of use, storage and handling this product not in strict accordance with directions given herewith. (EPA 061594)