#### RIVERDALE

## DiBro GRANULAR WEED KILLER Contains Diuron and Bromacil

FEB - 5 1987

228-20

## KEEP OUT OF REACH OF CHILDREN

## CAUTION

SEE REAR PANEL FOR ADDITIONAL PRECAUTIONS

# PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

## CAUTION

May irritate eyes, nose, throat and skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Remove and wash contaminated clothing before re-use.

## STATEMENT OF PRACTICAL TREATMENT

If in Eyes: Flush with plenty of water. Get medical

attention if irritation persists.

If on Skin: Wash with plenty of soap and water.

#### **ENVIRONMENTAL HAZARDS**

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

EPA REG. NO. 228-

NET WEIGHT\_\_LBS.

EPA EST. NO. 228-'[L-1

## MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY CHICAGO HEIGHTS, ILLINOIS 60411-3699

(KEAK PANEL)

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

## NON-CROP WEED CONTROL - INDUSTRIAL USE ONLY

Riverdale DiBro Granular is a non-selective herbicide for controlling a wide range of annual and perennial weeds and grasses. It is recommended only for noncropland areas such as railroad right-of-ways and industrial areas. 2000 to 4000 pound applications of Riverdale DiBro Granular per acre usually results in a non-productive condition of the soil for a period of a year or more. The duration of non-productivity is dependent upon rainfall, soil type and other conditions.

Riverdale DiBro Granular should be applied as furnished with a seed spreader, a fertilizer spreader, a shaker-type applicator, or with any equipment which will distribute the chemical uniformly over the area to be treated. For the control of annual and most perennial weeds and grasses, apply DiBro Granular at the rate of 2000 to 4000 pounds per acre; on smaller plots 0.5 to 1.0 pound per 5 square feet. Repeat spot treatment may be required when deep-rooted perennial weeds are present. To obtain best results, Dibro Granules should be applied to the ground where it will be absorbed by the roots. Application made early in the season have been found to give results superior to applications at a later seasonal date. DiBro Granular may, however, be applied at any time of the year. Maximum effectiveness in arid regions is obtained when application is made just prior to the rainy season. Users should consult state agricultural experimental stations or extension service weed specialist for recommendations as to use in their particular area.

IMPORTANT: (NOT FOR RESIDENTIAL USE) Do not apply on or near valuable woody or herbaceous plants or on areas where their roots may extend, because of possible injury to such plants. Thoroughly clean spreading equipment with a suitable chemical cleaner before using for other purposes (or do not use same spreading equipment for other purposes). Do not use on croplands or any land to be used for subsequent cropping. Keep animals off treated areas:...:



## STORAGE AND DISPOSAL

STORAGE: Always store pesticides in a secured warehouse or storage building. Do not store near seeds, fertilizers, insecticides or fungicides. Do not contaminate water, food, or feed by storage, disposal or cleaning of equipment. PESTICIDE DISPOSAL: Open dumping is prohibited. that cannot be used according to label instructions must be disposed of according to applicable Federal, state or local procedures. Place in a closed, labeled container for proper disposal. CONTAINER DISPOSAL: Fiber Drum: Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of liner and drum in a sanitary landfill or by incineration if allowed by state and local authorities. Fiber Canister: Do not reuse empty canister. Discard empty canister in trash.

NOTICE: Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith.