

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

JUL 28 1994

Mr. Russell F. Sawyer
 Riverdale Chemical Company
 425 West 194th Street
 Glenwood, IL 60425-1584

Dear Mr. Sawyer:

SUBJECT: Riverdale Weed and Feed with MCPA & Mecoprop
 EPA Registration No. 228-203
 Waiver Request - Removal of Reentry Statement
 Your Submission Dated February 17, 1994

The Agency has reviewed your request to waive the reentry statement from the end-use product listed above. The labeling submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable with the following comments:

On the non-commercial label:

1. Delete the commercial uses from the label, i.e., all uses except for home lawns. The other uses are properly identified on the commercial label.

A stamped copy is enclosed for your records. Please submit five (5) final printed copies for the referenced label incorporating the above changes, before releasing the product for shipment.

Sincerely yours,

Joanne I. Miller
 Product Manager (23)
 Fungicide-Herbicide Branch
 Registration Division (7505C)

CONCURRENCES

| | | | | | | | |
|---------|--------------|--|--|--|--|--|--|
| SYMBOL | 7505C | | | | | | |
| SURNAME | M. J. Miller | | | | | | |
| DATE | 7/27/94 | | | | | | |

(Non-Commercial Label)

ACCEPTED
with COMMENTS
in EPA Letter Dated

(Front Panel)

JUL 28 1994

RIVERDALE

Under the Federal Insecticide, Fungicide, and Plant Growth Regulator Act as amended, for the pesticide registered under EPA Reg. No. 228-203

WEED AND FEED WITH MCPA & MECOPROP

FOR USE ON HOME LAWNS, PARKS, GOLF COURSES
AND OTHER RECREATIONAL TURF AREAS

THIS PRODUCT IS A PREMIUM GRADE TURF FOOD AND HERBICIDE
COMBINATION WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF
WEEDS WHILE PROVIDING BOTH FAST AND LONG FEEDING ACTION

Weeds contacted by this product gradually die and disappear from the lawn.

TREATS (OR COVERS) 5,000 SQ. FEET (50' x 100')

ACTIVE INGREDIENTS:

| | |
|--|--------|
| Isocetyl (2-ethylhexyl) Ester of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid* | 1.0% |
| Isocetyl (2-ethylhexyl) Ester of 2-Methyl-4-Chlorophenoxyacetic Acid** | 1.0% |
| INERT INGREDIENTS: | 98.0% |
| TOTAL | 100.0% |

ISOMER SPECIFIC AOAC METHOD, EQUIVALENT TO:

| | |
|---|-------|
| *2-(2-Methyl-4-Chlorophenoxy)propionic Acid | 0.62% |
| **2-Methyl-4-Chlorophenoxyacetic Acid | 0.60% |

GUARANTEED ANALYSIS:

| | |
|--|----|
| Total Nitrogen (N) | X% |
| X% Ammoniacal Nitrogen | |
| X% Urea Nitrogen | |
| Available Phosphoric Acid (P ₂ O ₅) | X% |
| Soluble Potash (K ₂ O) | X% |

Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated Urea, Urea, Triple Super Phosphate, and Muriate of Potash

Chlorine (Cl) Maximum X%

The urea nitrogen in this product has been coated to provide X% coated slow release urea nitrogen.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
AND STATEMENT OF PRACTICAL TREATMENT

NET WEIGHT LBS.

EPA REG. NO. 228-203

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584

(Back Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Avoid inhalation of dust and contact with eyes, skin, or clothing. Wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Get medical attention.
IF IN EYES: Flush eyes with water for 15 minutes. Get medical attention.
IF ON SKIN: Wash skin with soap and water.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect non-target plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

RIVERDALE WEED AND FEED WITH MCPA & MECOPROP is a remarkably effective lawn fertilizer which combines both fast and long lasting feeding action plus effective weed control. It is for use on Lawns, Ornamental Turf (not pastures), Parks, and Golf Fairways, Aprons, Tees and Roughs.

This product may be used on Bahia, Bermuda, Blue Grass, Fescue, Rye and Zoysia Lawns. USE CAUTION: Some bent varieties such as Arlington, Congressional and Seaside may be injured and some injury may occur to St. Augustine. DO NOT USE on Dichondra. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" on flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product was designed to control such broadleaf weeds as: Beggarticks, Bindweed, Black medic, Burcucumber, Burdock, Buttercup, Canadian thistle, Carpetweed, Catsear, Chervil, Chickweed, Cinquefoil, Clover, Cocklebur, Croton, Daisy, Dandelion, Dock, Dragonhead mint, Fat hen, Fennel, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Halbert-leaved saltbush, Hempnettle, Henbit, Hoary cress, Honeysuckle, Jimsonweed, Kochia, Knotweed, Lambsquarters, Marsh elder, Mexican poppy, Morningglory, Mustard, Nettle, Pennycress, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Prickly lettuce, Puncturevine, Purslane, Ragweed, Red root, Redstem, Shepherdspurse, Sicklepod, Sneezeweed, Sow thistle, Spanish needle, Stinkweed, Stitchwort, Sunflower, Thistle, Thornapple, Vetch, Whitebrush, White top, Wild carrot, Wild gooseberry, Wild jute, Wild marigold, Wild petunia, Wild radish, Wild sage, Wild garlic, Wintercress, Witchweed, Yellow charlock, Yellow daisy, Yellow rocket.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Apply when grass is moist (after a rain, sprinkling or dew). For most effective broadleaf weed killing action, avoid unnecessary disturbances, including watering, of treated areas for 48 hours after application. Watering will wash off weed kill material. Do not apply if rainfall is expected within 48 hours following treatment. Avoid contact with desirable shrubs, plants, vegetables, or flowers. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. Do not cut grass 1 to 2 days before and after application.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: The contents of this bag should be used on 5000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this 20 pound package at a suggested rate of 4 pounds (3.2 lb. for 16 lb. bag or 3.6 lb. for 18 lb. bag) per 1000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 sq. ft. and adjust to apply at the rate of 1 lb. (.8 lb. for 16 lb. bag or .9 lb. for 18 lb. bag) per 250 sq. ft. Clean spreader thoroughly after use.

SPREADER SETTINGS
For One Pass Application*

| <u>MAKE</u> | <u>SETTING</u> | <u>MAKE</u> | <u>SETTING</u> |
|-------------|----------------|------------------|----------------|
| Cyclone | 5-1/4 to 5-3/4 | Montgomery Wards | |
| Scotts 35 | 5-1/2 | 1091, 1093, 1095 | 6-3/4 |
| Scotts C-1 | J - L | Montgomery Wards | |
| Central | 7 - 8 | 1080, 1081 | 5 |
| K-Mart | 7 - 8 | Precision | 6 |

Sweep all fertilizer materials from concrete surfaces immediately after application. Do not wash material from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near other feed or foodstuffs. To prevent cross-contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides or seeds.

PESTICIDE DISPOSAL: To dispose of bag which contains unused product, securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discard in trash.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

(PR938 042194/RV 122193)

The following note will not be part of the actual label copy.

*NOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers.