## (Front Panel) .

## RIVERDALE®

TRIAMINE® II LIQUID WEED AND FEED

READY-TO-USE

FOR USE WITH A HOSE-END SPRAYER ON GRI AND ORNAMENTAL TURF NO MIXING

AUG 8 2003

Just Connect Hose, Turn On Water, and Sprayder the Federal Inspecial

Fungicide, and Rode

TREATS (OR COVERS) 5,000 to 7,500 SQ. Free amended, for the REATS (OR COVERS) 5,000 to 7,500 Sq. registered under (Read directions for different grasses registered under 1974 Reg. No. 228-20

## CONTAINS MCPA, MECOPROP, DICHLORPROP AND A X-X-X FERTILIZER

ACTIVE INGREDIENTS:	
Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	ક
Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy) propionic Acid** 3.26	ક્ર
Dimethylamine Salt of 2-(2,4-Dichlorophenoxy) propionic Acid***	윰
INERT INGREDIENTS:	ક
TOTAL $100.\overline{00}$	ક
Isomer Specific AOAC Method, Equivalent to:  *2-Methyl-4-Chlorophenoxyacetic Acid	•
GUARANTEED ANALYSIS: Total Nitrogen (N)	B

Fertilizer guaranteed by (Name of Fertilizer registrant).

Riverdale and Triamine are Registered Trademarks of Riverdale - A Nufarm Company

#### KEEP OUT OF REACH OF CHILDREN

## CAUTION

## SEE BACK AND SIDE PANELS FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

NET CONTENTS 1 QT. (32 FL. OZ.)

NET WEIGHT 2.3 LBS.

EPA Reg. No. 228-204

EPA EST. No. 228-IL-1

MANUFACTURED BY NUFARM AMERICAS, INC., BURR RIDGE, ILLINOIS, 60537-0866

(Back and Side Panels)

#### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Harmful if absorbed through skin. Wash thoroughly with soap and water after handling

#### FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

IF IN EYES:

Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

IF SWALLOWED:

Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

IF ON SKIN OR CLOTHING: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

### ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate nor pour the washwaters on the ground; spray or drain over a large area away from wells, other water sources and desirable plants. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes. If spills occur, collect the material and dispose of by following disposal instructions on this label.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Spray anytime during the growing season when weeds are actively growing (except as noted for S-Southern or Warm Season Grasses).

5,000 Sq. Ft. Coverage for (N-Northern or Cool Season Grasses) such as Blue Grass, Fescue, and Rye grasses.

7,500 Sq. FT. Coverage for (S-Southern or Warm Season Grasses) Bahia, common Bermuda, St. Augustine, Centipede, Zoysia and Bent grasses. Some temporary injury may occur with these grasses. In Spring, do not apply until grass has reached full green-up. In Fall, exercise care when grass growth is going into dormancy. When applying to Bent grass, height of grass should be 1-1/2" or more. Note: Some hybrid Bermuda grasses may be sensitive to this product. Cortact your local Extension Service weed control specialist.

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing  $\frac{*(No. \text{ of } lbs.)}{*(No. \text{ of } lbs.)}$  pounds of nitrogen per 1,000 square feet (optional  $P_2O_5$  and  $K_2O$  levels) and Triamine<sup>TM</sup> to control listed weeds. Please contact your local extension office for questions regarding a regular fertilizer program.

\*Based on Inert fertilizer claimed.

## USE PRECAUTIONS

IMPORTANT: Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray exposed roots of desirable plants. Do not spray on Dichondra, Lippia, nor on grasses where desirable Clovers are present. Do not apply to newly seeded grasses until they have been mowed 3 times. Do not cut grass 1 to 2 days before and after application. Do not reseed or plant stolons for 3 to 4 weeks after use. If lawn needs watering, water thoroughly before application. Do not water for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours. Avoid fine mists. Use a lawn type sprayer with a coarse spray as wind drift is less likely. Spray entire lawn, wetting weeds and lawn grasses. Do not exceed recommended rates because damage to turf may occur. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Do not apply when temperature exceeds 85°F. Shake well before using.

## WEEDS CONTROLLED BY TRIAMINE® II LIQUID WEED AND FEED

Alder, Beggartick, Bindweed, Black medic, Blue lettuce, Buckhorn, Burcucumber, Burdock, Buttercup, Canada thistle, Carpetweed, Catsear, Chervil, Chickweed, Chicory, Clover, Cocklebur, Croton, Dandelion, Dock, Dogfennel, Dragonhead mint, Dwarf nettle, English daisy, Fanweed, Fat hen, Field bindweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Halbertleaved salt bush, Hawkweed, Healall, Heartleaf drymary, Hempnettle, Henbit, Hoary cress, Honeysuckle, Jimsonweed, Kochia, Knotweed, Lambsquarters, Mallow, Marshelder, Mexican poppy, Miners lettuce, Moneywort, Morningglory, Mustard, Nutgrass, Oxalis, Pennycress, Pennywort, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Prickly lettuce, Puncture vine, Purslane, Ragweed, Red clover, Red root, Red sorrel, Red stem, Sheep sorrel, Shepherdspurse, Smartweed, Sneezeweed, Sowthistle, Spanishneedles, Speedwell, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Sunflower, Sweet clover, Thornapple, Toadflax, Tree-of-heaven, Velvet leaf, Veronica, Vetch, Violet, Virginia buttonweed, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild gooseberry, Wild jute, Wild lettuce, Wild marigold, Wild petunia, Wild onion, Wild radish, Wild sage, Wild strawberry, Willow, Witchweed, Woodsorrel, Yarrow, Yellow charlock-mustard, Yellow daisy, Yellow rocket, and other broadleaf weeds.

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. Some hard-to-control broadleaf weeds may require a second (or an additional) treatment in 3-4 weeks.

#### (HOSE-END SPRAYER INSTRUCTIONS)

HOW TO APPLY: Use any hose-end sprayer that is designed for use on lawns and makes a minimum of 15 gallons of finished spray. Such sprayers are Ortho's Orthomatic Lawn Sprayer, Ortho's and Black Leaf's Lawn Sprayers (dilution ratio of 60 to 1). The following are adjustable/dial sprayers:

MANUFACTURER	<u>SETTING</u>
Super K-Gro	20
Melnor #94 & #95	11
Melnor #96 & #97	13
Gilmour #362	4
Gilmour #486	12

SPECIAL NOTE: Some adjustable sprayers do not hold 32 oz. The above settings are based on using 32 oz. per 5,000 or 7,500 sq. ft. The entire contents of this bottle must be used to make the proper recommended application rate.

Determine area to be sprayed. Apply at a rate of 32 oz. (1 quart) per 5,000 sq. ft. on Cool Season Grasses or 7,500 sq. ft. for application on S-Southern or Warm Season Grasses. Fill sprayer jar with appropriate amount of this product. Do not dilute this product with water when filling sprayer jar. The water will automatically mix with this product during spraying. Attach sprayer to garden hose while holding sprayer at waist level. Point in direction away from face and body. Walk back and forth at a steady pace. Overlap slightly for uniform coverage.

# (DISPOSABLE SPRAYER AND BOTTLE INSTRUCTIONS)

HOW TO APPLY: Connect spray nozzle to hose. Remove bottle cap and screw bottle into nozzle. While holding sprayer at waist level, point in direction away from face and body. Turn on water at the tap. Place finger over hole on right side of nozzle (or turn on/off

lever to the "ON" position) and this product will mix automatically at the proper ratio. Walk back and forth at steady pace. Overlap slightly for uniform coverage.

## COVERAGE MEASURING GUIDE

(Marks on left for 1 qt. disposable bottle Not to scale.)

1000 Sq. Ft. 1500

2000 Sq. Ft. 3000

3000 Sq. Ft. 4500

4000 Sq. Ft. 6000

5000 Sq. ft. 7500

Cool Season Warm

(N)

Season (S)

# Environmental **Benefits** (Grasscycling

. . .

Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment: Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lown.

- ✓ A 50' x 50' lawn produces enough. oxygen for a family of 4 every day
- ✓ Reduces Glore & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- ✓ Purifies Water & Modifies Temperature ✓ Improves Soil Condition & Controls Erosion.

# STORAGE AND DISPOSAL

**STORAGE:** Keep from freezing. To be stored in original container and placed in a locked area inaccessible to children. Do not store near feed or foodstuffs.

DISPOSAL: If empty - Do not reuse this container. Place in trash or offer for recycling if available. If partly filled - If product cannot be used as directed, or offered to others for their use, call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

NOTICE: Buyer assumes all responsibility for safety and use not in strict accordance with directions and instructions for use. (RV 032103)

The following note will not be part of the actual label copy.

Guaranteed (fortilizer) analysis and weight will vary among various commercial fertilizer manufacturers.