

(FRONT PANEL)

RIVERDALE

WEDESTROY^(R)

TRIAMINETM II

1064
PM 23
228-206

FOR SELECTIVE BROADLEAF WEED CONTROL
IN ORNAMENTAL LAWNS AND TURF GRASSES

CONTAINS MCPA, MECOPROP AND DICHLORPROP

ACCEPTED
JUN 2 1987
EPA Reg. No. 228-206

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*.....	17.15%
Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid**.....	16.94%
Dimethylamine Salt of 2-(2,4-Dichlorophenoxy)propionic Acid***.....	16.69%
INERT INGREDIENTS.....	49.22%
TOTAL.....	100.00%

By Isomer Specific A.O.A.C. Method, Equivalent to:

*2-methyl-4-chlorophenoxyacetic acid.....	14.0%, 1.3 lbs/gal.
**2-(2-methyl-4-chlorophenoxy)propionic acid.....	14.0%, 1.3 lbs/gal.
***2-(2,4-dichlorophenoxy)propionic acid.....	14.0%, 1.3 lbs/gal.

KEEP OUT OF REACH OF CHILDREN
DANGER

See Side Panel For Additional Precautionary Statements

STATEMENT OF PRACTICAL TREATMENT

- IF IN EYES: Immediately flush eyes with plenty of water for 15 minutes and get medical attention at once.
- IF SWALLOWED: Drink one or two glasses of water. Induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention immediately.
- IF ON SKIN: Wash promptly with soap and water and rinse thoroughly.

Weedestroy and Triamine are Trademarks of Riverdale Chemical Co. Patent Applied For
E.P.A. REG. NO. 228- NET CONTENTS GAL. E.P.A. EST. NO. 228-IL-1

MANUFACTURED BY
RIVERDALE CHEMICAL COMPANY
CHICAGO HEIGHTS, ILLINOIS 60411-3699

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER

Corrosive. Causes eye damage. Do not get in eyes, on skin or clothing. Wear goggles or face shield and rubber gloves when handling. If eye exposure occurs, get immediate medical attention. Avoid inhalation or spray mists. Harmful or fatal if swallowed. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not contaminate domestic or irrigation waters. Do not apply when weather conditions favor drift from target areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

GENERAL INFORMATION

WEEDESTROY TRIAMINE II is a stable solution containing a mixture of three herbicides. It is non-flammable. It is for use on Lawns, Ornamental Turf(not pastures) and Golf Fairways, Aprons, Tees and Roughs.

USE PRECAUTIONS

CARE MUST BE TAKEN TO AVOID DRIFT OF SPRAY MISTS TO NON-TARGET susceptible plants such as cotton, vegetables, flowers, ornamentals, shrubs, trees and other desirable plants. Do not spray near these plants. Do not spray on carpet grass, dichondra, nor on grasses where desirable clovers are present. Use a lawn type sprayer with coarse spray as wind drift is less likely. Avoid fine mists. Do not spray roots of desirable plants. Do not exceed specified application dosages for any area. Do not apply to newly seeded grasses until well established. Do not spray when air temperature is greater than 85°F. Reseed no sooner than three to four weeks after application of this product. Clean and rinse spray equipment using soap or detergent and water and rinse thoroughly before reuse for other sprays. Failure to observe the above precautions may result in turf injury.

(LEFT SIDE PANEL - CONT'G)

WEEDS CONTROLLED BY WEEDESTROY TRIAMINE II

Black Medic, Buckhorn, Burdock, Buttercup, Canada Thistle, Carpet Weeds, Chickweed, Chicory, Clover, Cocklebur, Dandelion, Dock, Dog Fennel, English Daisy, Florida Pursley, Frenchweed, Ground Ivy, Hawkweed, Heal-All, Heartloaf Drymary, Henbit, Jimson Weed, Knotweed, Lambs-quarter, Mallow, Morning Glory, Mustard, Oxalis, Pennywort, Pepperweed, Pigweed, Plantain, Poison Ivy, Poison Oak, Purslane, Rag Weed, Red Clover, Red Sorrel, Sheep Sorrel, Smartweed, Speedwell, Spurge, Spur Weed, Stitchwort, Toadflax, Veronica, Wild Aster, Wild Carrot, Wild Garlic, Wild Geranium, Wild Lettuce, Wild Onion, Wild Radish, Wood Sorrel, Yarrow and many other broadleaf weeds.

ORNAMENTAL LAWNS AND TURF
(Cool Season Grasses)

For normal application, use Weedestroy Triamine II at a rate of 3-4 pints per acre in sufficient water for good coverage (40-80 gallons of water per acre or 1-2 gallons of water per 1000 square feet is normal). Use high rate for hard to kill weeds. Spray when air temperature is between 50-85°F. Avoid applying during excessively dry or hot periods unless irrigation is used. Do not irrigate within 24 hours of application. Do not mow one or two days before or after application.

(RIGHT SIDE PANEL)

Controlled Droplet Applicator:
Reduced rates of Weedestroy Triamine II must be used when grass is stressed from heat, drought, etc. Do not use this method on warm season grasses. Add 2 pints Weedestroy Triamine II to the herbicide container then fill with 3 pints of water, keeping container agitated. Spray entire contents over 33,000 square feet (approximately 3/4 acre). Do not overlap between spray patterns.

BENT GRASS TURF

On closely mowed bentgrass, apply Weedestroy Triamine II at a maximum rate of 2/3 ounce in 3 gallons of water per 1000 square feet (2-3 pints per acre) in May or Mid-August through September when weeds are actively growing. Exercise care to avoid overdosing Bentgrass or injury may result. Do not mow two days before or after application. Do not apply during hot or dry periods. Slight turf yellowing should disappear after about 1 week.

ST. AUGUSTINE AND CENTIPEDE GRASSES

Apply Weedestroy Triamine II at a maximum rate of 2-3 pints in up to 150 gallons of water per acre (2/3 - 1 1/3 oz. in 3 gallons of water per 1000 square feet.). Do not spray St. Augustine or Centipede grasses if stressed from heat, drought, etc. Slight turf yellowing should disappear after about 1 week.

STORAGE AND DISPOSAL

STORAGE: Always store pesticides in a secured warehouse or storage building. Do not store near seeds, fertilizers, insecticides or fungicides. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well ventilated areas. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high.

PESTICIDE DISPOSAL: If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Metal Containers - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by State and Local authorities. Plastic containers - Triple rinse(or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

WARRANTY

Riverdale warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant, and soil conditions, mixtures with other chemicals, and other factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.