

NOTICE OF PESTICIDE

(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

Riverdale Team 1.75 Lawn Weeds
and Feed

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Riverdale Chemical Company
220 East 17th Street
Chicago Heights, IL 60411

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an endorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(A) provided that you:

1. Submit/cite all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
2. Add the phrase "EPA Registration No. 228-208" to your label before you release the product for shipment.
3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Robert J. Taylor
Product Manager (25)
Fungicide-Herbicide Branch
Registration Division (TS-767C)

Enclosures

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

Robert J. Taylor

DATE

9/3/87

EPA Form 2270-6 (Rev. 7-76)

PREVIOUS EDITION MAY BE USED UNTIL SUPPLY IS EXHAUSTED

15500; I; Erumela; T-1; KEMCO; 9/1/87; 9/11/87; TG; VO; BM; EK; CB

RIVERDALE
TEAM^(R) 1.15 LAWN WEED AND FEED

A Dual Action Pre-Emergent Crabgrass Killer containing Benefin (Balan^(R)) -Trifluralin and a Premium Turf Fertilizer.

For use on established Bluegrass (perennial), Ryegrass (perennial), Fescue, Bahiagrass, Bermudagrass, Centipedegrass, St. Augustine Grass, and Zoysiagrass for the control of Crabgrass and other listed Annual Weed Grasses on Home Lawns, Golf Courses, Parks and other Ornamental & Recreational Turf.

ACTIVE INGREDIENTS:

N-butyl-N-ethyl-alpha-alpha-alpha, trifluoro-2, 6-dinitro-p-toluidine	0.77%
Alpha, alpha, alpha, trifluoro-2,6-dinitro-N, N-dipropyl-p-toluidine.....	0.38%
INERT INGREDIENTS.....	98.85%
TOTAL.....	100.00%

Team^(R) and Balan^(R) - Registered Trademarks of The Elanco Products Company

GUARANTEED ANALYSIS:

Total Nitrogen.....	25%
1.0% Ammoniacal Nitrogen	
6.8% Sulfur Coated Urea	
17.2% Urea Nitrogen	
Available Potash (P ₂ O ₅).....	3%
Soluble Potash (K ₂ O).....	3%

Primary Plant Foods derived from Urea, Sulfur Coated Urea, Ammonium Phosphate, Triple Super Phosphate, and Muriate of Potash

Chlorine (Cl) Maximum 3%

KEEP OUT OF REACH OF CHILDREN

WARNING

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT



EPA REG. NO. 228-

EPA EST. NO. 228-IL-1

NET WEIGHT POUNDS

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY
Chicago Heights, IL 60411



PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: Causes eye irritation. Do not get in eyes. Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust and contact with skin or clothing. The granular formulation may cause skin sensitization reactions in certain individuals. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Wash contaminated clothing before reuse.

STATEMENT OF PRACTICAL TREATMENT

- IF IN EYES:** Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Call a physician.
- IF SWALLOWED:** Call a physician or Poison Control Center. Drink promptly, a large quantity of milk, egg white, gelatin solution, or, if these are not available, large quantities of water. Avoid Alcohol.
- IF ON SKIN:** Wash with plenty of soap and water. Get medical attention if irritation persists.
- IF INHALED:** If breathing difficulty occurs, move individual to fresh air and get medical attention.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

BEST AVAILABLE COPY

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

WHERE TO APPLY

When applied according to these directions this product can be used safely on the following turfgrass varieties: PERENNIAL BLUEGRASS VARIETIES, PERENNIAL RYEGRASS VARIETIES, BERMUDAGRASS, BAHIA GRASS, CENTIPEDEGRASS, FESCUE VARIETIES, ST. AUGUSTINEGRASS & ZOYSIAGRASS lawns and ornamental turf. Will not harm most nearby ornamental trees and shrubs when used as directed.

WEEDS CONTROLLED

This product contains a combination of Benflin and Trifluralin pre-emergence Herbicide plus a 25-3-3 Lawn Food. It should be applied to established turf for the control of CRABGRASS (Smooth and Hairy), GOOSEGRASS (Silver Crab or Crowfoot), BARNYARDGRASS (Wet, Dry, and Yellow), FOXTAIL (Green and Yellow), and ANNUAL BLUEGRASS (Poa Annua) and provides a complete lawn fertilization.

WHEN TO APPLY

Apply to turf prior to germination of weed seeds since established weeds will not be controlled. Apply only to healthy established turf grass lawns.

IMPORTANT PRECAUTIONS:

DO NOT apply in the spring to Bermuda Grass which has been overseeded with winter grasses (annual rye) since thinning will occur. DO NOT use Dichondra, nor in newly sprigged areas of Bermudagrass, St. Augustinegrass or Centipede Grass, until these turfs are well established, nor to Bentgrass or Putting Greens. DO NOT apply in spring to Turfgrass planted the previous fall. This product may injure established Annual Bluegrass (Poa Annua) & Fine-Leaved Fescue lawns when used at higher rates than 160 lbs. per acre. (2.0 lb. A.I. per Acre). Do not overseed.

WHEN AND HOW TO TIME YOUR APPLICATION

Successful pre-emergence control of the annual grass weeds on this label requires proper timing of application. This product must be applied one to two weeks prior to the germination of annual grass weeds. All summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grass/weeds listed on this label, weed control results may be erratic or poor.

WATER THE LAWN AFTER APPLICATION: It is not necessary but optimum weed control performance will be obtained if treated areas are irrigated with 1/4 to 1/2 inch water soon after application. Any cultural practices that disturb the soil such as verticuting, should be done prior to application of this product.

"SPRING GERMINATING" SUMMER ANNUAL GRASSES: CRABGRASS, FOXTAIL, BARNYARDGRASS:

- A. Northern Cool Season Grasses - For pre-emergence control, apply 130 to 174 lbs. per acre, (equal to 1.5 to 2.00 lbs. Act Ingrid/acre) applied in the spring one to two weeks prior to the onset of conditions favorable for annual woody grass seed germination. 50 lbs. will "Wood and Feed" 12,500 to 16,250 sq. ft.
- B. Southern warm season turfgrasses - For pre-emergence control apply 174 to 261 lbs. per acre (equal to 2 to 3 lbs. Active Ingredient per acre), in the late winter or early spring one to two weeks prior to the onset of conditions favorable for annual woody grass seed germination. 50 lbs. will "Wood and Feed" 8,325 to 12,500 sq. ft. Use for 8,325 sq. ft. where turf is completely overrun by weeds. Use for 12,500 sq. ft. when weeds are sparse and the turf is in a Yearly Crabgrass Control Program.

"LATE SPRING, EARLY SUMMER GERMINATING" ANNUAL GRASSES: GOOSEGRASS (SILVER CRAB)

- Although classified as a summer annual grass, goosegrass normally germinates 4 to 6 weeks later than the crabgrass, foxtails and other summer annual grass species. Applications of this product should therefore, be timed accordingly.
- A. Northern cool season turfgrasses - For pre-emergence control apply 174 lbs. per acre (equal to 2 lbs. Active Ingredient per acre) of this product in late spring or early summer one to two weeks prior to the expected germination period of goosegrass. 50 lbs. will "Wood and Feed" 12,500 sq. ft.
- B. Southern warm season turfgrasses - For pre-emergence control, apply 174 to 261 lbs. per acre (equal to 2 to 3 lbs. Active Ingredient per acre) of this product in late spring or early summer one to two weeks, prior to the expected germination of goosegrass. 50 lbs. will "Wood and Feed" 8,325 to 12,500 sq. ft. Use for 8,325 sq. ft. where turf is completely overrun by weeds. Use for 12,500 sq. ft. when weeds are sparse and the turf is in a Yearly Lawn Maintenance Program.

