

pm 23

228-216

178

JAN 16 1990

Mr. Russell P. Sawyer
Riverdale Chemical Company
425 West 194th Street
Glenwood, IL 60425-1584.

Dear Mr. Sawyer:

Subject: Riverdale Triamine™ Premium 6666 Liquid Weed and Feed
EPA Registration No. 228-216
Your Application To Amend Registration Dated
September 19, 1989

The additional precautions "May cause eye irritation and Avoid contact with skin, eyes or clothing" are acceptable for the subject product labeling and are conditional acceptable under the Federal Insecticide, Fungicide, and Rodenticide Act as amended. The conditions of this acceptance of labeling amendments are as follows:

1. Correct the spelling of the word "boosts" under the General Information Section.
2. The labeling may only be used with an O. M. Scott supplemental registration, as its warranty statements are designed only for such use.
3. That you will submit/cite all data required for registration/reregistration of this product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.
4. Submit five (5) copies of the final printed labeling before releasing the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes

BEST AVAILABLE COPY

-2-

acceptance of these conditions. A stamped copy of the label is enclosed for your records.

Sincerely yours,

Joanne I. Miller
Acting Product Manager (23)
Fungicide/Herbicide Branch
Registration Division (H-7505G)

Enclosure

E.Wilson:Diskette 007:1-12-90

BEST AVAILABLE COPY

3 of 8

ACCEPTED
STATEMENTS
EPA Letter Dated:

JAN 16 1990

(Front Panel)

RIVERDALE

TRIAMINE™ PREMIUM 6666

LIQUID WEED AND FEED

Product of Triamine Insecticide,
Fertilizer and Herbicide Act
of 1975, for the pesticide
registered under EPA Reg. No.

228-216

LIQUID LAWN CARE

READY-TO-USE

FOR USE WITH A HOSE-END SPRAYER ON GRASS AND ORNAMENTAL TURF (TURF GRASS LAWNS)

NO MIXING

TREATS (COVERS) 4,250 TO 10,000 SQ. FT. IN A MATTER OF MINUTES

Gives your lawn a quick boost of important nutrients while controlling
broadleaf weeds. Attach to a garden hose-end sprayer and spray.

Read directions for different grasses.
A Quick and Easy Way to a Quality Lawn.

CONTAINS 2,4-D, MECOPROP, DICHLORPROP AND A 15-0-0 FERTILIZER

ACTIVE INGREDIENTS:+

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	2.29%
Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid**	2.30%
Dimethylamine Salt of 2-(2,4-Dichlorophenoxy)propionic Acid***	2.26%

INERT INGREDIENTS:	93.15%
TOTAL	100.00%

Isomer Specific by AOAC Method

*2,4-Dichlorophenoxyacetic Acid equivalent	1.9%	.18 lbs./gal.
**2-(2-Methyl-4-Chlorophenoxy)propionic Acid equivalent	1.9%	.18 lbs./gal.
***2-(2,4-Dichlorophenoxy)propionic Acid equivalent	1.9%	.18 lbs./gal.

GUARANTEED ANALYSIS:°

Total Nitrogen (N)	15%
15% Urea Nitrogen	

+TRIAMINE IS A TRADEMARK OF RIVERDALE CHEMICAL COMPANY

Patent Pending.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE AND BACK PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET CONTENTS: 1/2 GAL. (64 Fl. Oz.)

NET WEIGHT:° 4.6 LBS.

EPA REG. NO. 228-216

EPA EST. NO. 228-1L-1

BEST AVAILABLE COPY

4 of 8

Manufactured By

RIVERDALE CHEMICAL COMPANY

Glenwood, Illinois 60425-1584

(Back Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin. May cause eye irritation. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling.

- IF SWALLOWED: Get medical attention.
- IF IN EYES: Immediately flush eyes with plenty of water.
- IF ON SKIN: Wash with plenty of soap and water. If irritation develops, get medical attention.

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift from target areas.

GENERAL INFORMATION

Designed to control broadleaf weeds and give your lawn a quick boost of important nutrients. Controls dicot weeds during their active growth period. Guaranteed not to burn your lawn when used as directed. High analysis fertilizer featuring a patented nitrogen formula.

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed this product will supplement your regular fertilizer program by providing *(no. of lbs.) pounds of nitrogen per 1,000 square feet (optional P₂O₅ and K₂O levels) and Triamine™ to control listed weeds. Please consult your local extension office for questions regarding a regular fertilizer program.

*Based on Inert fertilizer claimed.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL BEFORE USING. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

BEST AVAILABLE COPY

(Back Panel cont.)

- 4,250 to 6,666 Sq. Ft. Coverage for (N-Northern or Cool Season grasses)
Bahia, Bermuda, Bluegrass, Fescue, Rye and Zoysia grasses.
- 8,500 to 10,000 Sq. Ft. Coverage for (S-Southern or warm season grasses)
(spring and fall application only) St. Augustine, Centipede
and Bent grasses. Some temporary injury may occur with these
grasses.

Do not exceed recommended rate because damage to lawn may occur. Do not apply to
Dichondra, Lippia and Clover lawns. Do not use this product for controlling weeds in
vegetable gardens and flower beds or around shrubs and ornamental plants.

WEEDS CONTROLLED

Black medic	Jimsonweed	Red sorrel
Buckhorn	Knotweed	Sheep sorrel
Buttercup	Lambsquarters	Speedwell
Canada thistle	Mallow	Spurge
Carpetweed	Morningglory	Spurweed
Chickweed	Mustard	Stitchwort
Chicory	Oxalis	Toadflax
Clover	Pennywort	Wild aster
Cocklebur	Pepperweed	Wild carrot
Dandelion	Pigweed	Wild garlic
Dogfennel	Plantain	Wild geranium
English daisy	Poison ivy	Wild lettuce
Florida pusley	Poison oak	Wild onion
Frenchweed	Purslane	Wild radish
Ground ivy	Ragweed	Wood sorrel
Healall	Red clover	Yarrow, and other broadleaf weeds.
Heartloaf drymary		
Henbit		

WHEN TO USE: Spray anytime during the growing season when weeds are actively growing
(except as noted for St. Augustine, Centipede and Bent grasses). If lawn is extremely
dry, water 1 to 2 days before spraying. Do not apply when temperatures are expected to
exceed 85°F, or if rain is expected to occur within 24 hours following application. On
new lawns wait until grass has reached a height of 2 inches before spraying. Also wait
3 weeks before putting in new grass or stolons.

After applying this product, many broadleaf weeds will begin to show visible signs of
leaf curl within a few days. Some hard-to-control broadleaf weeds may require a second
treatment in 3-4 weeks.

(HOSE-END SPRAYER INSTRUCTIONS)

HOW TO APPLY: Use any 32 oz. hose-end sprayer that is designed for use on lawns and
makes a minimum of 15 gallons of finished spray. Such sprayers are Ortho's Orthomatic
Lawn Sprayer, Ortho's and Black Leaf's Lawn Sprayers (dilution ratio of 60 to 1). The
following are adjustable/dial sprayers:

MANUFACTURER

Supe. K-Gro

SETTING

20

BEST AVAILABLE COPY

(Back Panel cont.)

Melnor #94 & #95	11
Melnor #96 & #97	13
Gilmour #362	4
Gilmour #486	12

Special Note: Some adjustable sprayers do not hold 32 oz. The above settings are based on using 64 oz. per 4,250 to 6,666 (N) or 8,500 to 10,000 (S) Sq. Ft. The entire contents of this bottle must be used to make the proper recommended application rate.

Determine area to be sprayed. Apply at a rate of 64 oz. (1/2 gal.) per 4,250 to 6,666 square feet or 8,500 to 10,000 for spring application on St. Augustine, Centipede or Bent grasses. Shake well and fill sprayer jar with appropriate amount of this product. Do not dilute this product with water when filling sprayer jar. The water will automatically mix with this product during spraying. Attach sprayer to garden hose while holding sprayer at waist level. Point in a direction away from face and body. Walk back and forth at a steady pace. Over-lap slightly for uniform coverage. Do not exceed recommended rate because damage to lawn may occur. Do not water fertilized lawn for at least 24 hours.

(DISPOSABLE SPRAYER AND BOTTLE INSTRUCTIONS)

HOW TO APPLY: Determine area to be sprayed. Apply at a rate of 64 oz. (1/2 gal.) per 4,250 to 6,666 square feet or 8,500 to 10,000 for spring application on St. Augustine, Centipede or Bent grasses. Shake well before applying. Connect spray nozzle to hose. Remove bottle cap and screw bottle into nozzle. Turn on the water at the tap while holding sprayer at waist level. Point in a direction away from face and body. Place finger over hole on right side of nozzle (or turn on/off lever to the "ON" position) and the Liquid Weed and Feed will mix automatically at the proper ratio. Walk back and forth at a steady pace. Overlap slightly for uniform coverage. Do not exceed recommended rate because damage to lawn may occur. Do not water fertilized lawn for at least 24 hours.

Marks on left side panel of bottle indicate square feet covered.

(OMS SPRAYER INSTRUCTIONS)

Follow these steps to maintain an attractive, quality lawn.

- Shake well before applying.
- Be sure that the siphon tube is tightly connected to the adapter to assure proper siphoning.
- Connect spray nozzle to your garden hose.
- Hold bottle by the neck, remove child-resistant cap.
- Attach bottle to nozzle.
- Avoid squeezing the bottle as contents may spray through the nozzle opening.
- Adjust on/off switch to the "OFF" position, then turn on water at the faucet.
- Beginning at a point furthest from your faucet, turn the on/off switch to the "ON" position.
- Holding the bottle in front of you at waist height, use a moderate, side-to-side swinging motion to apply the product. Use at the rate of 1.3 fl. oz./100 sq. ft. Each pass should cover approximately an 8 to 10 foot swath.
- Walk at a steady pace and overlap treated areas by at least one foot to insure proper coverage.
- Replace child-resistant cap for storage of any unused product.

BEST AVAILABLE COPY

768

(Back Panel cont.)

ILLUSTRATION ILLUSTRATION ILLUSTRATION ILLUSTRATION

- Use whenever broadleaf weeds such as dandelion, clover, plantain and chickweed are actively growing. Apply beginning in spring, after your second mowing, through the fall. Some hard-to-control broadleaf weeds may require a second treatment in 3-4 weeks.
- Do not water your lawn until 24 hours have passed since treatment. Also, this product should not be applied if rain is expected within 24 hours.
- Regular lawn watering is necessary to maintain a quality lawn. During periods of drought or hot, dry weather, do not use this product until the rains occur or the lawn is watered.

PRECAUTIONS

- Apply on a calm day to avoid spray drift to non-target plants.
- Wait until you mow three or four times before you use this product on new grass.
- Avoid applications near shrubs, flowers, fruit or vegetable plants as plant injury may occur.
- Delay planting grass seed, sodding or sprigging until after a heavy rain or watering and not sooner than three weeks after treatment.
- Not for use on creeping bentgrass, St. Augustinegrass, carpetgrass, dichondra, lippia or similar ground covers. (NOTE: OMS will not be using directions for "Southern" grasses.)

IF YOU HAVE ANY QUESTIONS

This product is designed to help you obtain a better lawn in a simple, easy manner. Should you have any questions about this product or about proper lawn care in general, you can get answers from one of Scotts lawn counselors. Simply dial this toll free number: 1-800-543-TURF (in Ohio 1-800-762-4010)

SCOTTS POLICY
No Quibble Guarantee

If for any reason you are not satisfied with results after using any Scotts product, you are entitled to get your money back. Simply U.P.C. CODE send us evidence of purchase and we will mail you a refund check promptly.

(Right Side Panel)

STORAGE: Keep from freezing. (For 25-2-3 fertilizer - Do Not Store above 95°F) To be stored in original container (when applicable: with childproof cap) and placed in a locked area inaccessible to children. Do not store near feed or foodstuff.

PESTICIDE DISPOSAL: Product remaining in original container should be disposed of by securely wrapping original container in several layers of newspaper and discarding in trash.

CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly before discarding in trash.

NOTICE

Seller warrants that this product conforms to the ingredient statement of the label. Since the conditions of use, such as weather, compatibility with other chemicals, and

0 of 8

(Right Side Panel cont.)

conditions of application equipment will vary, seller makes no claims other than those stated on this label.

The following not² will not be part of the actual label copy.

*Note: Guaranteed (fertilizer) analysis and weight will vary among various commercial fertilizer manufacturers.

(Left Side Panel)

(Scotts)

2000 Sq. Ft.

1666 Sq. Ft.

1063 Sq. Ft.

4000 Sq. Ft.

3333 Sq. Ft.

2125 Sq. Ft.

6000 Sq. Ft.

5000 Sq. Ft.

3188 Sq. Ft.

8000 Sq. Ft.

6666 Sq. Ft.

4250 Sq. Ft.

(S)

(N)

(Not to Scale)

BEST AVAILABLE COPY