



TELEPHONE (800) 345-3330  
(708) 754-3330  
FAX (708) 754-0314

C H E M I C A L C O M P A N Y

OFFICE: 425 WEST 194th STREET, GLENWOOD, IL 60425-1584  
PLANT: 220 EAST 17th STREET, CHICAGO HEIGHTS, IL 60411-3699

June 9, 1993

Document Processing Desk (Notification)  
Registration Division - H7505C  
U. S. Environmental Protection Agency  
401 M Street S.W.  
Washington, DC 20460

Dear Sir/Madam:

Subject: PR Notice 93-3 dated March 9, 1993  
Riverdale Triamine™ II Lawn Weed Killer  
EPA Reg. No. 228-217  
OPP Identifier No. 196232

Please find enclosed the necessary paperwork to notify the Agency that we are revising our label in accordance with the above subject PR Notice. Be advised that this product is part of our "John Doe" private label program. As such, we do not market this product under the Riverdale name. **Thus, there will not be any final printed label.** If there is anything else we should do at this time, please contact us immediately.

Sincerely yours,

RIVERDALE CHEMICAL COMPANY

By Russell F. Sawyer  
Regulatory Affairs Manager

RFS:ls

Enc.

**CPDA** Chemical Producers  
and Distributors  
Association



(A)		United States Environmental Protection Agency Office of Pesticide Programs (H7505C) Washington, DC 20460	<input type="checkbox"/> Registration <input type="checkbox"/> Amendment <input checked="" type="checkbox"/> Other	OPP Identifier Number  <span style="font-size: 1.2em;">196232</span>
		<b>Application for Pesticide:</b>		

**Section I**

1. Company/Product Number 228-217	2. EPA Product Manager Joanne I. Miller	3. Proposed Classification  <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) <span style="float: right;">TM</span> Riverdale Triamine II Lawn Weed Killer	PM#  23	
5. Name and Address of Applicant (Include ZIP Code)  RIVERDALE CHEMICAL COMPANY 425 West 194th Street Glenwood, IL 60425-1584  <input type="checkbox"/> Check if this is a new address		6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to:  EPA Reg. No. _____  Product Name _____

**Section II**

<input type="checkbox"/> Amendment - Explain below  <input type="checkbox"/> Resubmission in response to Agency letter dated _____  <input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____  <input type="checkbox"/> "Me Too" Application.  <input type="checkbox"/> Other - explain below.
---	---

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

To comply with PR Notice 93-3

BEST AVAILABLE COPY

NOTIFICATION  
 LABEL NOT REVERSED  
 PER PR NOTICE 93-3  
 DATE: 9/26/93

**Section III**

<b>1. Material This Product Will Be Packaged In:</b>					
Child-Resistant Packaging  <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging  <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging  <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container  <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
* Certification must be submitted.		If "Yes," Unit Package wgt. _____ No. per container _____	If "Yes," Package wgt. _____ No. per container _____		
3. Location of Net Contents Information  <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) of Retail Container		5. Location of Label Directions  <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed To Product  <input type="checkbox"/> Lithograph <input type="checkbox"/> Other ( _____ ) <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled					

**Section IV**

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)		
Name  Russell F. Sawyer	Title  Regulatory Affairs Manager	Telephone No. (Include Area Code)  708/754-3330
Certification  I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment, both under applicable law.		6. Date Application Received  (Stamp)
2. Signature  	3. Title  Regulatory Affairs Manager	
4. Typed Name  Russell F. Sawyer	5. Date  June 9, 1993	

RIVERDALE TRIAMINE™ II

LAWN WEED KILLER

FOR SELECTIVE BROADLEAF WEED CONTROL IN ORNAMENTAL LAWNS AND TURF GRASSES  
And Other Listed Sites

ONE QUART TREATS (COVERS) UP TO 20,000 SQ. FT.  
(See directions for different grasses.)

ACTIVE INGREDIENTS:†

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid* .....	4.78%
Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid** .....	4.72%
Dimethylamine Salt of 2-(2,4-Dichlorophenoxy)propionic Acid*** .....	4.65%

INERT INGREDIENTS: .....	85.85%
TOTAL .....	100.00%

Isomer Specific AOAC Method No. 6.321, Equivalent to:

*2-Methyl-4-Chlorophenoxyacetic Acid .....	3.9%	0.33 lbs./gal.
**2-(2-Methyl-4-Chlorophenoxy)propionic Acid .....	3.9%	0.33 lbs./gal.
***2-(2,4-Dichlorophenoxy)propionic Acid .....	3.9%	0.33 lbs./gal.

†TRIAMINE - TRADEMARK OF RIVERDALE CHEMICAL COMPANY

Patent Pending

KEEP OUT OF REACH OF CHILDREN

WARNING

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

STORAGE AND DISPOSAL

STORAGE: To be stored in original container and placed in an area inaccessible to children.

PRODUCT DISPOSAL: Partially filled container may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Rinse thoroughly before discarding in trash. Do not reuse empty container.

NET CONTENTS GAL.


EPA REG. NO. 228-217

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584



Revised A/O 3/9/93

PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear face shield or goggles, long pants, long-sleeved shirt, shoes and chemical resistant gloves when mixing, loading or applying this product. Wash nondisposable gloves thoroughly with soap and water before removing. Harmful if swallowed, inhaled, or absorbed through skin. Wash thoroughly after handling. Remove contaminated clothing and wash before reuse. Do not allow persons and pets on treated areas until sprays have dried.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Immediately flush eyes with fresh water for 15 minutes and get medical attention immediately.

IF ON SKIN: Wash skin with plenty of soap and water. Get medical attention if irritation persists.

IF SWALLOWED: Drink one or two glasses of water. Induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect non-target plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate nor pour the washwaters on the ground; spray or drain over a large area away from wells, other water sources and desirable plants. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes. If spills occur, collect the material and dispose of by following disposal instructions on this label.

Most cases of groundwater contamination involving phenoxy herbicides such as MCPA, MCPP and 2,4-DP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

GENERAL INFORMATION

TRIAMINE II LAWN WEED KILLER is a stable solution containing a mixture of three herbicides. It is non-flammable. It is for use on Lawns and Turf, also along Highways, Rights-of-Way, Drainage Ditchbanks and other similar non-crop areas contained within a residential locale.

## USE PRECAUTIONS

**IMPORTANT:** Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray exposed roots of desirable plants. Do not spray on Dichondra, Lippia, nor on grasses where desirable Clovers are present. Do not apply to newly seeded grasses until they have been mowed 3 times. Do not cut grass 1 to 2 days before and after application. Do not reseed for 3 to 4 weeks after use. If lawn needs watering, water thoroughly before application. Do not water for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours. Avoid fine mists. Use a lawn type sprayer with a coarse spray as wind drift is less likely. Spray entire lawn, wetting weeds and lawn grasses. Do not exceed recommended rates because damage to turf may occur. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Do not apply when temperature exceeds 90°F. Make new dilution for each use. Shake well before using.

## WEEDS CONTROLLED BY TRIAMINE™ II LAWN WEED KILLER

Alders, Beggarticks, Bindweed, Black medic, Blue lettuce, Buckhorn, Burcucumber, Burdock, Buttercup, Canada thistle, Carpetweed, Catsear, Chervil, Chickweed, Chicory, Clover, Cocklebur, Croton, Dandelion, Dock, Dogfennel, Dragonhead mint, Dwarf nettle, English daisy, Fanweed, Fat hen, Field bindweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Halbertleaved salt bush, Hawkweed, Healall, Heartloaf drymary, Hempnettle, Henbit, Hoary cress, Honeysuckle, Jimsonweed, Kochia, Knotweed, Lambsquarters, Mallow, Marshelder, Mexican poppy, Minerslettuce, Moneywort, Morningglory, Mustard, Nutgrass, Oxalis, Pennycress, Pennywort, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Prickly lettuce, Puncture vine, Purslane, Ragweed, Red clover, Red root, Red sorrel, Red stem, Sheep sorrel, Shepherdspurge, Smartweed, Sneezeweed, Sowthistle, Spanishneedles, Speedwell, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Sunflower, Sweet clover, Thornapple, Toadflax, Tree-of-heaven, Velvet leaf, Veronica, Vetch, Violet, Virginia buttonweed, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild gooseberry, Wild jute, Wild lettuce, Wild marigold, Wild petunia, Wild onion, Wild radish, Wild sage, Wild strawberry, Willow, Witchweed, Woodsorrel, Yarrow, Yellow charlock-mustard, Yellow daisy, Yellow rocket, and other broadleaf weeds.

## ORNAMENTAL LAWNS AND TURF

**Cool Season Grasses** - such as bluegrass, fescue, and rye: To kill dandelion, wild onion, pennywort, plantain, healall and similar weeds: use 8 to 10-1/2 ounces in 6 gallons of water and apply with pump or hose-end sprayer or sprinkling can to 2,500 square feet of lawn. For smaller areas - use 2 to 3 teaspoons in 2 quarts of water applied to 100 square feet. For larger areas - use 1 to 1-1/2 gallons in 50 gallons of water to 1 acre.

To kill undesirable weeds such as thistle, bindweed, dock and ragweed, use 8 to 10-1/2 ounces in 6 gallons of water (or 2 to 3 teaspoonfuls in 2 quarts of water), and apply directly to plants, wetting them thoroughly. Repeat as necessary. For larger areas use 1 gallon in 50 gallons of water.

To kill undesirable plants such as poison ivy and poison oak, use 8 to 10-1/2 teaspoonfuls in 2 quarts of water or 2 gallons in 12 1/2 gallons of water and branch plants when they are fully leaved. Repeat in about 4 weeks if necessary. Certain hard-to-kill weeds such as poison ivy, poison oak and English daisy may require a fall

**BEST AVAILABLE COPY**

6/10

application.

**Warm Season Grasses** - St. Augustine, Bahia, common Bermuda, Centipede, Zoysia, also for BENT: Use a one-half rate which is 4 to 5 ounces in 6 gallons of water to treat 2500 sq. ft. (2 to 3 teaspoonfuls in 4 quarts of water applied to 200 square feet). Make two applications 10 days apart. Do not spray grasses if stressed from heat, drought, etc. Slight turf yellowing should disappear after about 1 week. Note: Some hybrid Bermuda grasses may be sensitive to this product. Contact your local extension service weed control specialist.

**Roadsides** (including aprons and guard rails) rights-of-way, drainage ditchbanks and other similar non-crop areas: For control of broadleaf weeds, mix at a rate of 1 to 3 1/2 pints of Triamine II Lawn Weed Killer per 5 to 30 gallons of water. This mixture will cover one acre. Thoroughly saturate all weeds with spray mixture. Apply when plants come into full leaf (Spring) to the time plants begin to go dormant. Best results are obtained when weeds are young and actively growing. Do not cut weeds until herbicide has translocated throughout the plant causing root death. For small broadleaf weeds use the lower rate. Heavy, dense stands require the higher rate with high water volume.

**For Spot Treatment:** Put 3 teaspoons Triamine II Lawn Weed Killer into an empty quart container with trigger sprayer. Then add 30 oz. tap water. Adjust spray nozzle to give coarse spray. Aim at center of weed and spray to wet. Applicator Cap - Hold applicator end about 6 inches from surface to be treated and thoroughly wet. To obtain wider coverage raise container. One application should be sufficient. Effects begin to show after a few days and weeds gradually die. A repeat application may be required in 3-4 weeks for hard-to-control weeds.

**DISPOSABLE SPRAYER AND BOTTLE INSTRUCTIONS**

Determine area to be sprayed. Apply at a rate of 32 oz. (1 quart) per 10,000 sq. ft. for Cool Season grasses or 20,000 sq. ft. for Warm Season grasses. Connect spray nozzle to hose. Remove bottle cap and screw bottle into nozzle. While holding sprayer at waist level, point in direction away from face and body. Turn on the water at the tap. Place finger over hole on right side of nozzle (or turn on/off lever to the "ON" position) and the Lawn Weed Killer will mix automatically at the proper ratio. Walk back and forth at steady pace.

2000 Sq. Ft. 4000

(Marks on left for 1 qt. disposable bottle. Not to scale.)

4000 Sq. Ft. 8000

6000 Sq. Ft. 12000

8000 Sq. Ft. 16000

10000 Sq. Ft.	20000
Good Season	Warm Season

**NOTICE**

Buyer assumes risks of use, storage and handling of this product not in strict accordance with directions give herewith. (PR933 042194)

**BEST AVAILABLE COPY**

RIVERDALE TRIAMINE™ II

LAWN WEED KILLER

FOR SELECTIVE BROADLEAF WEED CONTROL IN ORNAMENTAL LAWNS AND TURF GRASSES  
And Other Listed Sites

ONE QUART TREATS (COVERS) UP TO 20,000 SQ. FT.  
(See directions for different grasses.)

ACTIVE INGREDIENTS:†

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid* .....	4.78%
Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid** .....	4.72%
Dimethylamine Salt of 2-(2,4-Dichlorophenoxy)propionic Acid*** .....	4.65%

INERT INGREDIENTS: .....	85.85%
TOTAL .....	100.00%

Isomer Specific AOAC Method, Equivalent to:

*2-Methyl-4-Chlorophenoxyacetic Acid .....	3.9%, 0.33 lbs./gal.
**2-(2-Methyl-4-Chlorophenoxy)propionic Acid .....	3.9%, 0.33 lbs./gal.
***2-(2,4-Dichlorophenoxy)propionic Acid .....	3.9%, 0.33 lbs./gal.

†TRIAMINE - TRADEMARK OF RIVERDALE CHEMICAL COMPANY

Patent Pending

KEEP OUT OF REACH OF CHILDREN

WARNING

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

NET CONTENTS GAL.

EPA REG. NO. 228-217

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY


GLENWOOD, ILLINOIS 60425-1584

PRECAUTIONARY STATEMENT

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Causes substantial but temporary eye injury. Do not get in eyes or on clothing. Wear face shield or goggles, long pants, long-sleeved shirt, shoes and chemical resistant gloves when mixing, loading or applying this product. Wash nondisposable gloves thoroughly with soap and water before removing. Harmful if



Revised A/O 3/9/93

**BEST AVAILABLE COPY**

swallowed, inhaled, or absorbed through skin. Wash thoroughly after handling. Remove contaminated clothing and wash before reuse. Do not re-enter or permit workers to re-enter treated areas until sprays have dried.

#### STATEMENT OF PRACTICAL TREATMENT

**IF IN EYES:** Immediately flush eyes with fresh water for 15 minutes and get medical attention immediately.

**IF ON SKIN:** Wash skin with plenty of soap and water. Get medical attention if irritation persists.

**IF SWALLOWED:** Drink one or two glasses of water. Induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

#### ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect non-target plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the high water mark. Do not contaminate water when disposing of equipment washwaters or rinsate nor pour the washwaters on the ground; spray or drain over a large area away from wells, other water sources and desirable plants. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes. If spills occur, collect the material and dispose of by following disposal instructions on this label.

Most cases of groundwater contamination involving phenoxy herbicides such as MCPA, Mecoprop and Dichlorprop have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

#### GENERAL INFORMATION

TRIAMINE II LAWN WEED KILLER is a stable solution containing a mixture of three herbicides. It is non-flammable. It is for use on Lawns, Turf (not pastures) and Golf Fairways, Aprons, Tees and Roughs, Parks, Cemeteries, along Highways, Rights-of-Way, Drainage Ditchbanks and other similar non-crop areas.

#### USE PRECAUTIONS

**IMPORTANT:** Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray exposed roots of desirable plants. Do not spray on Dichondra, Lippia, nor on grasses where desirable Clovers are present. Do not apply to newly seeded grasses until they have been mowed 3 times. Do not cut grass 1 to 2 days before and after application. Do not reseed for 3 to 4 weeks

BEST AVAILABLE COPY

after use. If lawn needs watering, water thoroughly before application. Do not water for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours. Avoid fine mists. Use a lawn type sprayer with a coarse spray as wind drift is less likely. Spray entire lawn, wetting weeds and lawn grasses. Do not exceed recommended rates because damage to turf may occur. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Do not apply when temperature exceeds 90°F. Make new dilution for each use. Shake well before using.

#### WEEDS CONTROLLED BY TRIAMINE™ II LAWN WEED KILLER

Alders, Beggarticks, Bindweed, Black medic, Blue lettuce, Buckhorn, Burcucumber, Burdock, Buttercup, Canada thistle, Carpetweed, Catsear, Chervil, Chickweed, Chicory, Clover, Cocklebur, Croton, Dandelion, Dock, Dogfennel, Dragonhead mint, Dwarf nettle, English daisy, Fanweed, Fat hen, Field bindweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Halbertleaved salt bush, Hawkweed, Healall, Heartloaf drymary, Hempnettle, Henbit, Hoary cress, Honeysuckle, Jimsonweed, Kochia, Knotweed, Lambsquarters, Mallow, Marshelder, Mexican poppy, Minerslettuce, Moneywort, Morningglory, Mustard, Nutgrass, Oxalis, Pennycress, Pennywort, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Prickly lettuce, Puncture vine, Purslane, Ragweed, Red clover, Red root, Red sorrel, Red stem, Sheep sorrel, Shepherdspurse, Smartweed, Sneezeweed, Sowthistle, Spanishneedles, Speedwell, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Sunflower, Sweet clover, Thornapple, Toadflax, Tree-of-heaven, Velvet leaf, Veronica, Vetch, Violet, Virginia buttonweed, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild gooseberry, Wild jute, Wild lettuce, Wild marigold, Wild petunia, Wild onion, Wild radish, Wild sage, Wild strawberry, Willow, Witchweed, Woodsorrel, Yarrow, Yellow charlock-mustard, Yellow daisy, Yellow rocket, and other broadleaf weeds.

#### ORNAMENTAL LAWNS AND TURF

**Cool Season Grasses** - such as bluegrass, fescue, and rye: To kill dandelion, wild onion, pennywort, plantain, healall and similar weeds: use 8 to 10-1/2 ounces in 6 gallons of water and apply with pump or hose-end sprayer or sprinkling can to 2,500 square feet of lawn. For smaller areas - use 2 to 3 teaspoons in 2 quarts of water applied to 100 square feet. For larger areas - use 1 to 1-1/2 gallons in 50 gallons of water to 1 acre.

To kill undesirable weeds such as thistle, bindweed, dock and ragweed, use 8 to 10-1/2 ounces in 6 gallons of water (or 2 to 3 teaspoonfuls in 2 quarts of water), and apply directly to plants, wetting them thoroughly. Repeat as necessary. For larger areas use 1 gallon in 50 gallons of water.

To kill undesirable plants such as poison ivy and poison oak, use 8 to 10-1/2 teaspoonfuls in 2 quarts of water or 2 gallons in 12 1/2 gallons of water and drench plants when they are fully leaved. Repeat in about 4 weeks if necessary. Certain hard-to-kill weeds such as poison ivy, poison oak and English daisy may require a Fall application.

**Warm Season Grasses** - St. Augustine, Bahia, common Bermuda, Centipede, Zoysia, also for BENT: Use a one-half rate which is 4 to 5 ounces in 6 gallons of water (2 to 3 teaspoonfuls in 4 quarts of water applied to 200 square feet). Make two applications 10 days apart. Do not spray grasses if stressed from heat, drought, etc. Slight turf yellowing should disappear after about 1 week. Note: Some hybrid Bermuda grasses may be sensitive to this product. Contact your local extension service weed control

BEST AVAILABLE COPY

specialist.

Roadsides (including aprons and guard rails) rights-of-way, drainage ditchbanks and other similar non-crop areas: For control of broadleaf weeds, mix at a rate of 1-1/3 to 4 gallons of Triamine II Lawn Weed Killer per 50 to 300 gallons of water. This mixture will cover one acre. Thoroughly saturate all weeds with spray mixture. Apply when plants come into full leaf (Spring) to the time plants begin to go dormant. Best results are obtained when weeds are young and actively growing. Do not cut weeds until herbicide has translocated throughout the plant causing root death. For small broadleaf weeds use the lower rate. Heavy, dense stands require the higher rate with high water volume.

**For Spot Treatment:** Put 3 teaspoons Triamine II Lawn Weed Killer into an empty quart container with trigger sprayer. Then add 30 oz. tap water. Adjust spray nozzle to give coarse spray. Aim at center of weed and spray to wet. Applicator Cap - Hold applicator end about 6 inches from surface to be treated and thoroughly wet. To obtain wider coverage raise container. One application should be sufficient. Effects begin to show after a few days and weeds gradually die. A repeat application may be required in 3-4 weeks for hard-to-control weeds.

**STORAGE AND DISPOSAL**

**STORAGE:** Always use original container to store pesticides in a secured warehouse or storage building. Do not store near seeds, fertilizers, insecticides or fungicides. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well-ventilated area. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not contaminate water, food, or feed by storage or disposal.

**PESTICIDE DISPOSAL:** If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic containers are also disposable by incineration, or if allowed by state and local authorities by burning. If burned, stay out of smoke.

**WARRANTY**

Riverdale warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant, and soil conditions, mixtures with other chemicals, and factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

(PR933 042194)

