

<b>US ENVIRONMENTAL PROTECTION AGENCY</b> <b>OFFICE OF PESTICIDES PROGRAMS</b> <b>REGISTRATION DIVISION (75-767)</b> <b>WASHINGTON, DC 20460</b>	EPA REGISTRATION 228-218	DATE OF ISSUANCE JUN 11 1987
	TERM OF ISSUANCE Conditional	
	NAME OF PESTICIDE PRODUCT Riverdale Triamine II Premium 75 Liquid	

**NOTICE OF PESTICIDE:**  REGISTRATION  
 REREGISTRATION  
*(Under the Federal Insecticide, Fungicide,  
and Rodenticide Act, as amended)*

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

Riverdale Chemical Company  
220 East 17th Street  
 Chicago Heights, IL 60411-3699

**NOTE:** Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an endorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

This product is conditionally registered in accordance with FIFRA section 3(c)(7)(a) provided that you

1. Submit/owe all data required for registration/reregistration of your product under FIFRA section 3(c)(5) when the Agency requires all registrants of similar products to submit such data.

2. Make the following changes listed below before you release the product for shipment:

- a. Add the phrase "EPA Registration No. 228-218."
- b. In the Hazards to Humans section, incorporate the following statements:

Causes eye irritation. Harmful if inhaled.

If inhaled: Remove victim to fresh air and medical attention.

3. Submit five (5) copies of your final printed labeling before you release the product for shipment. Refer to the A-79 enclosure for a further description of final printed labeling.

ATTACHMENT IS APPLICABLE

**BEST AVAILABLE COPY**

SIGNATURE OF APPROVING OFFICIAL

JUN 11 1987

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

We note that EPA Registration No. 6305-13 has been transferred to Akzo Zout Chemie Nederland bv, EPA Registration No. 38117-11.

A stamped copy of the label is enclosed for your records.

Richard F. Mountfort *RFM*  
Product Manager (23)  
Fungicide-Herbicide Branch  
Registration Division (TS-767C)

Enclosures

**BEST AVAILABLE COPY**

(Front Panel)

RIVERDALE

TRIAMINE™ II PREMIUM 75 LIQUID

WEED AND FEED

READY-TO-USE.

FOR USE WITH A HOSE-END SPRAYER ON GRASS AND ORNAMENTAL TURF

NO MIXING

TREATS 5,000 to 7,500 SQ. FT.

Read directions for different grasses.

CONTAINS MCPA, MECOPROP AND DICHLORPROP AND A 15-0-0 FERTILIZER

ACC  
with comments  
JUN 11 1987  
Read directions for different grasses.  
280-210

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	1.41%
Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid**	1.39%
Dimethylamine Salt of 2-(2,4-Dichlorophenoxy)propionic Acid***	1.37%

INERT INGREDIENTS:	95.83%
TOTAL	100.00%

Isomer Specific by AOAC Method

*2-Methyl-4-Chlorophenoxyacetic Acid equivalent	1.15%	.11 lbs/gal.
**2-(2-Methyl-4-Chlorophenoxy)propionic Acid equivalent	1.15%	.11 lbs/gal.
***2-(2-Dichlorophenoxy)propionic Acid equivalent	1.15%	.11 lbs/gal.

GUARANTEED ANALYSIS:

Total Nitrogen (N)	15%
15% Urea Nitrogen	

† TRIAMINE - A TRADEMARK OF RIVERDALE CHEMICAL COMPANY

Patent Applied For.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE AND BACK PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 228-

EPA EST. NO. 228-IL-1

— NET CONTENTS: 75 Fl. Oz.

NET WEIGHT: 5.37 Lbs.

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

Chicago Heights, Illinois 60411-3699

BEST AVAILABLE COPY

(Rear Panel)

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin. Wash thoroughly with soap and water after handling.

- IF SWALLOWED: Get medical attention.
- IF IN EYES: Immediately flush eyes with plenty of water.
- IF ON SKIN: Wash with plenty of soap and water.

If irritation develops, get medical attention

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from target areas.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.


READ ENTIRE LABELING. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

- 5,000 Sq. Ft. Coverage for Bahia, Bermuda, Blue Grass, Fescue, Rye, and Zoysia Grasses.
- 7,500 Sq. Ft. Coverage for (spring and fall application only) St. Augustine, Centipede and Bent Grasses. Some temporary injury may occur with these grasses.

Do not exceed recommended rate because damage to lawn may occur. Do not apply to Dichondra, Lippia and Clover lawns. Do not use this product for controlling weeds in vegetable gardens and flower beds or around shrubs or ornamental plants.

WEEDS CONTROLLED

- |                   |              |                 |
|-------------------|--------------|-----------------|
| Black Medic       | Jimson Weed  | Speedwell       |
| Buckhorn          | Knotweed     | Spurge          |
| Buttercup         | Lambsquarter | Spur Weed       |
| Canada Thistle    | Mallow       | Stitchwort      |
| Carpetweed        | Morningglory | Toadflax        |
| Chickweed         | Mustard      | Wild Aster      |
| Chicory           | Oxalis       | Wild Carrot     |
| Clover            | Pennywort    | Wild Garlic     |
| Cocklebur         | Pepperweed   | Wild Geranium   |
| Dandelion         | Pigweed      | Wild Lettuce    |
| Dog Fennel        | Plantain     | Wild Onion      |
| English Daisy     | Poison Ivy   | Wild Radish     |
| Florida Pussley   | Poison Oak   | Wood Sorrel     |
| Frenchweed        | Purslane     | Yarrow, and     |
| Ground Ivy        | Ragweed      | other broadleaf |
| Healall           | Red Clover   | weeds           |
| Heartleaf Drymary | Red Sorrel   |                 |
| Henbit            | Sheep Sorrel |                 |



(Rear Panel cont.)

**WHEN TO USE:** Spray anytime during the growing season when weeds are actively growing (except as noted for St. Augustine, Centipede and Bent Grasses). If lawn is extremely dry, water 1 to 2 days before spraying. Do not apply when temperatures are expected to exceed 85°F., or if rain is expected to occur within 24 hours following application. On new lawns wait until grass has reached a height of 2 inches before spraying. Also wait 3 weeks before putting in new grass seed or stolons. After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. Some hard-to-control broadleaf weeds may require a second treatment in 3-4 weeks.

**DISPOSABLE SPRAYER AND BOTTLE INSTRUCTIONS**

**HOW TO APPLY:** Determine area to be sprayed. Apply at a rate of 75 oz. per 5,000 square feet or 7,500 for spring/fall application on St. Augustine, Centipede or Bent Grasses. The entire contents of this bottle must be used to make the proper recommended application rate. Shake well before applying. Connect spray nozzle to hose. Remove bottle cap and screw bottle into nozzle. Turn on the water at the tap. Place finger over hole on right side of nozzle and the Weed and Feed 15-0-0 will mix automatically at the proper ratio. Point in direction away from face and body. While holding sprayer at waist level walk back and forth at a steady pace. Over-lap slightly for uniform coverage. Do not exceed recommended rate because damage to lawn may occur. Do not water fertilized lawn for at least 24 hours.

Marks on left side panel of bottle indicate square feet covered.

(Left Side Panel)

(Right Side Panel)

**COVERAGE MEASURING GUIDE**  
(Not to Scale)

<u>1000 Sq. Ft.</u>	<u>1875 Sq. Ft.</u>
<u>2000 Sq. Ft.</u>	<u>3750 Sq. Ft.</u>
<u>3000 Sq. Ft.</u>	
	<u>6525 Sq. Ft.</u>
<u>4000 Sq. Ft.</u>	
<u>5000 Sq. Ft.</u>	<u>7500 Sq. Ft.</u>

**STORAGE:** Keep from freezing. To be stored in original container and placed in a locked area inaccessible to children. Do not store near food or feedstuff.

**PESTICIDE DISPOSAL:** Product remaining in original container should be disposed of by securely wrapping original container in several layers of newspaper and discarding in trash.

**CONTAINER DISPOSAL:** Do not reuse container. Rinse thoroughly before discarding in trash.

**NOTICE**

Seller warrants that this product conforms to the ingredient statement of the label. Since the conditions of use, such as weather, compatibility with other chemicals, and conditions of application equipment will vary, seller makes no claims other than those stated on this label.