

PM 23 228-220

1043

Riverdale

1.25% HEXAZINONE LIQUID

READY-TO-USE WEED AND BRUSH KILLER

A non-selective (postemergent) liquid for use on industrial plant sites, highways, railroads, and pipeline rights-of-way, storage areas, parking lots and similar areas.

ACTIVE INGREDIENT:

Hexazinone [3-cyclohexyl-6-(dimethylamino)-1-methyl-1, 3, 5-triazine-2,4(1H, 3H)-dione] 1.25%

INERT INGREDIENTS:

98.75%

TOTAL 100.00%

U.S. Patents 3,902,887 & 3,983,116

ACCEPTED
DEC 1 1989
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under 228-220
EPA Reg. No.

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

EPA REG. NO. 228-220

EPA EST. NO. 228-IL-1

NET CONTENTS GAL.

MANUFACTURED BY
RIVERDALE CHEMICAL COMPANY
GLENWOOD, ILLINOIS 60425-1584

BEST AVAILABLE COPY

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

May irritate eyes, nose, throat and skin. Avoid contact with skin, eyes, or clothing. Avoid breathing spray mist. Remove and wash contaminated clothing before re-use.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. *(washes hands, neck and clothes).*

ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands. Do not contaminate water when disposing of equipment washwaters. Do not apply this product through any type of irrigation system.

IMPORTANT: Injury to or loss of desirable trees or other plants may result from failure to observe the following recommendations. Do not apply or drain or flush equipment on or near desirable trees or other plants, or on areas where their roots may extend, or in locations where the chemical may be washed or moved into contact with their roots. Do not use on lawns, walks, driveways, tennis courts, or similar areas. Prevent drift of spray to desirable plants.

Thoroughly clean all traces of 1.25% Hexazinone Liquid from application equipment immediately after use. Flush tank, pump, hoses and boom with several changes of water after removing nozzle tips and screens (clean these parts separately).

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL** before using.

GENERAL INFORMATION: 1.25% Hexazinone Liquid is to be used as a spray for non-selective weed and brush control on non-cropland areas. It is non-corrosive to equipment. 1.25% Hexazinone Liquid is an effective general herbicide providing both contact and residual control of many weeds. At rates recommended for non-cropland areas, it provides control of many annual and biennial weeds, brush, woody vines, and except for Johnsongrass, is effective for control of most perennial weeds. It is absorbed through the roots and foliage. For brush control, soil application is most effective.

Moisture is required to activate 1.25% Hexazinone Liquid in the soil. Best results are obtained when weeds are less than 2 inches in height, soil is moist at time of application and 1/4 to 1/2 inch of rainfall occurs within 2 weeks after application. Foliar application to weeds is most effective under conditions of high temperature (80°F.), high humidity and good soil moisture. Foliar application when vegetation is dormant or semi-dormant may not be effective.

On herbaceous plants, symptoms usually appear within 2 weeks after application under warm, humid conditions, while 4 to 6 weeks may be required when weather is cool. If rainfall after application is inadequate to activate 1.25% Hexazinone Liquid in the soil, plants may recover from contact effects and continue to grow.

On woody plants, symptoms usually appear within 3 weeks after sufficient rainfall has carried the herbicide into the root zone, during periods of active growth. Defoliation and subsequent refoliation may occur, but susceptible plants are killed. The degree of control and duration of effect will vary with the amount of chemical applied, rainfall, temperature, weed and brush species, soil texture and other conditions.

For broadcast treatments using ground equipment apply 1.25% Hexazinone Liquid as a spray just before or soon after weed emergence. Do not apply to frozen or snow-covered soil. Before spraying, calibrate equipment to uniformly and thoroughly cover the vegetation and soil in a measured area to be treated. Add the proper amount of 1.25% Hexazinone Liquid to a spray tank. A suitable surfactant such as Dupont Surfactant WK may then be added at the rate of 1 quart per 100 gals. of spray to improve wetting and/or contact activity. After thoroughly mixing the spray solution, a suitable defoaming agent such as DOW Corning "Antifoam A" or Hodag "Antifoam F-1" may also be added at the rate of 1 to 5 oz. per 100 gals. of spray should foaming become a problem. Agitate thoroughly to disperse the defoaming agent.

Use a fixed boom sprayer properly calibrated to a constant speed and rate of delivery. For application with a handgun sprayer be sure to obtain thorough coverage. For treating small non-crop areas, a tank-type hand sprayer may be used.

BEST AVAILABLE COPY

DIRECTIONS FOR USE (cont'd)

GENERAL WEED AND BRUSH CONTROL—NON-CROPLAND AREAS: Railroad, Highway, Utility and Pipeline Rights-of-Way, Petroleum Tank Farms, Storage Areas, Industrial Plant Sites, Drainage Ditchbanks and other similar areas.

Use the higher levels of the recommended dosage ranges under the following conditions: on hard-to-kill species; on fine-textured soils (clay, clay loam); on soils containing more than 5% organic matter or carbon; and on coarse textured soils (sand, loamy sand, gravelly soils) where rainfall exceeds 40 inches per year. Use the lower levels of the dosage ranges if the growing season is short.

WEED CONTROL: Apply 50 to 100 gals. per acre for season long control of many annual, biennial and perennial weeds including Bahiagrass*, Bermudagrass*, Bluegrass, Broomsedge, Camphorweed, Canada thistle*, Chickweed, Clovers, Dewberry, Dogfennel*, Fescue*, Fingergrass, Foxtail, Guineagrass, Heath aster, Honeysuckle, Lantana, Marehail*, Natalgrass, Plantain, Prickly lettuce, Ragweed, Smutgrass, Spanish needle, Vaseygrass, Wild blackberry*, and Wild carrot.

For short term control (up to 3 months) of the above weeds, apply 17 to 43 gals. per acre. Application at 17 to 43 gals. per acre also provides short term control of many annual, biennial and perennial weeds including Barnyardgrass, Bindweed*, Bouncing bet*, Broomegrass, Buffalo grass*, Burdock, Cocklebur, Crabgrass, Crown vetch, Curly dock*, Dandelion*, Dogbane*, Fiddleneck, Flaree, Flabane, Goatsbeard vine, Goldenrod, Lespedeza, Milkweed*, M. Nutsedge*, Orchardgrass*, Oxalis, Paragrass, Pigweed, Purslane, Quackgrass*, Ryegrass, Smartweed, Spurge, Trumpet creeper*, Wild oats, Wild parsnip and Wild star thistle.

*Considered hard-to kill.

For small area—1½ gals. per 1000 sq. ft. is approximately 66 gals. per acre.

BRUSH CONTROL: Apply to soil during the period between late winter and early summer for control of Blackgum, Black cherry, Elm, Flowering dogwood, Green ash, Hawthorn, Hickory, Oaks (except liveoak), Persimmon, Sumac, Sweetgum, and Willow. In areas where the soil normally remains frozen during the winter and spring rains are usually inadequate for soil activation, a fall or winter treatment may be applied before the soil freezes.

Note: For effective brush control and prevention of damage to desirable vegetation: do not apply to brush standing in water; do not use water from treated ditches for irrigation; do not use where desirable trees or shrubs are immediately adjacent to the treated area.

Broadcast Treatment: Apply 33 to 66 gals. of 1.25% Hexazinone Liquid per acre as a coarse spray. Direct the spray to the soil beneath woody plants to be controlled.

Basal Soil Treatment: Apply 1.25% Hexazinone Liquid with a handgun applicator. This equipment delivers a thin stream of predetermined volume. Apply at the rate of 2 to 3 gals. per 1000 sq. ft. Direct the spray as a straight stream to the soil area surrounding the root collar area of the woody plants to be controlled.

STORAGE AND DISPOSAL

STORAGE: Always store pesticides in a secured warehouse or storage building. Do not store near seed, fertilizers, insecticides or fungicides. Do not contaminate water, food, or feed by storage, disposal or cleaning of equipment.

PESTICIDE DISPOSAL: Open dumping is prohibited. Waste or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state or local procedures. Place in a closed, labeled container for proper disposal.

CONTAINER DISPOSAL: Triple rinse (or equivalent) the container. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by State and local authorities, by burning. If burned stay out of smoke.

NOTICE OF WARRANTY

Seller warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with the use of this product. Ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of seller. In no case shall seller be liable for consequential, special or indirect damage resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. SELLER MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

BEST AVAILABLE COPY