

RIVERDALE

25% DACTHAL DUST

For FLOWERS and VEGETABLES

Use as a pre-emergence weed killer.

ACTIVE INGREDIENT:

DCPA (dimethyl tetrachloroterephthalate) 25%

INERT INGREDIENTS: 75%

TOTAL 100%

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS


CONTENTS 12 OUNCES

EPA Reg. No. 228-222

EPA Est. No. 228-IL-1

Manufactured By
RIVERDALE CHEMICAL COMPANY
Glenwood, IL 60425-1584

(Back Panel)



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Avoid contact with eyes or clothing. Always wash hands, face and arms thoroughly with soap and water after handling and before smoking, eating, drinking or toileting.

IF IN EYES: Flush with plenty of water. Call a physician if irritation persists.

IF ON SKIN: Wash off with soap and water.

USE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE CLOTHING: Wear long-sleeved shirt and long-legged pants and chemical resistant gloves.

AFTER USE: Before removing gloves, wash them with soap and water. Take off all work clothes and shoes. Shower using soap and water. Do not reuse contaminated clothing. Clothing worn during use must be laundered separately from household articles.



ENVIRONMENTAL HAZARDS

Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters.

Groundwater Advisory: Tetrachloroterephthalic acid, a breakdown product of Dacthal, is known to leach through soil as a result of agricultural and turf uses and has been found in groundwater which may be used for drinking water. Users are advised not to apply Dacthal to sand and loamy sand soils where the water table (groundwater) is close to the surface and where these soils are very permeable, i.e. well drained. Your local agricultural agencies can provide further information on the type of soil in your area and the location of groundwater used for drinking water.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Cultivate soil to provide weed free and uniform surface. Do not disturb surface after applying 25% Dacthal Dust.

STRAWBERRIES: For new plantings, apply five tablespoons of 25% Dacthal Dust as a broadcast soil application in sufficient water to cover 100 sq. ft. at transplanting. For established plantings apply three and one-half to five tablespoons as a broadcast application in sufficient water to cover 100 sq. ft. in the fall or early spring. Do not apply after first bloom.

VEGETABLES: SEEDED - Broccoli, Cabbage, Cauliflower, Onions, Snap Beans; TRANSPLANTED - Broccoli, Cabbage, Cauliflower, Onions, Sweet Potatoes, Tomatoes, Strawberries.

CONTROLS THESE WEEDS: Carpetweed, Common chickweed, Purslane, Hairy crabgrass, Goose grass, Green foxtail, Yellow foxtail, Spotted spurge, Prostrate spurge, Spiny pigweed, Dodder and others.

VEGETABLE CROPS: Apply five tablespoons of 25% Dacthal Dust in sufficient water to cover 100 square feet. Application of 25% Dacthal Dust should be made to the soil surface along with or directly following seeding. When transplanting, apply either immediately after transplanting or some time prior to the germination and emergence of susceptible weed species. For tomatoes spray overall four to six weeks after transplanting to weed-free soil.

LAWNS: Apply to established lawns at 6 tablespoons in sufficient water to cover 100 sq. ft. before germination of weed, grass seeds. Sprinkle thoroughly to carry 25% Dacthal Dust into the soil. Do not apply to newly seeded grass.

FLOWERS: Apply uniformly at the rate of five to six tablespoons in sufficient water to cover 100 sq. ft. Apply overall at time of transplanting for new plantings and in the fall or early spring for established plantings.

25% Dacthal Dust is specifically recommended for weed control with the following: Azalea, Barberry, Euonymus, Forsythia, Japanese andromeda, Juniper, Lantana, Pachistima, Pine, Rose, Tree peony, Variegated privet, Wilson ivy, Yew, Ageratum, Alyssum, Aster, Babysbreath, Bleeding heart, Chrysanthemum, Candles larkspur, Coleus, Columbine, Coral

bells, Dahlia, Evening primrose, Fern leaf yarrow, Feverfew, Forget-me-not, Gladiolus, Golden Marguerite, Golden tuft, Lupine, Marigold, Mourning bride, Mourning pinks, Petunia, Poker plant, Purple coneflower, Rose moss, Sahrisk scarlet sage, Snapdragon, Verbena, Zinnia.

DO NOT USE ON THE FOLLOWING PLANTS: Bugle-weed, Button pink, Carnation, Geum, Germande mesembryanthemum, Pansy, Phlox, Sweet William, Telanthera, Violet.

STORAGE AND DISPOSAL

STORAGE: Store in original container and place in a cool, dry location in an area inaccessible to children. Keep container tightly sealed when not in use.

PESTICIDE DISPOSAL; Partially filled containers may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse container. Discard container in trash.

NOTICE: Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith. (RS 060689/FV 022692)