

PM23

228-223

TELEPHONE (800) 345-3330  
(708) 754-3330  
FAX (708) 754-0314

145



OFFICE: 425 WEST 194th STREET, GLENWOOD, IL 60425-1584  
PLANT: 220 EAST 17th STREET, CHICAGO HEIGHTS, IL 60411-3699

February 22, 1993

### CERTIFICATION

I, being an authorized representative of Riverdale Chemical Company certify that all containers of Riverdale Triamine<sup>TM</sup> Granular Weed Killer, EPA Reg. No. 228-223 produced by June 15, 1994, will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products. I further certify this company by January 1, 1995, will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products.

Sincerely,

RIVERDALE CHEMICAL COMPANY

*James K. Champion*  
James K. Champion  
President

JKC:ls

**CPDA** Chemical Producers  
and Distributors  
Association



RIVERDALE

TRIAMINE™

GRANULAR WEED KILLER

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE WHICH CONTROLS  
A WIDE SPECTRUM OF BROADLEAF WEEDS.

Weeds Controlled by this product gradually die and  
disappear from the lawn.

CONTAINS 2,4-D, MECOPROP AND DICHLORPROP.

Treats (or covers) 5,000 Sq. Ft. (50' x 100')

ACTIVE INGREDIENTS:†

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	0.425%
Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid**	0.428%
Dimethylamine Salt of 2-(2,4-Dichlorophenoxy)propionic Acid***	0.423%
INERT INGREDIENTS:	98.724%
TOTAL	100.000%

ISOMER SPECIFIC AOAC METHOD, EQUIVALENT TO:

* 2,4-Dichlorophenoxyacetic Acid	0.35%
** 2-(2-Methyl-4-Chlorophenoxy)propionic Acid	0.35%
*** 2-(2,4-Dichlorophenoxy)propionic Acid	0.35%

†TRIAMINE™ - Trademark of Riverdale Chemical Company

Patent Pending

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS  
AND STATEMENT OF PRACTICAL TREATMENT

NET WEIGHT 20 LBS.


EPA REG. NO. 228-223

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584



(Back Panel)

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Avoid inhalation of dust and contact with eyes, skin, or clothing. Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

#### STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Get Medical Attention.

IF IN EYES: Flush Eyes with Water for 15 Minutes. Get medical attention.

IF ON SKIN: Wash skin with Soap and Water.

#### ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

TRIAMINE<sup>TM</sup> GRANULAR WEED KILLER is a stable mixture of three herbicides. It is for use on Lawns, Ornamental Turf, Parks, also Golf Fairways, Aprons, Tees and Roughs.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to control many broadleaf weeds in lawns; such as:

Alders, Artichoke, Aster, Austrian fieldcress, Beggarticks, Bidden, Bindweed, Bitterweed, Bitter winter cress, Black medic, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckhorn, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common sowthistle, Creeping jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hawkweed, Healall, Heartloaf, drymary, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Indiana

4 of 5

mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarters, Locoweed, Lupines, Mallow, Marshelder, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Nutgrass, Orange hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, Sagebrush, St. Johnswort, Sheep sorrel, Shepherdspurse, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedles, Spatterdock, Speedwell, Spurge, Spurweed, Stinkweed, Stitchwort, Sumac, Sunflower, Sweet clover, Tarweed, Thistle, Toadflax, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Woodsorrel, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds.

**WHEN TO APPLY:** Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. Do not mow 1 to 2 days before and after application. Apply when grass is moist (after a rain, sprinkling or dew). For most effective broadleaf weed killing action, avoid unnecessary disturbances, including watering, of treated areas for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

**NOTE:** When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

**HOW TO APPLY:** This 20 pound bag should be used on 5000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds per 1000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 sq. ft. and adjust to apply at the rate of 1 lb. per 250 sq. ft. Clean spreader thoroughly after use.

#### SPREADER SETTINGS FOR

ONE PASS APPLICATION:°

MAKE

SETTING

Central Calibrated  
Cyclone Broadcast  
Jackson  
Sears Broadcast  
Sears  
Scotts Calibrated

5

3-3/4

5

3

5

4

• • •

• • •

• • •

• • •

• • •

• • •

• • •

• • •

• • •

• • •

• • •

### STORAGE AND DISPOSAL

**STORAGE:** Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near other feed or foodstuffs. To prevent cross-contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides or seeds.

**PRODUCT DISPOSAL:** To dispose of bag which contains unused product, securely wrap original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Do not reuse empty bag. Fold empty bag and discard in trash.

**NOTICE:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions. (ERP 061594)

The following note will not be part of the actual label copy.

\*NOTE: Spreader Settings will vary among various commercial fertilizer manufacturers.