998.993

04/20/2006

01

Please read instructions on reverse before completing form.	<u>. </u>		Form A	pproved. OMB No. 2070-0060
United States Environmental Protection Agency Washington, DC 20460		An An	gistration nendment her	OPP Identifier Number
Applica	tion for Pestici	de - Section I		
1. Company/Product Number 228-223		Product Manager 9 Miller	3.	Proposed Classification
4. Company/Product (Name) Riverdale Triamine Granular Weed Killer	PM# 23	<u>, , , , , , , , , , , , , , , , , , , </u>	[L	
5. Name and Address of Applicant (Include ZIP Code) Nufarm Americas, Inc. 150 Harvester Drive Suite 200 Burr Ridge, IL 60527 Check if this is a new address	(b)(i), rr to: EPA F	ny product is similar (leg. No	or identical in	th FIFRA Section 3(c)(3) composition and labeling
	Section - I	ct Name		
Amendment - Explain below. Resubmission in response to Agency letter dated X Notification - Explain below.		Finel printed labels in Agency letter dated "Me Too" Application Other - Explain below.		NOTIFICATION APR 2 0 2006
Explanation: Use additional page(s) if necessary. (For section I and Section II.) Label notification per PR notice 98-10, Section II and part H. See cover letter for more details. This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or confidential statement of formula of this product. I understand that it is a violation of 18 U.S.C. Sec. 1001 to willfully make any false statements to EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under sections 12 and 14 of FIFRA. Section - III				
1. Material This Product Will Be Packaged In:				. <u></u>
Child-Resistant Packaging Unit Packaging Yes Yes No No Certification must Unit Packaging wgt. No. per Unit Packaging wgt. contained	Water Soluble P Yes No H "Yes" Package wgt	No. per container	Type of Contain Metai Plasti Glass Paper Other	c
3. Location of Net Contents Information 4. Size(s)	l Retail Container	5. Locatio	n of Label Direc	tions
6. Manner in Which Label is Affixed to Product Liti	hograph ber glued inciled	Other		
	Section - IN	1		
1. Contact Point (Complete items directly below for identifica			y, to process ti	his applicatics.
Name George Meindl	Title Registration I	Vanager	Telepho	one No, (include Aree Code)
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete, I acknowledge that any knowlinglly false or misleading statement may be punishable by fine or imprisonment or both under applicable law. (Staraped)				
2. Signifiane Mul	₃. ті⊎∙ Registrati	on Manage	r	
4. Typed Nama George Meindl	^{5. Date} 4/10/20	06		

EPA Form 8570-1 (Rev. 3-94) Previous editions are obsolete.

Clean 4



(Commercial Label)

TRIAMINE[®] GRANULAR WEED KILLER

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS

Weeds controlled by this product gradually die and disappear from the lawn.

CONTAINS 2,4-D, MECOPROP-p AND DICHLORPROP-p.

Treats (or covers) 12,500 Sq. Ft. (100' x 125')

GET THE OPTICAL ADVANTAGE®

ACTIVE INGREDIENTS:

Piverdale®

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid* Dimethylamine Salt of (+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**‡ Dimethylamine Salt of (+)-R-2-(2,4-Dichlorophenoxy)propionic Acid***‡ INERT INGREDIENTS:	0.212% 0.209%
TOTAL:	100.000%
ISOMER SPECIFIC METHOD, EQUIVALENT TO: *2,4-Dichlorophenoxyacetic Acid	. 0.175%

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG, NO. 228-223

NET WEIGHT 50 LBS.

EP4 FST. NO. 228-1L-1

MANUFACTURED BY NUFARM AMERICAS INC., BURR RIDGE, ILLINOIS 60527-0866*

....

20060404.optional

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Avoid inhalation of dust and contact with eyes, skin, or clothing. Wear long pants, long sleeved shirt, socks, shoes and gloves when loading and applying this product. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until dust has settled.

FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

- IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
- IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
- IF ON SKIN Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters, nor pour the washwater on the ground; spray or drain over a large area away from wells and other water sources.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D, MCPP and 2,4-DP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

TRIAMINE[©] GRANULAR WEED KILLER is a stable mixture of three herbicides. It is for use on Ornamental Turf Lawns (Residential, Industrial and Institutional), Parks, Cemeteries, Athletic Fields and Golf Courses (Fairways, Aprons, Tees and Roughs) and similar turf areas.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to control many broadleaf weeds such as: Alder, Artichoke, Aster, Austrian fieldcress, Beggartick, Biden, Bindweed, Bitterweed, Bitter wintercress, Black medic, Blessed thistle, Blue lettuce, Box eider, Broomweed, Buckhorn, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common sowthistle, Creeping jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hawkweed, Healali, Heartleaf drynary, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarter, Locoweed, Lupine, Mallow, Marshelder, Mexicanweed, Morningglory, Muck thistle, Mustard, Nettle, Nutgrass, Orange hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Peppe.weed, Figweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prick'y tettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, Sagebrush, St. Johnswort, Sheep sorrel, Shepherdspurse, Smartweed, Stitchwort, Sumac, Sunflower, Sweet clover, Tarweed, Thistle, Tox4^{nex}, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Wild aster, Wild carret, Wild

garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Woodsorrel, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. Do not mow 1 to 2 days before and after application, Apply when grass is moist (after a rain, sprinkling or dew). For most effective broadleaf weed killing action, avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 50 pound bag should be used on 12,500 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is equal to one application.

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound per 250 square feet. Clean spreader thoroughly after use.

SPREADER SETTINGS **ONE PASS APPLICATION®**

MAKE	<u>SETTING</u>
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Environmental **Benefits** (Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

approximately 20% of the volume in our oxygen for a family of 4 every day nation's landfills. To do your part to vReduces Glare & Reduces Noise reduce this amount, leave grass clippings on your lawn. Increased VTraps Dust, Smoke Particles & Other mowing height and frequency will produce smaller grass clippings that will √Purifies Water & Modifies Temperature decompose more readily so nutrients go back into your lawn.

Grass clippings and yard waste are \sqrt{A} 50' x 50' lawn produces enough

Pollutants

√Improves Soil Condition & Controls Erosion

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near open containers of fertilizer, seed or other pesticide. Do not stack more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticides wastes are toxic. If container is damaged or if pesticide has leaked, clean up all spilled material. Improper disposal of excess pesticide, spray mixtures or rinsate is a violation of Federal law and may contaminate groundwater. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay away from smoke.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions. (RV040406N)


Riverdale, Triamine and The Optical Advantage are registered trademarks of Nufarm Americas Inc.

The following note will not be part of the actual label copy.

PNOTE: Spreader Settings will vary among various commercial fertilizer manufacturers.

OPTIONAL LANGUAGE

- 1. Weed Illustrations.
- 2. Kills dandelions and many other weeds.
- 3. The easy way to kill broadleaf weeds in your lawn.
- 4. Controls a wide spectrum of broadleaf weeds including dandelions, henbit and oxalis.



NOTIFICATION APR 2 0 2006

(Residential Label)

TRIAMINE[®] GRANULAR WEED KILLER

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE WHICH CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS.

Weeds controlled by this product gradually die and disappear from the lawn.

CONTAINS 2,4-D, MECOPROP-p AND DICHLORPROP-p.

Treats (or covers) 5,000 Sq. Ft. (50' x 100')

GET THE OPTICAL ADVANTAGE®

ACTIVE INGREDIENTS:

iverdale [®]

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid* Dimethylamine Salt of (+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**‡	
Dimethylamine Salt of (+)-R-2-(2,4-Dichlorophenoxy)propionic Acid***‡	0.209%
INERT INGREDIENTS:	
	00.00010
ISOMER SPECIFIC METHOD, EQUIVALENT TO:	
*2,4-Dichlorophenoxyacetic Acid	0.350%
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid.	
***(+)-R-2-(2,4-Dichlorophenoxy)propionic Acid ±CONTAINS THE SINGLE ISOMER FORMS OF MECOPROP.p AND DICHLORPROP.p.	0.175%

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE BACK PANEL FOR FIRST AID AND ADDITIONAL PRECAUTIONARY STATEMENTS

EPA REG. NO. 228-223

NET WEIGHT 20 LBS.

EPA ES1, NO. 229-P.-1

MANUFACTURED BY NUFARM AMERICAS INC., BURR RIDGE, ILLINOIS 60527-0866

20060404.optional

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Avoid inhalation of dust and contact with eyes, skin, or clothing. Wear long pants, long-sleeved shirt, socks and shoes when applying this product. After using this product remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until dust has settled.

FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

- IF IN EYES: Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
- IF SWALLOWED: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
- IF ON SKIN Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

TRIAMINE® GRANULAR WEED KILLER is a stable mixture of three herbicides. It is for use on all residential turf.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to control many broadleaf weeds in lawns such as: Alder, Artichoke, Aster, Austrian fieldcress, Becoartick, Biden, Bindweed, Bitterweed, Bitter wintercress, Black medic, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckhorn, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeebeaed, Common sowthistle, Creeping Jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hawkweed, Healall, Heartleaf drymary, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarter, Locoweed, Lupine, Mallow, Marshelder, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Nutgrass, Orange hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce. Primrose. Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, Sagebrush, St. Johnswort, Sheep sorrel, Shepherdspurse, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterucck, Speedwell, Spurce, Spurced, Stinkweed, Stitchwort, Sumac, Sunflower, Sweet clover, Tarweed, Thistle, Toacilax, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild rape, Wild strawberry, Wild s.veet potato, Willow, Witchweed, Woodsorrel, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Avoid contact with desirable shrubs, plants, vegetables, or flowers. Do not reseed for at least 3 to 4 weeks after application. Dc not apply to newly seeded or sodded lawn until after 3 mowings. Do not mow 1 to 2 days before and after application. Apply when grass is moist (after a rain, sprinkling or dew). For most effective broadleaf weed killing action, avoid unnecessary disturbances, including watering of treated areas for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours following treatment. Possible retreatment in approximately

30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 20 pound bag should be used on 5.000 square feet. Apply evenly with a lawn spreader as a one pass application. A two pass application, criss-crossing the area at half the rate each time is equal to one application.

The following settings are approximate for applying contents of this package at a suggested rate of 4 pounds per 1,000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 square feet and adjust to apply at the rate of 1 pound per 250 square feet. Clean spreader thoroughly after use.

SPREADER SETTINGS **ONE PASS APPLICATION[®]**

MAKE	<u>SETTING</u>
Central Calibrated	5
Cyclone Broadcast	3-3/4
Jackson	5
Sears Broadcast	3
Sears	5
Scotts Calibrated	4

Besides increasing property values, did you know a well-maintained lawn Environmental provides the following benefits to your home and environment: **Benefits**

(Grass

isscycling Logo)	approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will	 √Reduces Glare & Reduces Noise √Traps Dust, Smoke Particles & Other Pollutants √Purifies Water & Modifies Temperature
	go back into your lawn.	√Improves Soil Condition & Controls Erosion

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near other feed or foodstuffs. To prevent crosscontamination, do not store near open containers of fertilizer, seed or other pesticide.

DISPOSAL: If empty - Do not reuse this container. Place in trash or offer for recycling if available. If partly filled -If product cannot be used as directed, or offered to others for their use, call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to (RV040406N) label instructions.

Riverdale, Triamine and The Optical Advantage are registered trademarks of Nufarm Americas Inc.

The following note will not be part of the actual label copy.

[®]NOTE: Spreader Settings will vary among various commercial fertilizer manufacturers.

.

OPTIONAL LANGUAGE

1. Weed Illustrations.

r

.

Weed inductions.
 Kills dandelions and many other weeds.
 The easy way to kill broadleaf weeds in your lawn.
 Controls a wide spectrum of broadleaf weeds including dandelions, henbit and oxalis.



Nufarm Americas, Inc. George Meindl Registration Manager 150 Harvester Drive, Suite 200 Burr Ridge, IL 60527 Phone: 630.455-2017 Fax: 630.455.2030 george.meindl@us.nufarm.com

April 10, 2006

Via Overnight Courier

Document Processing Desk (NOTIF) Office of Pesticide Programs (7504C) U. S. Environmental Protection Agency Room 266A, Crystal Mall 2 1801 Bell Street Arlington, VA 22202-4501

Subject: Riverdale® Triamine® Granular Weed Killer EPA Reg. No. 228-223 Label notification per PR98-10

Dear Ms. Hobgood:

Nufarm Americas, Inc would like to add optional language to subject product's label. This proposed language is consistent with PRN98-10 section II (Labeling notifications) and part H (Use of Symbols and Graphics). We have added the following as optional label language:

- 1. Weed illustrations
- 2. Kills dandelions and many other weeds.
- 3. The easy way to kill broadcast weeds in your lawn.
- 4. Controls a wide spectrum of broadleaf weeds including dandelions, henbit and oxalis.

The addition of weed illustrations and the 3 sentences above would not be considered false and misleading.

To process this request please find enclosed the following:

- Application for Pesticide Registration (EPA form 8570-1)
- Revised labeling with areas of change clearly identified (1 copy)
- Revised labeling clean (1 copy)

If you should have any questions regarding this matter, please feel free to contact me at 630.455.2017 or email at <u>george.meindl@us.nufarm.com</u>.

Sincerely, George Meindl

Registration Manager Nufarm Americas, Inc.

H:\228-223\label notification per PRN98-10 "weed illustrations" and optional language submitted 4.10.2006.doc