

RIVERDALE

TRIAMINE™ II

SPOT WEED KILLER

READY TO USE

CONTROLS DANDELIONS, CLOVER, CHICKWEED AND MANY OTHER LISTED WEEDS ON LAWNS AND ORNAMENTAL TURF

ACTIVE INGREDIENTS:†

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid*	. . . . .	.331%
Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid**	. . . . .	.328%
Dimethylamine Salt of 2-(2,4-Dichlorophenoxy)propionic Acid***	. . . . .	.324%
INERT INGREDIENTS . . . . .		<u>99.017%</u>
	TOTAL . . . . .	100.000%

Isomer Specific AOAC Method, Equivalent to:

*2-Methyl-4-Chlorophenoxyacetic Acid . . . . .	.27%	.023 lbs/gal.
**2-(2-Methyl-4-Chlorophenoxy)propionic Acid . . . . .	.27%	.023 lbs/gal.
***2-(2,4-Dichlorophenoxy)propionic Acid . . . . .	.27%	.023 lbs/gal.

†TRIAMINE IS A TRADEMARK OF RIVERDALE CHEMICAL COMPANY

Patent Pending

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

NET CONTENTS 32 FL. OZ. (1 QT.)


EPA REG. NO. 228-224

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584



(Left Side Panel)

## PRECAUTIONARY STATEMENTS

## HAZARDS TO HUMANS &amp; DOMESTIC ANIMALS

## CAUTION

Avoid inhalation of spray mist and contact with eyes or clothing. Causes moderate eye irritation. Wash thoroughly with soap and water after handling.

## STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush with plenty of water. Call a physician if irritation persists.  
IF ON SKIN: Wash skin with plenty of soap and water.

## ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters, nor pour the washwaters on the ground; spray or drain over a large area away from wells and other water sources and desirable plants. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes. If spills occur, collect the material and dispose of by following disposal instructions on this label.

## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

## USE PRECAUTIONS

IMPORTANT: Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray exposed roots of desirable plants. Do not spray on Dichondra, Lippia, nor on grasses where desirable Clovers are present. Do not apply to newly seeded grasses until they have been mowed 3 times. Do not cut grass 1 to 2 days before and after application. Do not reseed for 3 to 4 weeks after use. If lawn needs watering, water thoroughly before application. Do not water for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours. Avoid fine mists. Do not exceed recommended rates because damage to turf may occur. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Do not apply when temperature exceeds 90°F. Shake well before using.

## WEEDS CONTROLLED BY TRIAMINE II SPOT WEED KILLER

Alders, Beggarticks, Bindweed, Black medic, Blue lettuce, Buckhorn, Burcucumber, Burdock, Buttercup, Canada thistle, Carpetweed, Catsear, Chervil, Chickweed, Chicory, Clover, Cocklebur, Croton, Dandelion, Dock, Dogfennel, Dragonhead mint, Dwarf nettle, English daisy, Fanweed, Fat hen, Field bindweed, Florida pusley, Frenchweed, Galinsoga, Goats-beard, Goldenrod, Ground ivy, Halbertleaved salt bush, Hawkweed, Healall, Heartloaf drynary, Hempnettle, Henbit, Hoary cress, Honeysuckle, Jimsonweed, Kochia, Knotweed, Lambsquarters, Mallow, Marshelder, Mexican poppy, Minerslettuce, Moneywort, Morningglory, Mustard, Nutgrass, Oxalis, Pennycress, Pennywort, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Prickly lettuce, Puncture vine, Purslane, Ragweed, Red clover, Red root, Red sorrel, Red stem, Sheep sorrel, Shepherdspurse, Smartweed, Sneezeweed, Sowthistle, Spanishneedles, Speedwell, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Sunflower, Sweet clover, Thornapple, Toadflax, Tree-of-heaven, Velvet leaf, Veronica, Vetch, Violet, Virginia buttonweed, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild gooseberry, Wild jute, Wild lettuce, Wild marigold, Wild petunia, Wild onion, Wild radish, Wild sage, Wild strawberry, Willow, Witchweed, Woodsorrel, Yarrow, Yellow charlock-mustard, Yellow daisy, Yellow rocket, and other broadleaf weeds.

BEST AVAILABLE COPY

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. Some hard-to-control broadleaf weeds may require a second (or an additional) treatment in 3-4 weeks.

**HOW TO USE:** Trigger Spray - Adjust spray nozzle to give coarse spray. Aim at center of weed and spray to wet. Applicator Cap - Hold applicator end about 6 inches from surface to be treated and thoroughly wet. To obtain wider coverage raise container. One application should be sufficient.

**TO PREPARE REFILL:** Add 3 teaspoons (1/2 oz.) Riverdale Triamine II Lawn Weed Killer to this empty quart container. Then add 30 oz. tap water and use according to directions in this label.

#### STORAGE AND DISPOSAL

**STORAGE:** Keep from freezing. To be stored in original container and placed in areas inaccessible to children.

**PESTICIDE DISPOSAL:** Product remaining in original container should be disposed of by securely wrapping original container in several layers of newspaper and discarding in trash.

**CONTAINER DISPOSAL:** Do not reuse empty container except as specified in directions to prepare refill. Discard empty container in trash.

**NOTICE:** Buyer assumes all risks of use, storage and handling this product not in strict accordance with directions given herewith. (PR938 042194/RV 122393)