

RU 23 228-226
1-135

FEB 06 1991

Mr. Russell F. Sawyer
Riverdale Chemical Company
425 West 194th Street
Glenwood, IL 60425-1584

BEST AVAILABLE COPY

Dear Mr. Sawyer:

Subject: Label Amendment- Homeowner label
Sweet Sixteen Weed and Feed with Triamine II
EPA Registration Number 228-219
Your Submission Dated January 22, 1991
Triamine II Weed and Feed
EPA Registration Number 228-225
Your Submission Dated January 28, 1991
3-Way Weed and Feed with Triamine II
EPA Registration Number 228-226 ✓
Your Submission Dated January 29, 1991

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable provided you add the following:

1. MCPA Homeowner Protective Clothing Statements:

"When applying this pesticide, use of a face shield or goggles is required."

- Please note, the Agency is aware of your waiver request to drop this statement. Until the decision on this waiver request is made, the statement is required.

2. Add the method number used in your most recent Isomer Specific AOAC Method testing.

A stamped copy is enclosed for your records. Please submit three (3) copies of your final printed labeling for each product, incorporating this amendment and the above changes.

Sincerely yours,

Joanne I. Miller

CONCURRENCES Acting Product Manager (23)

SYMBOL	H7505C				Fungicide-Herbicide Branch
SURNAME	R. Udins				Registration Division (H7505C)
DATE	Enclosures 2/5/91				

ACCEPTED
with COMMENTS
in EPA Letter Dated

(Homeowner Label)

435

FEB 06 1991

RIVERDALE

EXTRA
COPY

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.
228-226

3-WAY WEED AND FEED

WITH TRIAMINE™ II

THIS PRODUCT IS A PREMIUM GRADE HERBICIDE AND TURF FOOD COMBINATION WHICH
CONTROLS A WIDE SPECTRUM OF BROADLEAF WEEDS WHILE
PROVIDING BOTH FAST AND LONG FEEDING ACTION.

Weeds contacted by this product gradually die and disappear from the lawn.

Treats (or covers) 2,500 Sq. Ft. (25' x 100')

ACTIVE INGREDIENTS:†

Dimethylamine Salt of 2-Methyl-4-Chlorophenoxyacetic Acid* ^{MCPA}	0.159%
Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid** ^{MCPP}	0.157%
Dimethylamine Salt of 2-(2,4-Dichlorophenoxy)propionic Acid*** ^{2,4-DB}	0.155%
INERT INGREDIENTS:	99.529%
TOTAL	100.000%

Isomer Specific AOAC Method, Equivalent to:

*2-Methyl-4-Chlorophenoxyacetic Acid	0.13%
**2-(2-Methyl-4-Chlorophenoxy)propionic Acid	0.13%
***2-(2,4-Dichlorophenoxy)propionic Acid	0.13%

GUARANTEED ANALYSIS:°

Total Nitrogen (N)	XZ
XZ Ammoniacal Nitrogen	
XZ Urea Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	XZ
Soluble Potash (K ₂ O)	XZ

Primary Plant Foods derived from Ammonium Phosphate, Sulfur Coated Urea, Triple Super Phosphate, and Muriate of Potash.

Chlorine (Cl) Maximum	XZ
-----------------------------	----

The urea nitrogen in this product has been coated to provide XZ coated slow release urea nitrogen.

†TRIAMINE™ II - Trademark of Riverdale Chemical Company

Patent Pending

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

NET WEIGHT 20 LBS.

EPA REG. NO. 228-226

EPA EST. NO. 228-1L-1

Revised 1/29/91

BEST AVAILABLE COPY

385

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY
GLENWOOD, ILLINOIS 60425-1584

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes eye irritation. Avoid contact with eyes, skin, or clothing. Harmful if absorbed through skin. Wear long pants, long-sleeved shirt and shoes when applying this product. Wash thoroughly with soap and water after handling. Do not allow persons and pets on treated areas until dusts have settled.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Get medical attention.
IF IN EYES: Flush eyes with water for 15 minutes. Get medical attention.
IF ON SKIN: Wash skin with soap and water.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect non-target plants. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water when disposing of equipment washwaters or rinsate. Do not contaminate water used for irrigation or domestic purposes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

3-WAY WEED AND FEED WITH TRIAMINE II is a stable mixture of three herbicides. It is for use on Lawns and Ornamental Turf.

This product may be used on Bahia, common Bermuda, Blue Grass, Centipede, Fescue, Rye, St. Augustine, and Zoysia grass. DO NOT USE on Dichondra and Carpet grass. DO NOT USE this product where desirable clovers are present or allow the product to "wind drift" onto flowers, vegetables, ornamentals, shrubs, trees or other desirable plants.

This product is designed to promote lush green growth and control many broadleaf weeds in lawns such as:

- Alders, Arrowhead, Beggarsticks, Bindweed, black medic, Blue lettuce, Buckhorn,
- Burcucumber, Burdock, Buttercup, Canada thistle, Carpetweed, Catseye, Chevil,
- Chickweed, Chicory, Clover, Cocklebur, Croton, Dandelion, Dock, Dogfennel, Dragonhead
- mint, Dwarf nettle, English daisy, Fanweed, Fat hen, Field bindweed, Florida pusley,
- Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Halbertleaved salt bush,
- Hawkweed, Healall, Heartloaf drymary, Hempnettle, Henbit, Hoary cress, Honeysuckle,
- Jimsonweed, Kochia, Knotweed, Lambsquarters, Mallow, Marshelder, Mexican poppy,

BEST AVAILABLE COPY

4-15

Minerslettuce, Moneywort, Morningglory, Mustard, Nutgrass, Oxalis, Pennycress, Pennywort, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Prickly lettuce, Puncture vine, Purslane, Ragweed, Red clover, Red root, Red sorrel, Red stem, Sagebrush, Sheep sorrel, Shepherdspurse, Sicklepod, Smartweed, Sneezeweed, Sowthistle, Spanishneedles, Speedwell, Spurge, Spurweed, Stinging nettle, Stinkweed, Stitchwort, Sunflower, Sweet clover, Thornapple, Toadflax, Tree-of-heaven, Velvet leaf, Veronica, Vetch, Violet, Virginia buttonweed, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild gooseberry, Wild jute, Wild lettuce, Wild marigold, Wild petunia, Wild onion, Wild radish, Wild sage, Wild strawberry, Willow, Witchweed, Woodsorrel, Yarrow, Yellow charlock-mustard, Yellow daisy, Yellow rocket, and other broadleaf weeds.

WHEN TO APPLY: Best results are obtained when weeds are growing actively in Spring or early Fall. Apply when grass is moist (after a rain, sprinkling or dew). For most effective broadleaf weed killing action, avoid unnecessary disturbances, including watering, of treated areas for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours following treatment. Avoid contact with desirable shrubs, plants, vegetables, or flowers. Possible retreatment in approximately 30 days may be necessary if lawn is heavily overrun with weeds or if adverse low moisture conditions prevail and weeds are in a state of poor growth. Do not exceed recommended rates because damage to turf may occur. Do not reseed for at least 3 to 4 weeks after application. Do not apply to newly seeded or sodded lawn until after 3 mowings. Do not cut grass 1 to 2 days before and after application.

NOTE: When applying to Southern Grasses - under conditions of inadequate moisture, high temperature and high humidity, over-application may cause minor transitory discoloration on Centipede and St. Augustine grasses. It is recommended that during hot, dry conditions, application rates should be reduced 50% to avoid possible dehydration or browning of grass.

HOW TO APPLY: This 20 pound bag should be used on 2500 square feet. Apply evenly with a lawn spreader as a one pass applicator. A two pass application, criss-crossing the area at half the rate each time is also acceptable.

The following settings are approximate for applying contents of this package at a suggested rate of 8 pounds per 1000 square feet. Settings are based on an average walking speed. Reduce setting for slower speed; raise setting for higher speed. Check your spreader on 250 sq. ft. and adjust to apply at the rate of 2 lb. per 250 sq. ft. Clean spreader thoroughly after use.

SPREADER SETTINGS

ONE PASS APPLICATION: °

MAKE

SETTING

- Cyclone
- Scott's 35
- Scott's C-1
- Central
- K-Mart
- Montgomery Ward's
(1091, 1093, 1095)
- Sears Broadcast

- 6 - 7
- 5 1/2
- J - L
- 7 - 8
- 7 - 8
- 6-3/4
- 6 - 7

BEST AVAILABLE COPY

595

Sweep all fertilizer materials from concrete surfaces immediately after application. Do not wash material from concrete surfaces because fertilizers which contain iron may cause discoloration upon oxidation.

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in locked storage area inaccessible to children and domestic animals. Do not store near other feed or foodstuffs. To prevent cross-contamination, do not store near other fertilizers, or near herbicides, insecticides, fungicides or seeds.

PRODUCT DISPOSAL: To dispose of bag which contains unused product, securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty bag. Fold empty bag and discard in trash.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions. (RS 030191/RV 012991)

The following note will not be part of the actual label copy.

°NOTE: Guaranteed (fertilizer) Analysis and Spreader Settings will vary among various commercial fertilizer manufacturers.

