

RIVERDALE

TRIAMINE™ PREMIUM 3333

LIQUID WEED AND FEED

Ready-To-Use

FOR USE WITH A HOSE-END SPRAYER ON GRASS AND ORNAMENTAL TURF

NO MIXING

Just Connect to Hose, Turn on Water, and Spray

TREATS (COVERS) 3,333 TO 5,000 SQ. FT.
(Read directions for different grasses.)

CONTAINS 2,4-D, MECOPROP, DICHLORPROP AND A X-X-X FERTILIZER

ACTIVE INGREDIENTS:†

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	1.14%
Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid**	1.15%
Dimethylamine Salt of 2-(2,4-Dichlorophenoxy)propionic Acid***	1.13%

INERT INGREDIENTS:	96.58%
TOTAL	100.00%

Isomer Specific AOAC Method No. 6.321, Equivalent to:


*2,4-Dichlorophenoxyacetic Acid95%	.09 lbs./gal.
**2-(2-Methyl-4-Chlorophenoxy)propionic Acid95%	.09 lbs./gal.
***2-(2,4-Dichlorophenoxy)propionic Acid95%	.09 lbs./gal.

GUARANTEED ANALYSIS:°

Total Nitrogen (N)	XZ
XZ Urea Nitrogen	

TRIAMINE IS A TRADEMARK OF RIVERDALE CHEMICAL COMPANY

Patent Pending.



KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE AND BACK PANELS FOR ADDITIONAL PRECAUTIONARY STATEMENTS AND STATEMENT OF PRACTICAL TREATMENT

NET CONTENTS: 1/2 GAL. (64 Fl. Oz.)

NET WEIGHT:° 4.75 LBS.

EPA REG. NO. 228-237

EPA EST. NO. 228-IL-1

Revised A/O 6/12/92

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear long pants, long-sleeved shirt, shoes and chemical resistant gloves when handling or applying this product. Wash nondisposable gloves thoroughly with soap and water before removing. Wash thoroughly with soap and water after handling. Do not allow persons and pets on treated areas until sprays have dried.

STATEMENT OF PRACTICAL TREATMENT

- IF SWALLOWED: Get medical attention.
- IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.
- IF ON SKIN: Wash with plenty of soap and water. If irritation develops, get medical attention.

ENVIRONMENTAL HAZARDS

BEST AVAILABLE COPY

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters nor pour the washwaters on the ground; spray or drain over a large area away from wells, other water sources and desirable plants. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes. If spills occur, collect the material and dispose of by following disposal instructions on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Spray anytime during the growing season when weeds are actively growing (except as noted for S-Southern or Warm Season Grasses).

- 3333 Sq. Ft. Coverage for (N-Northern or Cool Season Grasses) such as Blue Grass, Fescue, and Rye grasses.
- 5000 Sq. Ft. Coverage for (S-Southern or Warm Season Grasses) Bahia,

common Bermuda, St. Augustine, Centipede, Zoysia and Bent grasses. Some temporary injury may occur with these grasses. In Spring, do not apply until grass has reached full green-up.

In Fall, exercise care when grass growth is going into dormancy. When applying to Bent grass, height of grass should be 1-1/2" or more. Note: Some hybrid Bermuda grasses may be sensitive to this product. Contact your local Extension Service weed control specialist.

Designed to control broadleaf weeds and give your lawn a quick boost of important nutrients. Controls dicot weeds during their active growth period. Guaranteed not to burn your lawn when used as directed. High analysis fertilizer featuring a patented nitrogen formula.

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed this product will supplement your regular fertilizer program by providing *(No. of lbs.) pounds of nitrogen per 1,000 square feet (optional P₂O₅ and K₂O levels) and Triamine™ to control listed weeds. Please contact your local extension office for questions regarding a regular fertilizer program.
*Based on Inert fertilizer claimed.

USE PRECAUTIONS

IMPORTANT: Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray exposed roots of desirable plants. Do not spray on Dichondra, Lippia, nor on grasses where desirable Clovers are present. Do not apply to newly seeded grasses until they have been mowed 3 times. Do not cut grass 1 to 2 days before and after application. Do not reseed or plant stolons for 3 to 4 weeks after use. If lawn needs watering, water thoroughly before application. Do not water for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours. Avoid fine mists. Use a lawn type sprayer with a coarse spray as wind drift is less likely. Spray entire lawn, wetting weeds and lawn grasses. Do not exceed recommended rates because damage to turf may occur. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Do not apply when temperature exceeds 85°F. Shake well before using.

WEEDS CONTROLLED BY TRIAMINE™ PREMIUM 3333 LIQUID WEED AND FERTILIZER

Alders, Artichoke, Aster, Austrian fieldcress, Beggarticks, Bident, Bindweed, Bitterweed, Bitter winter cress, Black medic, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckhorn, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common sowthistle, Creeping jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hawkweed, Healall, Heartloaf drymary, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarters, Locoweed, Lupines, Mallow, Marshelder, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Nutgrass, Orange hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed,

BEST AVAILABLE COPY

Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, St. Johnswort, Sheep sorrel, Shepherdspurse, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedles, Spatterdock, Speedwell, Spurge, Spurweed, Stinkweed, Stitchwort, Sumac, Sunflower, Sweet clover, Tarweed, Thistle, Toadflax, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Woodsorrel, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds.

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. Some hard-to-control broadleaf weeds may require a second treatment in 3-4 weeks.

(HOSE-END SPRAYER INSTRUCTIONS)

HOW TO APPLY: Use any 32 oz. hose-end sprayer that is designed for use on lawns and makes a minimum of 15 gallons of finished spray, such sprayers are Ortho's Orthomatic Lawn Sprayer, Ortho's and Black Leaf's Lawn Sprayers (dilution ratio of 60 to 1). The following are adjustable/dial sprayers:

<u>MANUFACTURER</u>	<u>SETTING</u>
Super K-Gro	20
Melnor #94 & #95	11
Melnor #96 & #97	13
Gilmour #362	4
Gilmour #486	12

Special note: Some adjustable sprayers do not hold 32 oz. The above settings are based on using 64 oz. per 3333 square feet on N-Northern or Cool Season grasses or 5000 sq. ft. for application on S-Southern or Warm Season grasses. The entire contents of this bottle must be used to make the proper recommended application rate.

Determine area to be sprayed. Apply at a rate of 64 oz. (1/2 gal.) per 3333 square feet on N-Northern or Cool Season grasses or 5000 for application on S-Southern or Warm Season grasses. Fill sprayer jar with appropriate amount of this product. Do not dilute this product with water when filling sprayer jar. The water will automatically mix with this product during spraying. Attach sprayer to garden hose while holding sprayer at waist level. Point in direction away from face and body. Walk back and forth at a steady pace. Overlap slightly for uniform coverage.

(DISPOSABLE SPRAYER AND BOTTLE INSTRUCTIONS)

HOW TO APPLY: Connect spray nozzle to hose. Remove bottle cap and screw bottle into nozzle. While holding sprayer at waist level, point in direction away from face and body. Turn on water at the tap. Place finger over hole on right side of nozzle (or turn on/off lever to the "ON" position) and this product will mix automatically at the proper ratio. Walk back and forth at steady pace. Overlap slightly for uniform coverage.

(Left Side Panel)

COVERAGE MEASURING GUIDE

Marks on left side panel of bottle indicate square feet covered.
(Not to Scale)

<u>833 Sq. Ft.</u>	<u>1000 Sq. Ft.</u>
<u>1666 Sq. Ft.</u>	<u>2000 Sq. Ft.</u>
<u>2500 Sq. Ft.</u>	<u>3000 Sq. Ft.</u>
	<u>4000 Sq. Ft.</u>
<u>3333 Sq. Ft.</u> Cool Season (N)	<u>5000 Sq. Ft.</u> Warm Season (S)

(Right Side Panel)

STORAGE AND DISPOSAL

STORAGE: Keep from freezing. To be stored in original container and placed in a locked area inaccessible to children. Do not store near feed or foodstuff.

PESTICIDE DISPOSAL: Product remaining in original container should be disposed of by securely wrapping original container in several layers of newspaper and discarding in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Rinse thoroughly before discarding in trash.

NOTICE: Buyer assumes all responsibility for safety and use not in strict accordance with directions and instructions for use. (RS 0300191/RV 061292)

The following will not be part of the actual label copy.
 *Note: Guaranteed (fertilizer) analysis and weight will vary among various commercial fertilizer manufacturers.