

228-237

01/13/2005

(Front Panel)

ACCEPTED

15

RIVERDALE®

TRIAMINE® PREMIUM 3333

LIQUID WEED AND FEED

Ready-To-Use

JAN 13 2005  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

228-237

FOR USE WITH A HOSE-END SPRAYER ON ORNAMENTAL TURF (GRASS) LAWNS

NO MIXING

Just Connect to Hose, Turn on Water, and Spray

TREATS (COVERS) 3,333 TO 5,000 SQ. FT.  
(Read directions for different grasses.)

CONTAINS 2,4-D, MECOPROP-p, DICHLORPROP-p AND A X-X-X FERTILIZER

GET THE OPTICAL ADVANTAGE®

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	1.143%
Dimethylamine Salt of (+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**†	0.574%
Dimethylamine Salt of (+)-R-2-(2,4-Dichlorophenoxy)propionic Acid***†	0.566%
INERT INGREDIENTS:	97.717%
TOTAL	100.00%

Equivalent to:

*2,4-Dichlorophenoxyacetic Acid	0.950%, 0.090 lbs./gal.
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid †	0.475%, 0.045 lbs./gal.
***(+)-R-2-(2,4-Dichlorophenoxy)propionic Acid †	0.475%, 0.045 lbs./gal.

†CONTAINS THE SINGLE ISOMER FORMS OF MECOPROP-p AND DICHLORPROP-p.

GUARANTEED ANALYSIS:°

Total Nitrogen (N)	X%
X% Urea Nitrogen	
Fertilizer guaranteed by (Name of fertilizer registrant)	

Riverdale and Triamine are Registered Trademarks of Nufarm Americas Inc.

KEEP OUT OF REACH OF CHILDREN

CAUTION

SEE SIDE AND BACK PANELS FOR FIRST AID  
AND ADDITIONAL PRECAUTIONARY STATEMENTS

NET CONTENTS: ½ GAL. (64 Fl. Oz.)

NET WEIGHT:° 4.68 LBS.

EPA REG. NO. 228-237

EPA EST. NO. 228-IL-1

MANUFACTURED BY NUFARM AMERICAS INC., BURR RIDGE, ILLINOIS 60527-0866

Revised 3/21/03 Changed First Aid Statement per PR 2001-1, the Disposal Statement per PR 2001-6 and some Environmental hype all via Amendment.

A/O 4/25/03 Changed per EPA's letter dated 2/25/03.

3/15/04 Corrected the Disposal Statement via Notification.

4/29/04 Converted Mecoprop and Dichlorprop to Optical Isomer via Amendment.

8/3/04 Corrected per EPA's letters dated 5/17/04 and 7/15/04.

(Side Panels)

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS****CAUTION**

Harmful if absorbed through skin. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wear long pants, long-sleeved shirt, socks, shoes and chemical resistant gloves when handling or applying this product. After using this product, wash non-disposable gloves thoroughly with soap and water before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

**FIRST AID STATEMENT**

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN  
OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.

**ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters nor pour the washwaters on the ground; spray or drain over a large area away from wells, other water sources and desirable plants. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes. If spills occur, collect the material and dispose of by following disposal instructions on this label.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

Spray anytime during the growing season when weeds are actively growing (except as noted for S-Southern or Warm Season Grasses).

3333 Sq. Ft. Coverage for (-Northern or Cool Season Grasses) such as Blue Grass, Fescue, and Rye grasses.

5000 Sq. Ft. Coverage for (S-Southern or Warm Season Grasses) Bahia, common Bermuda, St. Augustine, Centipede, Zoysia and Bent grasses.

Some temporary injury may occur with these grasses. In Spring, do not apply until grass has reached full green-up.

In Fall, exercise care when grass growth is going into dormancy. When applying to Bent grass, height of grass should be 1-1/2" or more. Note: Some hybrid Bermuda grasses may be sensitive to this product. Contact your local Extension Service weed control specialist.

Designed to control broadleaf weeds and give your lawn a quick boost of important nutrients. Controls dicot weeds during their active growth period. Guaranteed not to burn your lawn when used as directed. High analysis fertilizer featuring a patented nitrogen formula.

A lawn typically will utilize 1 to 4 pounds of nitrogen per 1,000 square feet per growing season. When used as directed, this product will supplement your regular fertilizer program by providing \*(No. of lbs.) pounds of nitrogen per 1,000 square feet (optional P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O levels) and Triamine™ to control listed weeds. Please contact your local extension office for questions regarding a regular fertilizer program.

\*Based on Inert fertilizer claimed.

#### USE PRECAUTIONS

**IMPORTANT:** Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray exposed roots of desirable plants. Do not spray on Dichondra, Lippia, nor on grasses where desirable Clovers are present. Do not apply to newly seeded grasses until they have been mowed 3 times. Do not cut grass 1 to 2 days before and after application. Do not reseed or plant stolons for 3 to 4 weeks after use. If lawn needs watering, water thoroughly before application. Do not water for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours. Avoid fine mists. Use a lawn type sprayer with a coarse spray as wind drift is less likely. Spray entire lawn, wetting weeds and lawn grasses. Do not exceed recommended rates because damage to turf may occur. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Do not apply when temperature exceeds 85°F. Shake well before using.

#### WEEDS CONTROLLED BY TRIAMINE® PREMIUM 3333 LIQUID WEED AND FEED

Alder, Artichoke, Aster, Austrian fieldcress, Beggartick, Bident, Bindweed, Bitterweed, Bitter winter cress, Black medic, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckhorn, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common sowthistle, Creeping jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hawkweed, Healall, Heartleaf drymary, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarters, Locoweed, Lupines, Mallow, Marshelder, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Orange hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, St. Johnswort, Sheep sorrel, Shepherdspurse, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedles, Spatterdock, Speedwell, Spurge, Spurweed, Stinkweed, Stitchwort, Sumac, Sunflower, Sweet clover, Tarweed, Thistle, Toadflax, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Woodsorrel, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds.

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. Some hard-to-control broadleaf weeds may require a second treatment in 3 to 4 weeks.

#### (HOSE-END SPRAYER INSTRUCTIONS)

**HOW TO APPLY:** Use any 32 oz. hose-end sprayer that is designed for use on lawns and makes a minimum of 15 gallons of finished spray, such sprayers are Ortho's Orthomatic Lawn Sprayer, Ortho's and Black Leaf's Lawn Sprayers (dilution ratio of 60 to 1). The following are adjustable/dialsprayers:

<u>MANUFACTURER</u>	<u>SETTING</u>
Super K-Gro	20
Melnor #94 & #95	11
Melnor #96 & #97	13
Gilmour #362	4
Gilmour #486	12

Special note: Some adjustable sprayers do not hold 32 oz. The above settings are based on using 64 oz. per 3333 square feet on -Northern or Cool Season grasses or 5000 sq. ft. for application on S-Southern or Warm Season grasses. The entire contents of this bottle must be used to make the proper recommended application rate.

Determine area to be sprayed. Apply at a rate of 64 oz. (1/2 gal.) per 3333 square feet on -Northern or Cool Season grasses or 5000 for application on S-Southern or Warm Season grasses. Fill sprayer jar with appropriate amount of this product. Do not dilute this product with water when filling sprayer jar. The water will automatically mix with this product during spraying. Attach sprayer to garden hose. While holding sprayer at waist level, point in direction away from face and body. Walk back and forth at a steady pace. Overlap slightly for uniform coverage.

**(DISPOSABLE SPRAYER AND BOTTLE INSTRUCTIONS)**

HOW TO APPLY: Connect spray nozzle to hose. Remove bottle cap and screw bottle into nozzle. While holding sprayer at waist level, point in direction away from face and body. Turn on water at the tap. Place finger over hole on right side of nozzle (or turn on/off lever to the "ON" position) and this product will mix automatically at the proper ratio. Walk back and forth at steady pace. Overlap slightly for uniform coverage.

**COVERAGE MEASURING GUIDE**

Marks on left side panel of bottle indicate square feet covered.  
(Not to Scale)

<u>833 Sq. Ft.</u>	<u>1000 Sq. Ft.</u>
<u>1666 Sq. Ft.</u>	<u>2000 Sq. Ft.</u>
<u>2500 Sq. Ft.</u>	<u>3000 Sq. Ft.</u>
<u>3333 Sq. Ft.</u>	<u>5000 Sq. Ft.</u>
Cool Season (N)	Warm Season (S)

**Environmental Benefits**

(Grasscycling Logo)

Besides increasing property values, did you know a well-maintained lawn provides the following benefits to your home and environment:

*Grass clippings and yard waste are approximately 20% of the volume in our nation's landfills. To do your part to reduce this amount, leave grass clippings on your lawn. Increased mowing height and frequency will produce smaller grass clippings that will decompose more readily so nutrients go back into your lawn.*

- ✓ A 50' x 50' lawn produces enough oxygen for a family of 4 every day
- ✓ Reduces Glare & Reduces Noise
- ✓ Traps Dust, Smoke Particles & other Pollutants
- ✓ Purifies Water & Modifies Temperature
- ✓ Improves Soil Condition & Controls Erosion

**STORAGE AND DISPOSAL**

**STORAGE:** Keep from freezing. To be stored in original container and placed in a locked area inaccessible to children. Do not store near feed or foodstuff.

**DISPOSAL:** If empty - Do not reuse this container, except as a hose-end sprayer bottle only when using Riverdale Triamine Premium 3333 Liquid Weed and Feed. Place in trash or offer for recycling if available. If partly filled - If product cannot be used as directed, call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

**NOTICE**

Buyer assumes all responsibility for safety and use not in strict accordance with directions and instructions for use. (RV 042904C)

The following will not be part of the actual label copy.

Note: Guaranteed (fertilizer) analysis and weight will vary among various commercial fertilizer manufacturers.