

7 of 7

the nearest EPA Regional Office for guidance.


CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by procedures approved by state and local authorities.

Local conditions may affect the use of herbicides. State Agricultural Experiment Station or Extension Service weed specialists in many states issue recommendations to fit local conditions.

WARRANTY

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. The exclusive remedy of user or buyer and the limit of liability of Riverdale Chemical Company is the purchase price paid for the quantity of product involved.

(ERP 102393)



OFFICE: 425 WEST 194th STREET, GLENWOOD, IL 60425-1584
PLANT: 220 EAST 17th STREET, CHICAGO HEIGHTS, IL 60411-3699

April 30, 1993

Ms. Joanne I. Miller
Product Manager (23)
Registration Division - H7505C
U.S. Environmental Protection Agency
Crystal Mall Bldg. #2
1921 Jefferson Davis Highway
Arlington, VA 22202

Dear Ms. Miller:


Subject: 2,4-D Exposure Reduction Program Label Revisions Due 3/1/93
Label Corrections

Following the submission of approximately sixty (60) labels, we conducted an audit and found, for more reasons than we care to go in to, it became necessary to correct the majority of the labels previously submitted.

We have made two copies of each corrected label. We respectfully request that they be entered in to our registration file as replacements. We apologize for any inconvenience this may cause the Agency.


Sincerely yours,

RIVERDALE CHEMICAL COMPANY


By Russell F. Sawyer
Regulatory Affairs Manager

RFS:ls

Enc.



TRIAMINE™ 8000
LAWN WEED KILLER

FOR SELECTIVE BROADLEAF WEED CONTROL IN ORNAMENTAL LAWNS AND TURF GRASSES
And Other Listed Sites

When using a hose-end sprayer, just connect hose,
turn on water and spray.

ONE QUART COVERS UP TO 16,000 SQUARE FEET
(See directions for different grasses.)

ACTIVE INGREDIENTS:†

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	3.6%
Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid**	3.7%
Dimethylamine Salt of 2-(2,4-Dichlorophenoxy)propionic Acid***	3.6%
INERT INGREDIENTS:	89.1%
TOTAL	100.0%

Isomer Specific AOAC Method No. 6.321, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid	3.0%, 0.26 lbs./gal.
**2-(2-Methyl-4-Chlorophenoxy)propionic Acid	3.0%, 0.26 lbs./gal.
***2-(2,4-Dichlorophenoxy)propionic Acid	3.0%, 0.26 lbs./gal.

†TRIAMINE - TRADEMARK OF RIVERDALE CHEMICAL COMPANY

Patent Pending

KEEP OUT OF REACH OF CHILDREN

WARNING

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
AND STATEMENT OF PRACTICAL TREATMENT

NET CONTENTS

EPA REG. NO. 228-239

EPA EST. NO. 228-IL-1

MANUFACTURED BY


RIVERDALE CHEMICAL COMPANY
GLENWOOD, ILLINOIS 60425-1584

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Do not get in eyes. Causes substantial but temporary eye injury. Harmful if swallowed. Avoid contact with skin or clothing. Avoid breathing vapors or spray mist. Wear goggles or face shield, long pants, long-sleeved shirt, socks, shoes and chemical resistant gloves when mixing, loading or applying this product or repairing or cleaning equipment used with this product. For aerial applicators

Revised A/O 1/14/93 Corr.



3 of 12

in an enclosed cockpit and applicators applying this product from a tractor that has a completely enclosed cab, eye protection is not required. Wash hands, face and arms with soap and water as soon as possible after mixing, loading or applying this product. Wash hands, face and arms with soap and water before eating, smoking or drinking. Wash hands, face and arms before using toilet. After work, wash non-disposable gloves thoroughly with soap and water before removing, remove all clothing and shower using soap and water. Do not reuse clothing worn during the previous day's mixing and loading or application of this product without cleaning first. Clothing must be kept and washed separately from other household laundry. Remove saturated clothing as soon as possible and shower. When applying to turf the maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried.

If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron. If this container is five gallons or more in capacity, then a mechanical system (probe and pump) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

STATEMENT OF PRACTICAL TREATMENT

- IF IN EYES: Immediately flush eyes with fresh water for 15 minutes and get medical attention immediately.
- IF ON SKIN: Wash skin with plenty of soap and water. Get medical attention if irritation persists.
- IF SWALLOWED: Drink one (1) or two (2) glasses of water. Induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters nor pour the washwater on the ground; spray or drain over a large area away from wells, other water sources and desirable plants. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes. If spills occur, collect the material and dispose of by following disposal instructions on this label.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D, MCPP and 2,4-DP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

GENERAL INFORMATION

TRIAMINE 8000 LAWN WEED KILLER is a stable solution containing a mixture of three herbicides. It is non-flammable. It is for use on Lawns and Turf (not pastures) and Golf Fairways, Aprons, Tees and Roughs, Parks, Cemeteries, along Highways, Rights-of-Way, Drainage Ditchbanks and other similar non-crop areas.

USE PRECAUTIONS

IMPORTANT: Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray exposed roots of desirable plants. Do not spray on Dichondra, Lippia, nor on grasses where desirable Clovers are present. Do not apply to newly seeded grasses until they have been mowed 3 times. Do not reseed for 3 to 4 weeks after use. If lawn needs watering, water thoroughly before application. Do not water for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours. Avoid fine mists. Use a lawn type sprayer with a coarse spray as wind drift is less likely. Spray entire lawn, wetting weeds and lawn grasses. Do not exceed recommended rates because damage to turf may occur. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Do not apply when temperature exceeds 90°F. Make new dilution for each use. Shake well before using.

WEEDS CONTROLLED BY TRIAMINE 8000 LAWN WEED KILLER

Alders, Artichoke, Aster, Austrian fieldcress, Beggarticks, Bidden, Bindweed, Bitterweed, Bitter winter cress, Black medic, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckhorn, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common sow thistle, Creeping jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hawkweed, Healall, Heartloaf drymary, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarters, Locoweed, Lupines, Mallow, Marshelder, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Nutgrass, Orange hawkweed, Oxalis, Parsnip, Pennygrass, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, St. Johnswort, Sheep sorrel, Shepherdspurse, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedles, Spatterdock, Speedwell, Spurge, Spurweed, Stinkweed, Stitchwort, Sumac, Sunflower, Sweet clover, Tarweed, Thistle, Toadflax, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Woodsorrel, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds.

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. Some hard-to-control broadleaf weeds may require a second (or additional) treatment in 3-4 weeks.

O NTAL LAWNS AND TURF

Cool Season Grasses - such as bluegrass, fescue, and rye: To kill dandelion, wild onion, pennywort, plantain, healall and similar weeds: use 8 ounces in 6 gallons of water and apply with pump or hose-end sprayer or sprinkling can to 2,000 square feet of lawn. For smaller areas - use 3 teaspoons in 2 quarts of water applied to 80 square feet. For larger areas - use 1 gallon in 50 gallons of water to 4/5 acre (35,000 sq. ft.).

5212

To kill undesirable weeds such as thistle, bindweed, dock and ragweed, use 8 ounces in 4-3/4 gallons of water (or 3 teaspoonfuls in 1-2/3 quarts of water), and apply directly to plants, wetting them thoroughly. Repeat as necessary. For larger areas use 1 gallon in 40 gallons of water.

To kill undesirable plants such as poison ivy and poison oak, use 8 teaspoonfuls in 1-2/3 quarts of water or 2 gallons in 10 gallons of water and drench plants when they are fully leaved. Repeat in about 4 weeks if necessary. Certain hard-to-kill weeds such as poison ivy, poison oak and English daisy may require a Fall application.


Warm Season Grasses - St. Augustine, Bahia, common Bermuda, Centipede, Zoysia, also for BENT: Use a one-half rate which is 4 ounces in 6 gallons of water to treat 2000 sq. ft. (3 teaspoonfuls in 4 quarts of water applied to 160 square feet). Make two applications 10 days apart. Do not spray grasses if stressed from heat, drought, etc. Slight turf yellowing should disappear after about 1 week.
 Note: Some hybrid Bermuda grasses may be sensitive to this product. Contact your local extension service weed control specialist.

Roadsides (including aprons and guard rails) rights-of-way, drainage ditchbanks and other similar non-crop areas: For control of broadleaf weeds, mix at a rate of 1 to 3-1/4 pints of TRIAMINE 8000 LAWN WEED KILLER per 5 to 30 gallons of water. This mixture will cover 3500 sq. ft. Thoroughly saturate all weeds with spray mixture. Apply when plants come into full leaf (Spring) to the time plants begin to go dormant. Best results are obtained when weeds are young and actively growing. Do not cut weeds until herbicide has translocated throughout the plant causing root death. For small broadleaf weeds use the lower rate. Heavy, dense stands require the higher rate with high water volume.

For Spot Treatment: Put 4-1/2 teaspoons TRIAMINE 8000 LAWN WEED KILLER into an empty quart container with trigger sprayer. Then add 30 oz. tap water. Adjust spray nozzle to give coarse spray. Aim at center of weed and spray to wet. Applicator Cap - Hold applicator end about 6 inches from surface to be treated and thoroughly wet. To obtain wider coverage raise container. One application should be sufficient. Effects begin to show after a few days and weeds gradually die. A repeat application may be required in 3-4 weeks for hard-to-control weeds.

HOSE-END SPRAYER INSTRUCTIONS:

Use a hose-end sprayer designed for lawns. Put recommended Amount of Lawn Weed Killer (Column 1) in bottle and add water to indicated level (Column 2), and spray on area indicated (Column 3). For example, to spray 2,500 sq. feet measure 10 oz. in jar and add water to the 10-gallon level and mix. Turn on hose and spray as evenly as possible over 2,500 square feet.



Column 1 Amount Lawn Weed Killer to use: (Fl. Oz.)	Column 2 Fill Bottle to: (Gallons)	Column 3 Area to Treat: (Sq. Ft.)
5	5	1,250
10	10	2,500
15	15	3,750
20	20 (Full)	5,000

PRESSURE SPRAYER INSTRUCTIONS:

Measure indicated amount of Lawn Weed Killer into tank and dilute with water according to the following chart. Use a spray pattern which gives small droplets without any fine mist and cover the area indicated.

Amount Lawn Weed Killer to Use: (Fl. Oz.)	Amount of Water in Sprayer (Gallons)	Area to be Sprayed: (Sq. Ft.)
1	1	250
2	2	500
3	3	750
4	4	1,000

DISPOSABLE SPRAYER AND BOTTLE INSTRUCTIONS

Determine area to be sprayed. Apply at a rate of 32 oz. (1 quart) per 8,000 sq. ft. for Cool Season grasses or 16,000 sq. ft. for Warm Season grasses. Connect spray nozzle to hose. Remove bottle cap and screw bottle into nozzle. While holding sprayer at waist level, point in direction away from face and body. Turn on the water at the tap. Place finger over hole on right side of nozzle (or turn on/off lever to the "ON" position) and the Lawn Weed Killer will mix automatically at the proper ratio. Walk back and forth at steady pace.

2000	Sq. Ft.	4000
4000	Sq. Ft.	8000
6000	Sq. Ft.	12000
8000	Sq. Ft.	16000
Cool Season		Warm Season

(Marks on left for 1 qt. disposable bottle. Not to scale.)

STORAGE AND DISPOSAL

STORAGE: Always use original container to store pesticides in a secured warehouse or storage building. Do not store near seeds, fertilizers, insecticides or fungicides. Store at temperatures above 32°F. If allowed to freeze, remix before using. This does not alter this product. Containers should be opened in well-ventilated area. Keep container tightly sealed when not in use. Do not stack cardboard cases more than two pallets high. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Pesticide wastes are toxic. If container is damaged or if pesticide has leaked, contain all spillage. Absorb and clean up all spilled material with granules or sand. Place in a closed labeled container for proper disposal. Improper disposal of excess pesticide, spray mixtures, or rinsate is a violation of Federal law and may contaminate ground water. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for Guidance.


7 of 12

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic containers are also disposable by incineration, if allowed by state and local authorities by burning. If burned, stay out of smoke.

WARRANTY

Riverdale warrants that this herbicide conforms to the chemical description on its label. When used in accordance with label directions under normal conditions, this herbicide is reasonably fit for its intended purposes. Since timing, method of application, weather, plant, and soil conditions, mixtures with other chemicals, and factors affecting the use of this product are beyond our control, no warranty is given concerning the use of this product contrary to label directions or under conditions which are abnormal or not reasonably foreseeable. The user assumes all risks of any such use.

(ERP 102393)



(Non-Commercial Label)

8 of 12

RIVERDALE

TRIAMINE™ 8000
LAWN WEED KILLER

FOR SELECTIVE BROADLEAF WEED CONTROL IN ORNAMENTAL LAWNS AND TURF GRASSES
And Other Listed Sites

When using a hose-end sprayer, just connect hose,
turn on water and spray.

ONE QUART COVERS UP TO 16,000 SQUARE FEET
(See directions for different grasses.)

ACTIVE INGREDIENTS:†

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	3.6%
Dimethylamine Salt of 2-(2-Methyl-4-Chlorophenoxy)propionic Acid**	3.7%
Dimethylamine Salt of 2-(2,4-Dichlorophenoxy)propionic Acid***	3.6%
INERT INGREDIENTS:	89.1%
TOTAL	100.0%

Isomer Specific AOAC Method No. 6.321, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid	3.0%	0.26 lbs./gal.
**2-(2-Methyl-4-Chlorophenoxy)propionic Acid	3.0%	0.26 lbs./gal.
***2-(2,4-Dichlorophenoxy)propionic Acid	3.0%	0.26 lbs./gal.

†TRIAMINE - TRADEMARK OF RIVERDALE CHEMICAL COMPANY

Patent Pending

KEEP OUT OF REACH OF CHILDREN

WARNING

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS
AND STATEMENT OF PRACTICAL TREATMENT

STORAGE AND DISPOSAL

STORAGE: To be stored in original container and placed in area inaccessible to children.

PRODUCT DISPOSAL: Partially filled container may be disposed of by securely wrapping original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Rinse thoroughly before discarding in trash. Do not reuse empty container.

NET CONTENTS


EPA REG. NO. 228-239

EPA EST. NO. 228-IL-1

MANUFACTURED BY

RIVERDALE CHEMICAL COMPANY

GLENWOOD, ILLINOIS 60425-1584



9212

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING

Do not get in eyes. Causes substantial but temporary eye injury. Harmful if swallowed. Avoid contact with skin or clothing. Avoid breathing vapors or spray mist. Wear goggles or face shield, long pants, long-sleeved shirt, socks, shoes and chemical resistant gloves when mixing, loading or applying this product. After using this product, wash nondisposable gloves thoroughly with soap and water before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower. The maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron. If this container is five gallons or more in capacity, then a mechanical system (probe and pump) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

STATEMENT OF PRACTICAL TREATMENT

- IF IN EYES: Immediately flush eyes with fresh water for 15 minutes and get medical attention immediately.
- IF ON SKIN: Wash skin with plenty of soap and water. Get medical attention if irritation persists.
- IF SWALLOWED: Drink one (1) or two (2) glasses of water. Induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters nor pour the washwater on the ground; spray or drain over a large area away from wells, other water sources and desirable plants. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes. If spills occur, collect the material and dispose of by following disposal instructions on this label.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D, MCPP and 2,4-DP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

10 of 12

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

GENERAL INFORMATION

TRIAMINE 8000 LAWN WEED KILLER is a stable solution containing a mixture of three herbicides. It is non-flammable. It is for use on Lawns, Ornamental Turf, Parks, Cemeteries, Golf Fairways, Aprons, Tees, Roughs, also along Highways, Rights-of-Way, Drainage Ditchbanks and other similar non-crop areas.

USE PRECAUTIONS

IMPORTANT: Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray exposed roots of desirable plants. Do not spray on Dichondra, Lippia, nor on grasses where desirable Clovers are present. Do not apply to newly seeded grasses until they have been mowed 3 times. Do not reseed for 3 to 4 weeks after use. If lawn needs watering, water thoroughly before application. Do not water for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours. Avoid fine mists. Use a lawn type sprayer with a coarse spray as wind drift is less likely. Spray entire lawn, wetting weeds and lawn grasses. Do not exceed recommended rates because damage to turf may occur. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Do not apply when temperature exceeds 90°F. Make new dilution for each use. Shake well before using.

WEEDS CONTROLLED BY TRIAMINE 8000 LAWN WEED KILLER

Alders, Artichoke, Aster, Austrian fieldcress, Beggarticks, Biden, Bindweed, Bitterweed, Bitter winter cress, Black medic, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckhorn, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common sow thistle, Creeping jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hawkweed, Healall, Heartloaf drymary, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarters, Locoweed, Lupines, Mallow, Marshelder, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Nutgrass, Orange hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, St. Johnswort, Sheep sorrel, Shepherdspurse, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedles, Spatterdock, Speedwell, Spurge, Spurweed, Stinkweed, Stitchwort, Sumac, Sunflower, Sweet clover, Tarweed, Thistle, Toadflax, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Woodsorrel, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds.

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. Some hard-to-control broadleaf weeds may require a second (or additional) treatment in 3-4 weeks.

11 of 12

ORNAMENTAL LAWNS AND TURF

Cool Season Grasses - such as bluegrass, fescue, and rye: To kill dandelion, wild onion, pennywort, plantain, healall and similar weeds: use 8 ounces in 6 gallons of water and apply with pump or hose-end sprayer or sprinkling can to 2,000 square feet of lawn. For smaller areas - use 3 teaspoons in 2 quarts of water applied to 80 square feet. For larger areas - use 1 gallon in 50 gallons of water to 4/5 acre (35,000 sq. ft.).

To kill undesirable weeds such as thistle, bindweed, dock and ragweed, use 8 ounces in 4-3/4 gallons of water (or 3 teaspoonfuls in 1-2/3 quarts of water), and apply directly to plants, wetting them thoroughly. Repeat as necessary. For larger areas use 1 gallon in 40 gallons of water.

To kill undesirable plants such as poison ivy and poison oak, use 8 teaspoonfuls in 1-2/3 quarts of water or 2 gallons in 10 gallons of water and drench plants when they are fully leaved. Repeat in about 4 weeks if necessary. Certain hard-to-kill weeds such as poison ivy, poison oak and English daisy may require a Fall application.

Warm Season Grasses - St. Augustine, Bahia, common Bermuda, Centipede, Zoysia, also for BENT: Use a one-half rate which is 4 ounces in 6 gallons of water to treat 2000 sq. ft. (3 teaspoonfuls in 4 quarts of water applied to 160 square feet). Make two applications 10 days apart. Do not spray grasses if stressed from heat, drought, etc. Slight turf yellowing should disappear after about 1 week.

Note: Some hybrid Bermuda grasses may be sensitive to this product. Contact your local extension service weed control specialist.

Roadsides (including aprons and guard rails) rights-of-way, drainage ditchbanks and other similar non-crop areas contained within a residential locale: For control of broadleaf weeds, mix at a rate of 1 to 3-1/4 pints of TRIAMINE 8000 LAWN WEED KILLER per 5 to 30 gallons of water. This mixture will cover 3500 sq. ft. Thoroughly saturate all weeds with spray mixture. Apply when plants come into full leaf (Spring) to the time plants begin to go dormant. Best results are obtained when weeds are young and actively growing. Do not cut weeds until herbicide has translocated throughout the plant causing root death. For small broadleaf weeds use the lower rate. Heavy, dense stands require the higher rate with high water volume.

For Spot Treatment: Put 4-1/2 teaspoons TRIAMINE 8000 LAWN WEED KILLER into an empty quart container with trigger sprayer. Then add 30 oz. tap water. Adjust spray nozzle to give coarse spray. Aim at center of weed and spray to wet. Applicator Cap - Hold applicator end about 6 inches from surface to be treated and thoroughly wet. To obtain wider coverage raise container. One application should be sufficient. Effects begin to show after a few days and weeds gradually die. A repeat application may be required in 3-4 weeks for hard-to-control weeds.

HOSE-END SPRAYER INSTRUCTIONS:

Use a hose-end sprayer designed for lawns. Put recommended Amount of Lawn Weed Killer (Column 1) in bottle and add water to indicated level (Column 2), and spray on area indicated (Column 3). For example, to spray 2,500 sq. feet measure 10 oz. in jar and add water to the 10-gallon level and mix. Turn on hose and spray as evenly as possible over 2,500 square feet.

