

228-239

8/4/2003

Page 186 Jacket

AUG 4 2003

Russell F. Sawyer
Nufarm Americas, Inc.
Riverdale - A Nufarm Company
1333 Burr Ridge Parkway, Suite 125A
Burr Ridge, IL 60527-0866

Dear Mr. Sawyer:


SUBJECT: Riverdale Triamine 8000 Lawn Weed Killer
Label Amendment
EPA Registration No. 228-239 ✓
Your Application Dated February 21, 2003

The label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you:

1. Delete the section entitled "Residential Roadsides" (including aprons and guard rails), from the residential label.

The amended label supersede all previously accepted ones. A stamped copy is enclosed for your records. Please submit one copy of your printed label before you release the product for shipment.

Sincerely yours,



Joanne I. Miller
Product Manager (23)
Herbicide Branch
Registration Division (7505C)

Enclosure

(Residential Label)

276

RIVERDALE®

TRIAMINE® 8000 LAWN WEED KILLER

FOR SELECTIVE BROADLEAF WEED CONTROL IN ALL RESIDENTIAL TURF (OR LAWNS)
And Other Listed Sites

When using a hose-end sprayer, just connect hose,
turn on water and spray.

ONE QUART COVERS UP TO 16,000 SQUARE FEET
(See directions for different grasses.)

GET THE OPTICAL ADVANTAGE®

ACTIVE INGREDIENTS:

Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid*	3.6%
Dimethylamine Salt of (+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid**‡	1.8%
Dimethylamine Salt of (+)-R-2-(2,4-Dichlorophenoxy)propionic Acid***‡	1.8%
INERT INGREDIENTS:	92.8%
TOTAL	100.0%

Isomer Specific AOAC Method, Equivalent to:

*2,4-Dichlorophenoxyacetic Acid	3.0%, 0.26 lbs./gal.
**(+)-R-2-(2-Methyl-4-Chlorophenoxy)propionic Acid	1.5%, 0.13 lbs./gal.
***(+)-R-2-(2,4-Dichlorophenoxy)propionic Acid	1.5%, 0.13 lbs./gal.

‡CONTAINS THE SINGLE ISOMER FORMS OF MECOPROP-p AND DICHLORPROP-p.

Riverdale, Triamine and The Optical Advantage are Registered Trademarks of Riverdale - A Nufarm Company

KEEP OUT OF REACH OF CHILDREN

WARNING

SEE BACK PANEL FOR FIRST AID
AND ADDITIONAL PRECAUTIONARY STATEMENTS

NET CONTENTS

EPA REG. NO. 228-239


EPA EST. NO. 228-IL-1

MANUFACTURED BY NUFARM AMERICAS INC., BURR RIDGE, ILLINOIS, 60527-0866 .

ACCEPTED
with COMMENTS
In EPA Letter Dated
AUG 4 2003

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

228-239



Revised 2/21/03 Changed First Aid Statement per PR 2001-1 and Disposal Statement per PR 2001-6. Also added cemeteries and similar non-crop areas via Amendment.
6/13/03 Corrected Ornamental Sites per EPA's Request 5/28/03 Meeting.

PRECAUTIONARY STATEMENTS**HAZARDS TO HUMANS AND DOMESTIC ANIMALS****WARNING**

Causes substantial but temporary eye injury. Harmful if swallowed, inhaled or absorbed through skin. Do not get in eyes or on clothing. Avoid breathing spray mist and contact with skin. Wear goggles or face shield, long pants, long-sleeved shirt, socks, shoes and chemical resistant gloves when mixing, loading or applying this product. After using this product, wash nondisposable gloves thoroughly with soap and water before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower. After product has been diluted in accordance with the "Directions For Use" below, goggles or face shield are not required. When applying to turf, the maximum number of broadcast applications per treatment site is 2 per year. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

If this container is over one gallon and less than five gallons, then persons engaged in open pouring of this product must also wear coveralls or a chemical resistant apron. If this container is five gallons or more in capacity, do not open pour product from this container. A mechanical system (such as a probe and pump or spigot) must be used for transferring the contents of this container. If the contents of a non-refillable pesticide container are emptied, the probe must be rinsed before removal.

FIRST AID STATEMENT

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

- IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
- IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person.
- IF ON SKIN:
OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 to 20 minutes. Call a poison control center or doctor for treatment advice.
- IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.

ENVIRONMENTAL HAZARDS

This product is toxic to aquatic invertebrates. Drift or runoff may adversely affect aquatic invertebrates and non-target plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters nor pour the washwater on the ground; spray or drain over a large area away from wells, other water sources and desirable plants. Do not apply this product through any type of irrigation system. Do not contaminate water used for irrigation or domestic purposes. If spills occur, collect the material and dispose of by following disposal instructions on this label.

Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D, MCPP and 2,4-EP have been associated with mixing/loading and disposal sites. Caution should be exercised when handling these phenoxy pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.

GENERAL INFORMATION

TRIAMINE® 8000 LAWN WEED KILLER is a stable solution containing a mixture of three herbicides. It is non-flammable. It is for use on all Residential Turf.

USE PRECAUTIONS

IMPORTANT: Do not use this product for controlling weeds in flower or vegetable beds or around shrubs or ornamental plantings. Do not spray exposed roots of desirable plants. Do not spray on Dichondra, Lippia, nor on grasses where desirable Clovers are present. Do not apply to newly seeded grasses until they have been mowed 3 times. Do not reseed for 3 to 4 weeks after use. If lawn needs watering, water thoroughly before application. Do not water for 48 hours after application. Watering will wash off weed killing material. Do not apply if rainfall is expected within 48 hours. Avoid fine mists. Use a lawn type sprayer with a coarse spray as wind drift is less likely. Spray entire lawn, wetting weeds and lawn grasses. Do not exceed recommended rates because damage to turf may occur. Spray when air is calm to avoid spray drift that might injure desirable ornamental plants. Do not apply when temperature exceeds 90°F. Make new dilution for each use. Shake well before using.

WEEDS CONTROLLED BY TRIAMINE® 8000 LAWN WEED KILLER

Alder, Artichoke, Aster, Austrian fieldcress, Beggartick, Bideen, Bindweed, Bitterweed, Bitter wintercress, Black medic, Blessed thistle, Blue lettuce, Box elder, Broomweed, Buckhorn, Bull thistle, Burdock, Bur ragweed, Buttercup, Canada thistle, Carpetweed, Catnip, Chickweed, Chicory, Clover, Cockle, Cocklebur, Coffeebean, Coffeeweed, Common sowthistle, Creeping Jenny, Croton, Curly indigo, Dandelion, Dock, Dogbane, Dogfennel, Elderberry, English daisy, Flea bane (daisy), Flixweed, Florida pusley, Frenchweed, Galinsoga, Goatsbeard, Goldenrod, Ground ivy, Gumweed, Hawkweed, Healall, Heartleaf drymary, Hemp, Henbit, Hoary cress, Honeysuckle, Horsetail, Indiana mallow, Ironweed, Jewelweed, Jimsonweed, Kochia, Knotweed, Lambsquarter, Locoweed, Lupine, Mallow, Marshelder, Mexicanweed, Morningglory, Musk thistle, Mustard, Nettle, Nutgrass, Orange hawkweed, Oxalis, Parsnip, Pennycress, Pennywort, Peppergrass, Pepperweed, Pigweed, Plantain, Poison hemlock, Poison ivy, Poison oak, Pokeweed, Poorjoe, Povertyweed, Prickly lettuce, Primrose, Puncture vine, Purslane, Ragweed, Red clover, Red sorrel, Rush, Russian thistle, St. Johnswort, Sheep sorrel, Shepherdspurse, Smartweed, Sneezeweed, Southern wild rose, Sowthistle, Spanishneedle, Spatterdock, Speedwell, Spurge, Spurweed, Stinkweed, Stitchwort, Sumac, Sunflower, Sweet clover, Tarweed, Thistle, Toadflax, Tumbleweed, Velvet leaf, Veronica, Vervain, Vetch, Virginia buttonweed, Virginia creeper, Wild aster, Wild carrot, Wild garlic, Wild geranium, Wild lettuce, Wild onion, Wild radish, Wild rape, Wild strawberry, Wild sweet potato, Willow, Witchweed, Woodsorrel, Wormseed, Yarrow, Yellow rocket, and other broadleaf weeds.

After applying this product, many broadleaf weeds will begin to show visible signs of leaf curl within a few days. Some hard-to-control broadleaf weeds may require a second (or additional) treatment in 3 to 4 weeks.

ORNAMENTAL LAWNS AND TURF

Cool Season Grasses - such as Bluegrass, Fescue, and Rye: To kill Dandelion, Wild onion, Pennywort, Plantain, Healall and similar weeds: use 8 ounces in 6 gallons of water and apply with pump or hose-end sprayer or sprinkling can to 2,000 square feet of lawn. For smaller areas - use 3 teaspoons in 2 quarts of water applied to 80 square feet. For larger areas - use 1 gallon in 50 gallons of water to 4/5 acre (35,000 square feet).

To kill undesirable weeds such as Thistle, Bindweed, Dock and Ragweed, use 8 ounces in 4-3/4 gallons of water (or 3 teaspoons in 1-2/3 quarts of water), and apply directly to plants, wetting them thoroughly. Repeat as necessary. For larger areas, use 1 gallon in 40 gallons of water.

To kill undesirable plants such as Poison ivy and Poison oak, use 8 teaspoons in 1-2/3 quarts of water or 2 gallons in 10 gallons of water and drench plants when they are fully leaved. Repeat in about 4 weeks if necessary. Certain hard-to-kill weeds such as Poison ivy, Poison oak and English daisy may require a Fall application.

Warm Season Grasses - St. Augustine, Bahia, common Bermuda, Centipede, Zoysia, also for Bent: Use a one-half rate which is 4 ounces in 6 gallons of water to treat 2,000 square feet (3 teaspoons in 4 quarts of water applied to 160 square feet). Make two applications 10 days apart. Do not spray grasses if stressed from heat, drought, etc. Slight turf yellowing should disappear after about 1 week.

Note: Some hybrid Bermuda grasses may be sensitive to this product. Contact your local Extension Service weed control specialist.

Residential Roadsides (including aprons and guard rails), rights-of-way, drainage ditchbanks and other similar non-crop areas contained within a residential locale: For control of broadleaf weeds, mix at a rate of 1 to 3-1/4 pints of Triamine 8000 Lawn Weed Killer per 5 to 30 gallons of water. This mixture will cover 3,500 square feet. Thoroughly saturate all weeds with spray mixture. Apply when plants come into full leaf (Spring) to the time plants begin to go dormant. Best results are obtained when weeds are young and actively growing. Do not cut weeds until herbicide has translocated throughout the plant causing root death. For small broadleaf weeds, use the lower rate. Heavy, dense stands require the higher rate with high water volume.

For Spot Treatment: Put 4-1/2 teaspoons of Triamine 8000 Lawn Weed Killer into an empty quart container with trigger sprayer. Then add 30 ounces of tap water. Adjust spray nozzle to give coarse spray. Aim at center of weed and spray to wet. Applicator Cap - Hold applicator end about 6 inches from surface to be treated and thoroughly wet. To obtain wider coverage, raise container. One application should be sufficient. Effects begin to show after a few days and weeds gradually die. A repeat application may be required in 3 to 4 weeks for hard-to-control weeds.

HOSE-END SPRAYER INSTRUCTIONS: Use a hose-end sprayer designed for lawns. Put recommended Amount of Lawn Weed Killer (Column 1) in bottle and add water to indicated level (Column 2), and spray on area indicated (Column 3). For example, to spray 2,500 square feet, measure 10 ounces in jar and add water to the 10-gallon level and mix. Turn on hose and spray as evenly as possible over 2,500 square feet.

Column 1 Amount Lawn Weed Killer To Use: (Fluid Ounces)	Column 2 Fill Bottle To: (Gallons)	Column 3 Area To Treat: (Square Feet)
5	5	1,250
10	10	2,500
15	15	3,750
20	20 (Full)	5,000

PRESSURE SPRAYER INSTRUCTIONS: Measure indicated amount of Lawn Weed Killer into tank and dilute with water according to the following chart. Use a spray pattern which gives small droplets without any fine mist and cover the area indicated.

Amount Lawn Weed Killer To Use: (Fluid Ounces)	Amount of Water In Sprayer (Gallons)	Area To Be Sprayed: (Square Feet)
1	1	250
2	2	500
3	3	750
4	4	1,000

DISPOSAL SPRAYER AND BOTTLE INSTRUCTIONS: Determine area to be sprayed. Apply at a rate of 32 ounces (1 quart) per 8,000 square feet for Cool Season grasses or 16,000 square feet for Warm Season grasses. Connect spray nozzle to hose. Remove bottle cap and screw bottle into nozzle. While holding sprayer at waist level, point in direction away from

face and body. Turn on the water at the tap. Place finger over hole on right side of nozzle (or turn on/off lever to the "ON" position) and the Lawn Weed Killer will mix automatically at the proper ratio. Walk back and forth at steady pace.

2000	Square Feet	4000	(Marks on left for 1 quart disposable bottle. Not to scale.)
4000	Square Feet	8000	
6000	Square Feet	12000	
8000	Square Feet	16000	
Cool Season		Warm Season	

STORAGE AND DISPOSAL

STORAGE: To be stored in original container and placed in area inaccessible to children.

DISPOSAL: If empty - Do not reuse this container. Place in trash or offer for recycling if available. If partly filled - If product cannot be used as directed, or offered to others for their use, call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor (including toilet) or outdoor (including sewer) drain.

NOTICE

Buyer assumes all risks of use, storage and handling of this product not in strict accordance with directions given herewith. (RV 061303)

