

# SUPPLEMENTAL LABELING


READ AND FOLLOW THE ENTIRE LABEL BOOKLET FOR RIVERDALE 2,4-D 6 AMINE BEFORE PROCEEDING WITH THE USE DIRECTIONS CONTAINED IN THIS SUPPLEMENTAL LABELING.

"Label" as used in this supplemental labeling refers to the label booklet for this product and this supplement.



## 2,4-D 6 Amine

EPA Reg. No. 228-242



### Maximum Use Rates For Forestry and Non-Crop Sites

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in any manner inconsistent with its labeling.

This label must be in the possession of the user at the time of application.

All applicable directions and precautions in the label booklet for this product must be followed.

Also refer to additional "GENERAL INFORMATION" and "WEEDS CONTROLLED" in the label booklet. INJURY TO CROPS FROM THIS HERBICIDE MAY OCCUR. IF YOU ARE NOT PREPARED TO ACCEPT SOME DEGREE OF CROP INJURY, DO NOT USE THIS PRODUCT. Product should not be used in or near greenhouses.

#### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR, Part 170. Refer to label booklet under "Agricultural Use Requirements" in the Directions for Use section for information about this standard.

#### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. For Turf use, do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried.

**GENERAL WEED CONTROL** (Airfields, Roadsides, Vacant Lots, Drainage Ditchbanks, Fencerows, Industrial Sites, Rights-of-Way, and similar areas): Use 1-1/3 to 2.75 pints of product per acre. Apply when most annual broadleaf weeds are still young and growing vigorously. Apply when perennial and biennial weeds are actively growing and near the bud stage, but before flowering. For best results on Tansy ragwort and Musk thistle, treat in rosette stage, before bolting. A second application is usually needed for best results on Thistle, Nettle, and Bindweed. Treat Wild onion or garlic in early Spring and in Fall when they are young and growing actively. Mix 2-2/3 pints of this product per acre in adequate water. This may require a spray volume of 300 to 500 gallons of water per acre, depending on the stand. The addition of a wetting agent (spray adjuvant) is suggested. Usually 2-2/3 pints per acre will give adequate control. Do not use on herbaceous ground covers or creeping grass such as Bent. Legumes will usually be damaged or killed. Deep-rooted perennials may require repeat applications. Do not use on freshly seeded turf until grass is well established. Delay reseeding for 30 days.

**WOODY PLANT CONTROL:** To control woody plants susceptible to 2,4-D such as Alder, Buckbrush, Elderberry, Sumac, Cherokee rose, Japanese honeysuckle, Virginia creeper, Wild grape and Willow on non-crop areas such as rights-of-way, fence rows, roadsides and along ditchbanks, use 1-1/3 to 2 quarts of product per acre. This may require a spray volume 30 to 100 gallons of water per acre. Lower volume of water can be used unless applying through such equipment as Directa-Spra, Wobbler, Mini Wobbler, or Spirometer. Spray brush 5 to 8 feet tall after Spring foliage is well developed. Wet all parts of the plants thoroughly, including stem and foliage, to the point of runoff. This may require a spray volume of 300 to 500 gallons of water per acre where the brush is very dense and over 6 to 8 feet high. Spraying can be effective at anytime up to 3 weeks

ACCEPTED

FEB - 7 2006

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

228-242

before frost as long as soil moisture is sufficient for active growth of the brush. Control will be less effective in mid-Summer during hot, dry weather when soil moisture is deficient and plants are not actively growing. A wetting agent may be added to the spray if needed for increased effectiveness. Hard-to-control species may require re-treatment next season. In general, it is better to cut tall woody plants and spray sucker growth when 2 to 4 feet tall.

**Using this product and Tahoe 4E or Tahoe 3A tank mixtures for Non-Crop Areas:** Broadleaf Weed Control: Use 1-1/3 to 2-2/3 pints of this product per acre plus 2 to 6 pints Tahoe 4E (or 3 to 8 pints Tahoe 3A) per acre. For wider spectrum control of broadleaf weeds and woody plants, apply as a broadcast spray in adequate water. This may require 20 to 100 gallons of water per acre. Apply when broadleaf weeds are actively growing. Woody Plant Control Broadcast Foliar Spray: Use 2/3 gallons of this product per acre plus 1-1/2 to 3 quarts Tahoe 4E (or 2 to 4 quarts Tahoe 3A) per acre. Apply as a broadcast spray in adequate water. This may require 20 to 100 gallons of water per acre. Apply when woody plants are actively growing. Woody Plant Control High Volume Leaf-Stem Treatment with Ground Equipment: Use 2/3 to 2 -3/4 quarts of this product per acre plus 1-1/2 to 12 pints Tahoe 4E (or 2 to 16 pints Tahoe 3A) per acre. Mix 2/3 to 2 quarts of this product per acre plus 1-1/2 to 3 pints Tahoe 4 (or 2 to 4 pints Tahoe 3A) in enough water to make 100 gallons of spray. Higher spray volumes of 100 to 400 gallons of water per acre may be required depending on size and density of woody plants. Thoroughly wet all leaves, stems, and root collars of plants to be controlled. Woody Plant Control Aerial Application (Helicopter only): Use 2/3 gallons of this product per acre plus 3 to 4 quarts Tahoe 4E (or 4 to 6 quarts Tahoe 3A) per acre. Apply in adequate water (10 to 30 gallons) using drift control equipment such as Microfoil boom or an effective drift control agent such as Lo-Drift Spray Additive. Use the higher rates and volumes when plants are dense or under drought conditions.

**Using this product and Diablo Herbicide tank mixtures for Non-Crop Areas:** Annual Broadleaf Weeds: Use 1-1/3 to 2-2/3 pints of this Product per acre plus 1/2 to 1-1/2 pints Diablo. For wider spectrum control of broadleaf weeds and woody plants, apply as a broadcast spray in adequate water (20 to 100 gallons water per acre.) Apply when broadleaf weeds are actively growing. Use the higher rates when treating dense or tall vegetative growth. Perennial and Biennial Broadleaf Weeds: Use 2 to 3 pints of this product per acre plus 1/2 to 6 pints Diablo. Apply as a broadcast spray. Apply when broadleaf weeds are actively growing but prior to flowering. Use the lower rates for biennials less than 3 inches rosette diameter. Use the higher rate for perennial weeds or for biennial weeds past the 3 inch rosette stage. Woody Plant Control Broadcast, High Volume, Stem Foliage or Aerial Application: Use 2/3 gallons of this product per acre plus 2 to 8 quarts Diablo. Apply as a broadcast spray in enough water (20 to 100 gallons) or apply as a high volume stem foliage spray in enough volume to thoroughly wet leaves, stems, and root collars (100 to 400 gallons water per acre) or apply aerially in enough water to deliver total spray volume of 10 to 30 gallons water per acre using drift control

equipment such as Microfoil Boom or an effective drift control agent such as Lo-Drift Spray Additive. Use the higher rates and volumes when plants are dense or under drought conditions.

This product will either kill, control or suppress the weeds listed in the label booklet for this product. Some of these species may require repeat spot applications even under ideal conditions.

\* **Maximum Seasonal Application Rate to non-crop and forestry sites is 4 pounds 2,4-D Acid equivalent per acre per application site.**

#### NOTICE

Before buying or using this product, read "WARRANTY LIMITATIONS AND DISCLAIMER" in the label booklet. If terms are not acceptable, return unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by the Buyer or any other User constitutes acceptance of the terms under the "WARRANTY LIMITATIONS AND DISCLAIMER".

**Manufactured By  
Nufarm Americas Inc.  
Burr Ridge, IL 60521**

09/05