			•			
-*:	•	••••		•		

GOOD-HAY **PYRETHRIN CONCENTRATE** FOR FOGGING MACHINE OR DIRECT ANIMAL SPRAY ACTIVE INGREDIENTS

DAIRY

BARNS

CATTLE

Piperonyl Butoxido, Technical** 10.00% Pyrethrins ____ Inert ENGLOWING 57,5-7. NOL IVELUY. *Equivalent to 8% (butylcarbityl)

(6-prophypiperonyl) ether and 27 r lated compounds.

MIX with WATER or OIL

CAUTION: A. 1. 1917

KEEP OUT OF THE REACH OF CHILDREN See Additional Cautions on Back Panel

NET CONTENTS ONE PINT

GOOD-WAY PYRETHRIN-TYPE DAIRY SPRAY CONCENTRATE EMULSIFIABLE

 $Y^{I'}$

For the protection of dairy cattle from biting flies and lice and for control of flies in barns, stables and poultry houses. Protects against hornflies, stableflies, horseflies, houseflies, gnats, mosquitoes, and both blood sucking and biting lice.

Good-Way's Pyrethrin Emulsifiable Concentrate may be used at dilutions of 1 to 39 with water as a contact spray on diary cattle or other animals. When diluted 1 to 9 with water, it provides an excellent fly spray that can be applied in small quantities directly to animals or as a space treatment of barns. Using about a pint to a quart per animal to wet the hair. Projection is then afforded against horse flies, stable flies, horn flies and mosquitoes for several days with partial protection for several additional days.

For treatment of barns and animats with a single application, the concentrate may be diluted with 9 volumes of oil and used in mechanical opraying or foguing machines at one-half ounce per 1000 cubic feet of space,

DIRECTIONS

PROTECTION OF DAIRY ANIMALS

Hornflies, Mosquitoes and Gnats: Dilute one pint of Good-Way Pyrethrin Concentrate with 39 pints of waler and apply to wet the hair thoroughly, with particular attention to topline, underline, flanks, withers, ears and other infested areas. Repeat treatments at intervals of 2 weeks for small populations, or as needed when flies are emerging in big numbers.

Stableflies and Horseflies: Dilute one pint of Good-Way Pyrethrin Concentrate with 9 pints of water and apply a pint or more per head to wet the hair thoroughly, with particular attention to legs, flanks, barrel, topline and other body areas commonly attacked by these flies. Repeat treatments each week, or as needed.

Lice: Different dilutions of Good-Way Pyrethrin Concentrate are recommended for the most effective control of the active stages of biting lice (yellow) on beef and c'airy cattle. For best results from a single treatment, combining kills of young and adults of both biting (yellow) and sucking (blue) lice, a dilution of 1 to 100 is recommended. Apply to wet all portions of the body thoroughly, including the brush of the tail.

Poultry Lice: Using a dilution of 1 to 100 in water. dip birds by submerging in a barrel, tank or tub of the solution, holding the wings away from the body to insure thorough wetting of all infested areas. In order to avoid chilling the fowl, dip them on a warm day or in a warm building.

Horse, Sheep, Goat and Hog Lice: Dilute one pint of concentrate with 100 pints of water for control of motile stages of both sucking and biting lice. Spray the entire animal including the head and the brush of the tail.

Sheep "Tick" or Ked: Ure one pint of Good-Way Pyrethrin Concentrate in 100 pints of water and thoroughly wet all portions of the body by dipping, or by spraying with sufficient pressure and nozzle adjustment to give penetration of the wool. Treat at a rate to wet the animal.

Spinose Ear Tick: Good-Way Pyrethrin Concentrate at I to 24 with water and apply by individual treatment, syringing or pouring 1 to 2 ounces of material into each ear.

CAUTION-Cover and protect all milking utensils and milk from spray. Do not spray into or near an open flame. Do not smoke when spraying. Harmful if swallswed

874 188. 10. 228-247 EPA 857. 10. 238-TL-1

AND A CHINE AND LIVILLALE CHOICAL CONTANT GLERHOOD. 611210018 . 60425-1584/

Not Reviewed. Registrant claims to be in accordance with BEST AVAILABLE COPI Helipolise, Paga 2.12, Active to Inert Change in Status

